

DTIC FILE COPY

ornl

**OAK RIDGE
NATIONAL
LABORATORY**

MARTIN MARIETTA

AD-A224 521

OPERATED BY
MARTIN MARIETTA ENERGY SYSTEMS, INC.
FOR THE UNITED STATES
DEPARTMENT OF ENERGY

AD
ORNL/TM-11573

ARMY PROJECT ORDER NO: 89PP9921

REPORT NO: DOE IAG 1016-B123-A1

TITLE: CHARACTERIZATION OF EXPLOSIVES PROCESSING
WASTE DECOMPOSITION DUE TO COMPOSTING

PRINCIPAL INVESTIGATORS: W. H. Griest, A. J. Stewart, R. L. Tyndall,
C.-h. Ho, and E. Tan

CONTACTING ORGANIZATION: Oak Ridge National Laboratory
P. O. Box 2008, Bldg. 4500S, MS-6120
operated by
Martin Marietta Energy Systems, Inc.
for the
U.S. Department of Energy
under Contract DE-AC05-84OR21400

REPORT DATE: January 31, 1990

TYPE OF REPORT: Final, Phase I Report

PREPARED FOR: U.S. ARMY MEDICAL RESEARCH AND
DEVELOPMENT COMMAND
Fort Detrick, Frederick, Maryland 21701-5012

DISTRIBUTION STATEMENT: Approved for public release;
Distribution unlimited

The findings in this report are not to be construed as an official Department of the
Army position unless otherwise so designated by other authorized documents.

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

DTIC
S ELECTE D
JUL 06 1990
B

90 07 5 1990

This report has been reproduced directly from the best available copy.

Available to DOE and DOE contractors from the Office of Scientific and Technical Information, P.O. Box 62, Oak Ridge, TN 37831, prices available from (615) 576-8401, FTS 626-8401.

Available to the public from the National Technical Information Service, U.S. Department of Commerce, 5285 Port Royal Rd., Springfield, VA 22161.
NTIS price codes—Printed Copy: A09 Microfiche A01

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

**Characterization of Explosives Processing Waste
Decomposition Due to Composting**

FINAL, PHASE I REPORT

January 1990

**W. H. Griest, A. J. Stewart, R. L. Tyndall,
C.-h. Ho, and E. Tan**

DOE Interagency Agreement No. 1016-B123-A1

**Supported by
U.S. Army Medical Research and Development Command
Fort Detrick, Frederick, Maryland 21701-5012**

**Project Officer: Dr. Wayne R. Mitchell
U.S. Army Biomedical Research and Development Laboratory
Fort Detrick, Frederick, Maryland 21701-5010**

**OAK RIDGE NATIONAL LABORATORY
Oak Ridge, Tennessee 37831
operated by
MARTIN MARIETTA ENERGY SYSTEMS, INC.
for the
U.S. DEPARTMENT OF ENERGY
Under Contract No. DE-AC05-84OR21400**

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS NA		
2a. SECURITY CLASSIFICATION AUTHORITY NA			3. DISTRIBUTION/AVAILABILITY OF REPORT APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED.		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE NA					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) ORNL/TM-11573			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION OAK RIDGE NATIONAL LABORATORY		6b. OFFICE SYMBOL (if applicable)		7a. NAME OF MONITORING ORGANIZATION U.S. ARMY BIOMEDICAL RESEARCH AND DEVELOPMENT LABORATORY	
6c. ADDRESS (City, State, and ZIP Code) PO BOX 2008, BLDG 4500S, MS-6120 OAK RIDGE, TENNESSEE 37831-6120			7b. ADDRESS (City, State, and ZIP Code) FORT DETRICK, FREDERICK, MARYLAND 21701-5010		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION U.S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND		8b. OFFICE SYMBOL (if applicable) SGRD-RMI-S		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER PROJECT ORDER NO. 89PP9921	
8c. ADDRESS (City, State, and ZIP Code) FORT DETRICK, FREDERICK, MARYLAND 21701-5012			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO. 62720A	PROJECT NO. 3EI- 62720A835 ✓	TASK NO. 00
11. TITLE (Include Security Classification) CHARACTERIZATION OF EXPLOSIVES PROCESSING WASTE DECOMPOSITION DUE TO COMPOSTING. PHASE I REPORT.					
12. PERSONAL AUTHOR(S) W. H. GRIEST, A. J. STEWART, R. L. TYNDALL, C.-h. HO, E. TAN					
13a. TYPE OF REPORT FINAL, PHASE I		13b. TIME COVERED FROM 05/89 TO 12/89		14. DATE OF REPORT (Year, Month, Day) 90/01/31	
15. PAGE COUNT 182					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) EXPLOSIVES PROCESSING WASTE, EXPLOSIVES CONTAMINATED SOIL, DECONTAMINATION, COMPOSTING, CHARACTERIZATION, CHEMISTRY, TOXICITY RA III		
FIELD	GROUP	SUB-GROUP			
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Explosives-contaminated (primarily TNT, RDX, and HMX) sediments composted under mesophilic and thermophilic conditions at the Louisiana Army Ammunition Plant were subjected to regulatory leaching (Toxicity Characteristic Leaching Procedure and Synthetic Precipitation Leach Test) and organic solvent extraction followed by chemical characterization and toxicity tests. Chemical characterization included EPA Contract Laboratory Program Target compound list volatile and semivolatile organics, PCBs, pesticides, metals, and also explosive compounds and TNT metabolites. The toxicity testing utilized <i>Ceriodaphnia dubia</i> , fathead minnow larvae, and the Ames test. Regulatory compounds were not detected at significant concentrations in the leachates or extracts and the main constituents found were TNT, HMX, RDX, and two monoamino dinitrotoluenes at low to sub-ppm levels. Low toxicity was detected in the aqueous leachates, with the leachate from the mesophilic compost being more toxic. Particles recovered from the leachates had only low extractable toxicity. Considerably higher toxicity was found in the organic solvent extracts, because of lesser dilution than the regulatory leachates. Comparison of toxicity calculated per g of compost extracted or leached did not reveal any substantial toxicity in the compost which was not liberated by the regulatory leaching procedure, but the high dilutions (100g compost per 2L of leachate) considerably reduced the sensitivity of the Ames test unless the leachates were concentrated 100-fold. Considering the lack of a composted blank devoid of explosives (not included in the original compost engineering study design) and the age of the compost samples (ca. 10 months) before characterization these results must be considered as preliminary. The application of Phase I methods and experiences to Phase II studies of fresh samples including proper blanks and controls will provide a more conclusive test of explosives process waste detoxification by composting.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input checked="" type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL MRS. VIRGINIA M. MILLER			22b. TELEPHONE (Include Area Code) (301) 663-7325		22c. OFFICE SYMBOL SGRD-RMI-S

DD Form 1473, JUN 86

Previous editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE

UNCLASSIFIED

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

_____ Where copyrighted material is quoted, permission has been obtained to use such material.

_____ Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

W.H.Z. Citations of commercial organizations and trade names in this report do not constitute an official Department of the Army endorsement or approval of the products or services of these organizations.

W.H.Z. In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Animal Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

_____ For the protection of human subjects, the investigator(s) have adhered to policies of applicable Federal Law 45CFR46.

_____ In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

W.H.Z. Duest 6/5/90
PI Signature Date



Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

EXECUTIVE SUMMARY

Laboratory, pilot scale, and field demonstration studies show that composting can reduce the concentrations of explosives in contaminated soils and sediments, and they suggest that composting has the potential to be a viable option to incineration. However, the final forms of the transformed explosives compounds and the toxicity of the compost product have not been established. Such information is needed to determine if composting can decontaminate explosives-contaminated soils to products which can be safely disposed by surface application or burial. This project addresses these questions. The purpose of Phase 1, reported in this document, was to investigate the chemical and toxicological characteristics of the mesophilic and thermophilic composts produced from explosives-contaminated lagoon sediments at the Louisiana Army Ammunition Plant by Roy F. Weston, Inc. in the spring and summer of 1988.

Samples of the mesophilic and thermophilic composts were subjected to the U. S. Environmental Protection Agency (EPA) Synthetic Precipitation Leach Test (CCLT for "Clean Closure Leach Test") and the Toxicity Characteristic Leaching Procedure (TCLP). The leachates were analyzed for EPA Contract Laboratory Program volatile and semivolatile organics, PCBs and pesticides; several metals on the Target Compound List; and explosive compounds and 2,4,6-trinitrotoluene (TNT) metabolites. The aquatic toxicity and bacterial mutagenicity of the CCLT leachates were determined using *Ceriodaphnia dubia* and fathead minnow larvae, and the Ames test (respectively). Very little of importance was detected in the chemical characterization of the leachates except for low $\mu\text{g/mL}$ concentrations of TNT, hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX), octahydro-1,3,5,7-tetranitro-1,3,5,7-tetrazocine (HMX), and two monoaminodinitrotoluenes. Low toxicity was detected in the CCLT leachates, and mutagenicity was difficult to measure except when the leachate was concentrated 100-fold. The mesophilic compost leachate was slightly more toxic than that of the thermophilic compost. This is consistent with the greater biotransformation of the explosives in the thermophilic compost and the lower concentrations of such compounds found in the leachate, but the lack of a compost blank prevents assigning the cause(s) of the toxicity to explosives. Only minor levels of mutagenic activity were found to be associated with suspended particles which pass through the pressure filter in the EPA leaching protocols. Compost heterogeneity considerably complicated chemical analysis and toxicity testing, and it requires that relatively large masses of sample or large numbers of samples be tested.

Organic solvent extracts were considerably more toxic than the aqueous leachates on the basis of toxicity per mg of extractable matter or per mL of extract, but not on the basis of toxicity per g of compost leached or extracted. The CCLT does not appear to miss large reservoirs of aquatic organism toxicity or bacterial mutagenicity which are available to organic solvent extraction, but the high dilution of the leached material in the CCLT protocol makes determination of mutagenic activity difficult without preconcentration of the CCLT. Although a substantial portion of the bacterial mutagenic activity apparently could be accounted for by the TNT content of the extract, the lack of a blank compost prevents assignment of such activity to explosives compounds or their biotransformation products. Indeed, the relative mutagenic responses with two bacterial strains were different from those of TNT and most of its available metabolites. This observation suggests that compounds other than those identified contribute or affect the extractable mutagenicity. Hydrolysis of the organic solvent-extracted composts to release a labile bound fraction of transformed explosive compound products did not produce detectable toxicity, but the survivability of mutagens to the hydrolysis conditions remains a question.

Although the Phase 1 results showed low toxicity of the aqueous leachates of the composts, the age of these compost samples (ca. 10 months after composting was completed) and the lack of proper controls (such as the original sediment used for composting and a blank compost from uncontaminated sediment) for a toxicity investigation prevent definitive conclusions from being made regarding the detoxification of explosives-contaminated soils. The techniques developed and lessons learned from the Phase 1 studies will allow a more conclusive assessment to be made in Phase 2 with compost samples to be produced in a second field composting experiment scheduled for the spring of 1990.

TABLE OF CONTENTS

FOREWORD	vii
EXECUTIVE SUMMARY	1
TABLE OF CONTENTS	3
LIST OF TABLES	5
LIST OF FIGURES	7
1. INTRODUCTION	9
2. COMPOST SAMPLE PREPARATION AND CHEMICAL CHARACTERIZATION	10
2.1 Compost Acquisition and Sampling	10
2.2. Preparation and Chemical Characterization of Regulatory Leachates	10
2.2.1. Preliminary Trial of EPA Synthetic Precipitation Leach Test	10
2.2.2. Final Trial of EPA Synthetic Precipitation Leach Test	11
2.2.3. EPA Toxicity Characteristic Leaching Procedure	11
2.2.4. Chemical Characterization of Regulatory Leachates	13
2.2.5. Leachate and Compost Homogeneity	15
2.3. Preparation and Characterization of Organic Solvent Extracts	18
2.3.1. Comparison of Organic Solvents for Compost Extraction	18
2.3.2. Comparison of Organic Solvent Extraction Procedures	18
2.3.3. Preparation of Organic Solvent Extracts	21
2.3.4. Chemical Characterization of Organic Solvent Extracts	21
2.4. Preparation and Chemical Characterization of Hydrolyzed Bound Fraction	24
2.5. Preparation of Particle-Bound Organics Extract from CCLT Leachate	26
2.6. Analysis of Mesophilic Compost Wood Chips for Explosives and TNT Metabolites	26
2.7. Conclusions	27

TABLE OF CONTENTS (Cont'd)

3.	AQUATIC TOXICITY TESTS	28
3.1.	Introduction	28
3.2.	Methods	28
3.3.	Observations	29
3.3.1.	Tests of Preliminary CCLT Leachates	29
3.3.2.	Tests of Final CCLT Leachates	29
3.3.3.	Tests of Preliminary Acetonitrile Extracts	34
3.3.4.	Tests of Final Organic Solvent Extracts	36
3.3.5.	Tests of the Hydrolyzed Bound Fraction from Composts	36
3.4.	Conclusions	37
4.	AMES TESTS	39
4.1.	Introduction	39
4.2.	Materials and Methods	39
4.3.	Observations	40
4.3.1.	Confirmation of Strains	40
4.3.2.	Tests of Explosives Compounds and TNT Metabolites	40
4.3.3.	Tests of CCLT Leachates and Suspended Particles from Leachates	41
4.3.4.	Tests of Organic Solvent Extracts	44
4.3.5.	Tests of Hydrolyzed Bound Fraction	45
4.4.	Discussion and Conclusions	45
5.	CONCLUSIONS	48
	ACKNOWLEDGEMENTS	48
	REFERENCES	49
	APPENDIX	51
	DISTRIBUTION LIST	183

LIST OF TABLES

<u>Table</u>		
2-1	Recovery of Explosive Compounds and TNT Metabolites from Prefiltration of TCLP Leachates from Composts	12
2-2	Comparison of Gross Characteristics of CCLT and TCLP Leachates of Composts	13
2-3	Results of Explosives Compounds and TNT Metabolites Analysis of Compost Leachates	14
2-4	Variation in RDX and TNT Concentrations in Individual 4L Portions of Thermophilic Compost CCLT Leachates	16
2-5	Examination of Homogeneity of Compost Samples Used for Solvent Comparison Study	17
2-6	Comparison of Organic Solvents for Mesophilic Compost Extraction ...	19
2-7	Comparison of Solvent Extraction Methods for Composts	20
2-8	Parameters of Organic Solvent Extracts for Chemical and Toxicity Testing	22
2-9	Explosive Compound and TNT Metabolite Analysis of Organic Solvent Extracts and Hydrolyzed Bound Fraction	23
2-10	Parameters for Preparation of Hydrolyzed Bound Fractions for Chemical and Toxicity Testing	26
2-11	Characterization of Mesophilic Compost Wood Chips for Explosives and Biotransformation Products	27
3-1	Results of <u>Ceriodaphnia</u> Toxicity Tests of Preliminary CCLT Leachates in 2.5 mL (Upper) and 17-mL (Lower) Test Chambers	30
3-2	Results of <u>Ceriodaphnia</u> and Fathead Minnow Larvae Toxicity Tests of CCLT Leachates Prepared from Thermophilic and Mesophilic Composts	31
3-3	Summary of Results for Chemical Analyses of Control Water, Synthetic Rainwater, and Mesophilic and Thermophilic Leachates Performed in Conjunction with Toxicity Test	32
3-4	<u>Ceriodaphnia</u> Survival and Fecundity in Various Concentrations of Acetonitrile Extracts of Thermophilic (T) or Mesophilic (M) Composts Contaminated with Explosives	33

LIST OF TABLES - (Cont'd)

Table

3-5	Results of Ceriodaphnia Toxicity Tests of Pure Dimethylsulfoxide (DMSO), Acetonitrile Extracts of Thermophilic (T-A) and Mesophilic Compost, and Ethyl Acetate Extracts of Thermophilic(T-E) and Mesophilic (M-E) Compost	33
3-6	Results of Fathead Minnow Toxicity Tests of Acetonitrile Extracts of Mesophilic (M-A) and Thermophilic (T-A) Composts	34
3-7	Survival and Fecundity of Ceriodaphnia Tested with the Acid-Hydrolyzed Extracts of Thermophilic (T) and Mesophilic (M) Compost	35
3-8	Estimated Toxicity (toxic units/gram) of Thermophilic and Mesophilic Compost Based on Results of <u>Ceriodaphnia</u> Tests of Extracted Material Obtained Using Aqueous (CCLT) and Organic-Solvent (Acetonitrile and Ethyl Acetate) Extractants	38
4-1	Results from Ames Testing of Known Mutagens	41
4-2	Summary of Ames Mutagenicity Test for Explosives and Metabolites ...	42
4-3	Summary of Results for Ames Mutagenicity Test of the DMSO Concentrate of the Mesophilic and Thermophilic Compost CCLT Leachates	43
4-4	Ames Test of Mesophilic CCLT Leachates, Before and After Centrifugation, and Leachate Solids Extracted with DMSO	44
4-5	Ames Test of Preliminary Acetonitrile Extracts of Compost	45
4-6	Ames Test Result Summary for Final Organic Solvent Extracts of Composts	46
4-7	Comparison of Mutagenic Activities	47

LIST OF FIGURES

Figure

- 2-1 Capillary Column GC of Crude and Derivatized Compost Extracts. Elution of TNT is Labeled. Brackets Show Elution Range of Metabolites (crude) and TNT Plus Metabolites (TMS-Derivatized) 25

CHAPTER 1. INTRODUCTION

The environmental restoration of explosive compound-contaminated soils and sediments at U. S. Army installations will be a very expensive and time-consuming process (1). Decontamination of approximately 28 installations will require the treatment of at least 1,000,000 cubic yards of contaminated soil (total). Incineration is the only decontamination technology available at the present, but at a cost of \$200 to \$300 per ton, tens of millions of dollars will be required per site. Two existing incinerators each can treat less than one ton of soil per hour. Alternative technologies are clearly needed to reduce costs and time.

Composting is an attractive alternate to incineration, and laboratory (2), pilot scale (3), and field scale studies (4) have demonstrated that the concentrations of explosives in soil or sediment can be effectively reduced by biotransformation. The field composting experiment conducted at the Louisiana Army Ammunition Plant (LAAP) by Roy F. Weston, Inc. (4) reduced the concentrations of TNT, RDX, and HMX (see Appendix A-1 for list of abbreviations) in lagoon sediments by 99.9, 99.1, and 96.5% (respectively) in 153 days of composting under thermophilic conditions. Under mesophilic composting conditions, the percentage reductions were slightly lower (99.6, 94.8, and 86.9%, respectively).

Major questions to be answered include the ultimate fate of the biotransformed explosives and the toxicity of the compost product. This project has two objective, which address these questions. The first objective is to determine the chemical and toxicological characteristics of the two composts produced in the Roy F. Weston, Inc. ("Weston") experiments at the LAAP (4). This characterization is the subject of this Phase 1 report. The second objective, to be addressed in Phase 2 of this project, focuses on more fundamental issues associated with composting munitions wastes. These issues include identifying and determining toxic major biotransformation products of the explosives compounds, and determining the potential for long-term releases of organically-bound explosive compound products formed during composting. Phase 2 also provides for the chemical and toxicological characterization of materials from a second, more extensive field demonstration experiment planned for the spring of 1990 by Weston. This experiment includes the proper controls and samples for a more conclusive toxicity determination.

It must be strongly emphasized that the results presented in this report must be considered only as preliminary, and more of a methods development effort. This is because the Weston experiment was designed only from engineering considerations, and this Phase 1 chemical characterization and toxicity study was conducted afterward. Proper controls and blanks for toxicity investigations, such as the lagoon sediment used in composting and a control compost with uncontaminated sediment, were not a part of the engineering study. In addition, the samples characterized in Phase 1 had been stored ca. 10 months, and the effects of such storage are unknown. These considerations in no way detract from the engineering study, but they do limit the conclusions which can be drawn from this work. A conclusive characterization of toxicity associated with the composting option will not be available until the second field composting experiment and the Phase 2 studies are complete.

CHAPTER 2. COMPOST SAMPLE PREPARATION AND CHEMICAL CHARACTERIZATION

2.1. Compost Acquisition and Sampling

One 55 gallon drum each of the mesophilic (pile no. 3) and the thermophilic (pile no. 4) composts from the LAAP field composting experiment (4) were received from Weston on April 17, 1989. The contents of the drums were first sampled for the laboratory studies on April 20, 1989. The drums were rotated at ca. 12 rpm for 5 min by rolling back and forth along a sloped concrete walkway. The drums were found to be ca. 7/8 full upon opening. The interior surfaces of each carbon-steel drum had rusted, but the structural integrity of the drums was not breached. The composts probably have been contaminated with a small amount of the rust. Six 1 L widemouth bottles were filled with compost taken from the upper portion of each drum. The drums were then archived in a monitored cold-storage (4°C) unit on April 24, 1989.

The second sampling of the drums was conducted on June 30, 1989, to obtain more compost for the organic solvent extraction experiments (sections 2.3.1 and 2.3.2). The drums were opened inside the cold storage facility, and a light mould growth was noted on the surface of the compost. The top layer was scraped aside and six 1 L samples were taken from underlying compost.

The third and final sampling was carried out on August 10, 1989, to provide sufficient compost for the preparation of the organic solvent extracts. Before this sampling, the composts were mixed in the drums by rotating them about their horizontal axis for 2 hrs. at 20 rpm. Three L volumes of compost were collected from each drum. Two L were used for the organic solvent extracts and 1 L was archived at -20°C. The drums were returned to 4°C storage.

2.2. Preparation and Chemical Characterization of Regulatory Leachates

2.2.1. Preliminary Trial of EPA Synthetic Precipitation Leach Test

A preliminary leaching was conducted on samples of the two composts using the EPA Synthetic Precipitation Leach Test ("Clean Closure Leach Test" or CCLT, SW-846 method 1312, reference 5). This preliminary run was performed to determine the behavior of the composts in leaching, and to generate leachate samples for range-finding in the toxicity tests. The latter was necessary to estimate the volumes of leachate which would be required for the toxicology battery. The leachates also were analyzed for explosive compounds and TNT metabolites.

As per the EPA protocol, the samples were leached with the "western" synthetic precipitation fluid (ASTM Type II water adjusted to pH 5.0 with a mixture of sulfuric and nitric acids) because the composts were generated at a site west of the Mississippi. Two L of CCLT leachate were generated from 100 g of each compost. It was observed that the required 10 mm screening of the composts removed mainly wood chips from the samples. These chips were analyzed later for explosive compound contamination (see section 2.6.). The main problem encountered in the CCLT preparation was that the filtration step was quite difficult and time-consuming. The EPA protocol pressure filter plugged rapidly from the fine clay particles present in the composted soil, and only ca. 250 mL could be filtered before plugging. A preliminary treatment of the crude leachate was needed to prevent this. Several pretreatments of the leachate were investigated in working with this leaching test and with the EPA Toxicity Characteristic Leaching Procedure (see below). The best pretreatment method appears to be a centrifugation of the crude leachate at 350 XG for 10 min. followed by prefiltration through a 1-to 3-um porosity, binderless glass fiber (Nucleopore type P100);

filter before the pressure filtration step. Ca. 1 L of crude leachate were filtered through the prefilter and the EPA protocol pressure filter step before the filters required replacement.

These preliminary leachates were analyzed for explosive compounds and TNT metabolites by HPLC (method described in Appendix A-2, compound abbreviations are listed in Appendix A-1). The compounds identified and concentrations determined in the leachates from the mesophilic and thermophilic composts were TNT (0.2 and <0.05 $\mu\text{g/mL}$ for the mesophilic and thermophilic leachates, respectively), 2-A-4,6-DNT (0.1 and 0.01 $\mu\text{g/mL}$), 4-A-2,6-DNT (0.13 and 0.07 $\mu\text{g/mL}$), HMX (0.98 and 0.27 $\mu\text{g/mL}$), and RDX (9.2 and 3.3 $\mu\text{g/mL}$). None of the diaminomononitrotoluenes were identified although there were very small peaks at retention times which were very close to those of the standard compounds. Trinitrobenzyl alcohol, tetryl, and the hydroxyaminodinitrotoluene were not detected (< ca. 0.05 $\mu\text{g/mL}$). A peak close to that of the azoxydimer also was observed.

2.2.2. Final Trial of EPA Synthetic Precipitation Leach Test

The CCLT was conducted upon the mesophilic and thermophilic composts as per the EPA protocol (5) to produce the large volume of leachates necessary to conduct the chemical and toxicity testing battery. Our toxicity testing of the preliminary CCLT run (see following chapters) indicated that 28 L of each leachate would be needed for the chemical analyses and toxicity tests. Of this, 26 L were required for the toxicity tests, most of which was for the fathead minnow larvae test. The CCLT leachates were generated over a period of ca. two weeks, and required 15 two-L tumbler bottle loads. The Extraction Fluid No. 2 (ASTM Type II water adjusted to pH 5.0 with a 60/40 wt.% mixture of sulfuric/nitric acids) was used because the composts were generated from a site west of the Mississippi (LAAP) and presumably would be surface-applied at or near that site. The EPA protocol was followed, with the addition of centrifugation (350 to 1,100 XG for 30 min.) and prefiltration immediately before the pressure filtration step to prevent plugging of the EPA protocol pressure filter. Experiments (see section 2.2.3.) with the EPA Toxicity Characteristic Leaching Procedure leachates indicated no significant effects upon leached explosives and TNT metabolites from this preliminary centrifugation and filtration.

The CCLT leachates were collected in 4 L jugs after preparation, and were composited after all of the necessary leachate had been generated. Two composited aliquots were prepared. Aliquot no. 1 (designated "CCLT-1" in Table 2-2) was composited from the first 14 L, and aliquots were withdrawn for chemical characterization only. The compositing was continued, and the second aliquot was composed of all 28 L (identified as "CCLT-2" in Table 2-2). One-half of the second aliquot was composed of the first; thus the two aliquots were not true duplicates. The second aliquot was subjected to both chemical analyses and toxicity characterization. Chemical analysis included semivolatile organic compounds, PCBs, pesticides, and metals (inductively coupled plasma emission spectroscopy screen [ICP] only) by EPA SW-846 (6) and Contract Laboratory Program (CLP) (7) procedures. Explosive compounds and TNT metabolites were determined using high performance liquid chromatography (HPLC) (see Appendix A-2) after U. S. Army Cold Regions Research and Engineering Laboratory methods (8,9). Toxicity characterization consisted of fathead minnow and Ceriodaphnia dubia aquatic toxicity tests and the Ames bacterial mutagenicity test.

2.2.3. EPA Toxicity Characteristic Leaching Procedure

Samples of both composts were subjected to the Toxicity Characteristic Leaching Procedure (TCLP) per the EPA protocol (10). A total of 2 L of leachate were generated from 100 g of the thermophilic compost, and 4 L were generated from 200 g of the mesophilic compost. The leaching medium, TCLP extraction fluid no. 1 (pH 4.93 acetate buffer), was used because the pH in the preliminary test was < 5.0. The pH of the solution prepared from 96.5 mL of ASTM Type II water

and 5.0 g of compost stirred for 5 min. was 9.1 for the mesophilic compost and 8.7 for the thermophilic compost. Per the TCLP protocol, 3.5 mL of 1 N hydrochloric acid were added and stirred at 50° C for 10 min., and the pH was redetermined. For the mesophilic compost, the resulting pH was 2.3, and for the thermophilic compost, the pH was 2.2. Because this was < 5.0, extraction fluid no. 1 was required. The only deviation from the EPA protocol was the centrifuging and prefiltering conducted before the EPA protocol pressure filtration. This pretreatment was similar to that used for the CCLT leachates.

Samples of the TCLP leachate were analyzed by EPA SW-846 (6) and CLP (7) procedures for semivolatile organic compounds, pesticides, PCBs, and metals (ICP screen only). Explosive compounds and TNT metabolites also were determined by HPLC (Appendix A-2). No toxicity tests were conducted on these leachates because of interferences by the acetate buffer.

To investigate the possible consequences of the prefiltration, samples of the final leachates of both composts were spiked with known concentrations of explosive compounds and TNT metabolites. A 55 mL volume of each leachate was spiked with 1.0 mL of a 1 µg/mL (each compound) standard of explosives and metabolites in acetonitrile. The spiked solutions were filtered through a 57 mm diameter filter of the same type as used for the prefiltration to preserve the same leachate volume/filter diameter ratio. The spiked leachates were sampled before and after prefiltration. The leaching medium also was sampled before and after prefiltration to determine if the prefilter contaminated the leachates. The samples were analyzed by HPLC. As shown in Table 2-1, no significant changes in concentration were found for TNT, HMX, RDX, two diaminomononitrotoluenes, two monoaminodinitrotoluenes, and trinitrobenzyl alcohol, but a 15-20% decrease was observed for 4-hydroxyamino-2,6-dinitrotoluene, and the azoxydimer. The latter two compounds are more hydrophobic than the other species and may have sorbed on fine particles which were subsequently filtered out. No contamination by the filter was detected.

TABLE 2-1. RECOVERY OF EXPLOSIVE COMPOUNDS AND TNT METABOLITES FROM PREFILTRATION OF TCLP LEACHATES FROM COMPOSTS

Compound	Recovery ^a , %	
	Mesophilic	Thermophilic
2,6-DA-4-NT	99	95
2,4-DA-6-NT	100	99
2,4,6-TNBAIc	102	97
RDX	99	100
HMX	97	93
2-A-4,6-DNT	98	99
4-A-2,6-DNT	98	99
TNT	98	100
4-OHA-2,6-DNT	84	83
Azoxydimer	83	80

^a0.9 µg/mL spike added to leachates. Recovery is calculated by (100%) x (peak area after filtration)/(peak area before filtration).
N=1

2.2.4. Chemical Characterization of Regulatory Leachates

The analyses consisted of gross leachate characteristics, explosives compounds and TNT metabolites by HPLC, semivolatile organic compounds (the "SVOA") and PCBs/pesticides by the EPA CLP and SW-846 methods 3550 and 8270 for SVOA and 8080 for PCBs/pesticides, and a metals screen by ICP.

Gross Chemical Characteristics of Leachates:

Two types of gross measurements were conducted upon the leachates immediately after their generation. Leachate pH was measured using a glass combination electrode, and the extractable matter was estimated by evaporating and reweighing 10 mL of leachate in a tared aluminum pan. As shown in Table 2-2, the pH of the TCLP extracts increased slightly during the extraction, while the blank remained at 4.9. A greater pH increase was observed for the CCLT leachates, consistent with the much lesser buffering capacity of the CCLT extractant (a very dilute combination of nitric and sulfuric acids) versus that of the TCLP (sodium acetate/acetic acid buffer). Most of the extractable mass from the TCLP was contributed by the sodium acetate/acetic acid buffer. It appears that a greater mass was extracted from the thermophilic compost than from the mesophilic compost. This also was observed with the CCLT leachates. In fact, the masses extracted by the CCLT leaching are close to those for the TCLP if the blank is subtracted from the TCLP. Additional characterization of the CCLT leachates, conducted as a part of the toxicity testing protocol, is reported in Table 3-3 of the following chapter.

TABLE 2-2. COMPARISON OF GROSS CHARACTERISTICS OF CCLT AND TCLP LEACHATES OF COMPOSTS

Leaching Method	Compost	pH	Extractable Matter	
			mg/mL ^a	mg/g ^b
CCLT-Prelim.	Meso.	8.4	0.51	10
	Thermo.	8.4	0.95	19
CCLT-Final	Meso.	7.8	0.92	18
	Thermo.	7.8	1.19	24
	Blank	6.1	0.025	0.5
TCLP	Meso.	5.05	6.85	137
	Thermo.	5.05	7.04	141
	Blank	4.9	5.53	111

^amg of matter per mL of leachate

^bmg of matter per g of compost

Explosive Compounds and TNT Metabolites

Of the organic chemical analyses, only the HPLC analyses (method described in Appendix A-2) revealed the presence of appreciable organic compounds. The data shown in Table 2-3 (abbreviations are listed in Appendix A-1) indicate that low $\mu\text{g/mL}$ concentrations of RDX and HMX are leached by both the TCLP and CCLT. It is not clear why the TCLP was a more aggressive extractant than the CCLT for the mesophilic compost. We suspect that it may be due to inhomogeneity in the compost (see below), and that the 200 g of mesophilic compost taken for the TCLP contained a "rich" deposit of the explosives, while the much greater mass of compost (1400 g) taken for the CCLT averaged out such inhomogeneities. The thermophilic compost leachates contained lower concentrations of explosives in both the TCLP and CCLT leaching than did the mesophilic compost leachates. This is consistent with the lower concentrations of these compounds in the composts, as determined by Weston (4) and our own analysis (see below). Approximately 40 - 100% of the available explosives and metabolites were leached from the compost. For the CCLT, the percentages leached, as averages of the data for the mesophilic and thermophilic compost leachates, were HMX (100%), RDX (57%), TNT (72%), 2-A-4,6-DNT (43%), and 4-A-2,6-DNT (46%).

TABLE 2-3. RESULTS OF EXPLOSIVES COMPOUNDS AND TNT METABOLITES ANALYSIS OF COMPOST LEACHATES

Compound	Concentration in Leachate ^a , $\mu\text{g/mL}$					
	Mesophilic			Thermophilic		
	TCLP ^b	CCLT-1 ^b	CCLT-2 ^c	TCLP ^b	CCLT-1 ^c	CCLT-2 ^c
2,6-DA-4-NT	-	-	-	-	-	-
2,4-DA-6-NT	-	-	-	-	-	-
2,4,6-TNBAlc	-	-	-	-	-	-
RDX	16.6	6.1	6.2	0.5	2.0	1.2
HMX	1.5	0.8	0.8	-	0.6	0.2
TNB+DNB	-	-	-	-	-	-
2-A-4,6-DNT	0.13	0.08	0.08	-	0.11	-
4-A-2,6-DNT	0.11	0.18	0.17	-	0.40	0.09
2,6-DNT	-	-	-	-	-	-
2,4-DNT	-	-	-	-	-	-
TNT	5.3	3.0	3.7	0.1	0.90	1.0
Tetryl	-	-	-	-	-	-
4-OHA-2,6-DNT	-	-	-	-	-	-
Azoxydimer	-	-	-	-	-	-
Total ID	23.6	10.2	11.0	0.6	4.0	2.5

^a - =* = not detected (ca. 0.01-0.05 $\mu\text{g/mL}$). Concentrations < 0.1 $\mu\text{g/mL}$ are estimates.

^b n = 1

^c n = 2

Regulatory Organic Compounds

Appendix A-3 contains the reporting forms for the characterization of the TCLP leachates and Appendix A-4 contains those for the CCLT leachates. Briefly, the SVOA did not detect any of the 65 EPA Target Compound List (TCL) semivolatile organic constituents in the TCLP or CCLT leachates of either compost, with reporting limits of 11-65 $\mu\text{g/L}$ (depending upon the exact constituent). The SVOA Tentatively Identified Compounds ("TICs") included TNT and 2-5 compounds which were not identified by the NBS mass spectral library machine search. Comparison of their mass spectra with those of authentic standards identified two of these compounds as the monoaminodinitrotoluenes.

None of the 20 TCL pesticides were found in the TCLP leachates with reporting limits of 0.5 - 5 $\mu\text{g/L}$, and none of the 7 TCL PCBs were found at reporting limits of 5-10 $\mu\text{g/L}$ (Appendix A-3). In contrast, the analysis of the CCLT leachates (Appendix A-4) did reveal the presence of low concentrations of several pesticides, probably because a larger volume of leachate was analyzed, and the sensitivity was better. Three pesticides were determined at the same concentrations in both the mesophilic and thermophilic CCLT leachates: beta-BHC at 0.3 $\mu\text{g/L}$, gamma-BHC at 0.2 $\mu\text{g/L}$, and dieldrin at 0.01 $\mu\text{g/L}$. Gamma-chlordane was estimated at 0.12 $\mu\text{g/L}$ in the mesophilic leachate and at 0.04 $\mu\text{g/L}$ in the thermophilic leachate, which suggests that the composting may have reduced its concentration. In addition, a trace (0.02 $\mu\text{g/L}$) of heptachlor was estimated in one of the aliquots of the mesophilic leachate. These concentrations are too low to be significant, and the compounds most likely were not contributed by the explosives-contaminated sediment, but rather from the amendments used in composting. They would be highly site- and amendment-specific.

Volatile organic compounds also were determined in CCLT leachates prepared separately from the others. The zero headspace extractor was not used because it is difficult to operate and the history of the compost samples suggests that the effort required to use the zero headspace extractor would not be justified. The composts had been aerated (4) during composting and then they were stored in a partially filled drum for nearly one year before analysis. Most volatile organic compounds would be lost. Therefore, 2 g samples of the composts were shaken for 18 hrs with 42 mL of the CCLT extraction fluid in a 40 mL VOA vial. The CLP VOA surrogate standards were added to the vials before shaking. The analysis by the CLP purge and trap GC-MS method (similar to SW-846 method 8240) did not reveal any VOA different from the blank with reporting limits of 5-10 $\mu\text{g/L}$ (Appendix A-4). Although the surrogate standard recoveries were low for toluene and bromofluorobenzene, any volatiles present would have been detected.

Metals Screen by ICP

The results for the determination by ICP of 29 elements in the TCLP and CCLT leachates of the two composts and leaching blanks are listed in Appendix A-3 and A-4. The metals compositions of the mesophilic and thermophilic leachates were almost identical because they are contributed by the soil, and not the munitions, and are not bacterially degraded. They would be highly site-specific.

2.2.5. Leachate and Compost Homogeneity

The final report by Weston (4) shows that the composts are inhomogeneous with respect to explosives concentrations. Obtaining representative samples for study is important. This concern was addressed in the mixing of the contents of the drums before sampling. In the generation of the CCLT leachates for toxicity testing and chemical analysis, the relatively large masses of compost (1400 g) required to generate the 28 L of leachate allowed a considerable "averaging" of local inhomogeneities in the composts. This was demonstrated by HPLC analysis of individual 4 L

aliquots which were composited to prepare the CCLT leachate of the thermophilic compost. Table 2-4 shows the variability in the results for RDX and TNT. The concentrations for each ranged from "not detected" (ca. 0.05 $\mu\text{g/mL}$) to nearly 2 $\mu\text{g/mL}$, and the relative standard deviations were 60% for RDX and 122% for TNT. Thus, the final composited leachate "averaged" a wide variation in compost explosives concentrations.

TABLE 2-4. VARIATION IN RDX AND TNT CONCENTRATIONS IN INDIVIDUAL 4L PORTIONS OF THERMOPHILIC COMPOST CCLT LEACHATES

Aliquot	Concentration, ^a $\mu\text{g/mL}$	
	RDX	TNT
1	1.3	1.0
2	1.8	1.8
3	ND	0.2
4	0.8	1.1
5	0.9	0.6
6	0.7	ND
7	0.8	ND
---	---	---
Avg.	0.91	1.39
SD	0.54	1.67
RSD	60%	122%

^aND = not detected (use 0.05 $\mu\text{g/mL}$ for calculations).

Our previous experiences in preparing the TCLP and CCLT leachates, as well as Weston's analytical data on the compost piles, indicate that a lack of homogeneity in the composts must be considered in the design of experiments. Preliminary to the comparison of organic solvents for compost extraction, samples of the composts were mixed carefully and analyzed to establish their homogeneity as a baseline before the extractions were carried out. Ca. 1.5 kg of each compost were sieved to ≤ 10 mm per the TCLP and CCLT protocol, and then were sieved again to ≤ 3.35 mm. The sieved material was then tumbled for ca. 1 hr in the TCLP apparatus (without solvent), spread in a glass tray, and mixed again with a spatula. Attempts to homogenize the composts with a food homogenizer or with a grinder failed because the composts caked and plugged the equipment. The composts were not dried because of concerns that explosives metabolites could be oxidized.

Duplicate 5 g aliquots were taken from five points in the glass tray, and these were extracted with 20 mL of acetonitrile for 18 hrs in an ultrasonic bath, and were analyzed using HPLC. The data are included in Table 2-5. The relative standard deviations (RSD) ranged from 5 to 26% for all compounds (except for TNT) for the mesophilic compost, and from 8.5 to 53% for the thermophilic compost. The somewhat better homogeneity for the mesophilic compost may be due to its lesser

TABLE 2-5. EXAMINATION OF HOMOGENEITY OF COMPOST SAMPLES USED FOR SOLVENT COMPARISON STUDY

Compound	Concentration in Compost ^a , µg/g					
	Mesophilic			Thermophilic		
	Range	Avg ± SD (RSD)	Range	Avg ± SD (% RSD)	Range	Avg ± SD (% RSD)
RDX	77-196	129 ± 33.7 (26%)	12-36	23 ± 8.4 (37%)		
HMX	29-44	35 ± 5.3 (15%)	3-14	6.0 ± 3.2 (53%)		
2-A-4,6-DNT	3.3-4.4	3.7 ± 0.37 (10%)	1-2	1.87 ± 0.42 (23%)		
4-A-2,6-DNT	7.4-8.6	8.0 ± 0.44 (5.5%)	3.2-4.2	3.6 ± 0.31 (8.5%)		
TNT	30-3530	487 ± 1084 (223%)	8-1400	179 ± 430 (240%)		
(TNT) ^b	30-213	89 ± 68 (77%)	8-76	34 ± 24 (72%)		

^an = 10

^bTwo highest results deleted

moisture content, and the greater ease in mixing a less sticky compost. The RSDs for TNT in both composts were ca. 240% for all 10 data points, and dropped to ca. 75% when the two highest data points (for $n=8$) were deleted. The lesser precision for TNT suggests either considerable inhomogeneities in the distribution of the TNT in the composts or an analytical interference that itself is not constant. These RSDs are similar to those observed by Weston (4) over one year ago at the end of the field experiment at the LAAP. The concentration data, with the exception of TNT, are also similar to the Weston data. The TNT was higher in our measurements, again suggesting that the TNT peak is subject to an analytical interference, or that the CRREL extraction procedure was unable to recover all the TNT.

2.3. Preparation and Characterization of Organic Solvent Extracts

Organic solvent extracts were prepared from the two composts to provide material enriched in potentially toxic compounds for chemical analysis and toxicological testing. The aqueous leaching would not be expected to mobilize toxic organic compounds as readily as organic solvent extraction, and thus toxicity in the composts could be missed by the latter. The solvents to be used and the extraction method were first determined before preparing extracts for characterization.

2.3.1. Comparison of Organic Solvents for Compost Extraction

Seven solvents were compared for their ability to extract explosives and metabolites from the composts. For this purpose, the homogenized composts samples described immediately above were used. Triplicate 5 g aliquots of the mesophilic compost were extracted for 18 hrs in an ultrasonic bath using 20 mL of each solvent. The solvents were chosen because of their proven efficiency for explosives (i.e., acetonitrile), their extensive use in bioanalytical (i.e., ethyl acetate) or environmental (i.e., methanol, methylene chloride) analysis, or their good solubility characteristics (i.e., tetrahydrofuran, diethyl ether, and dimethyl sulfoxide [DMSO]). Explosives and metabolites analyses were performed by HPLC, and the total extractable matter was estimated by weighing the residue from evaporating 10 mL of the extract. The results are listed in Table 2-6. DMSO was not evaporated because of its relatively poor volatility. Of the seven solvents, acetonitrile and ethyl acetate appeared to offer the best recoveries, and were selected for preparing the extracts for chemical and toxicological characterization. It is not clear why TNT and one of the metabolites were not detected in the DMSO extraction.

2.3.2. Comparison of Organic Solvent Extraction Procedures

Several different extraction procedures have been used in these studies. These include the U.S. Army Cold Regions Research and Engineering Laboratory (CRREL) method (8), the U. S. EPA SW-846 method 3550 (6), and ORNL variations of both methods. Results of the applications of these methods to the composts are shown in Table 2-7. Basically, the ultrasonic extractions performed using a probe appear to be more efficient than those done with a bath, although compost inhomogeneity complicates comparison of the results. However, the data do indicate that the procedure (90 g of compost extracted three times with 300 mL of solvent [each time] using a probe-type sonicator) chosen for preparing the extracts for toxicological testing was at least as effective as the CRREL analytical method while providing nearly two orders of magnitude scaleup. Analysis of the extracts from each of the three extraction steps showed that 80% of the extracted explosives and metabolites were recovered on the first extraction, 16% on the second, and 4% in the third extraction. These results also suggest the completeness of the extraction.

TABLE 2-6. COMPARISON OF ORGANIC SOLVENTS FOR MESOPHILIC COMPOST EXTRACTION^a

Avg \pm Std. Dev. (n = 3 unless otherwise noted)					
Solvent	Total Extractable Matter, mg/g	Explosives and Metabolites, μ g/g			
		RDX	HMX	2-A-4, 6-DNT	4-A-2, 6-DNT
Acetonitrile ^b	3.5 ^c	129 \pm 34	35 \pm 5	3.7 \pm 0.4	8.0 \pm 0.4
Methanol	5.1 \pm 0.4	63 \pm 37	10 \pm 1.5	2.7 \pm 0.6	6.0 \pm 1
Diethyl Ether	2.1 \pm 0.6	165 \pm 41	8 \pm 1	4.0 \pm 0	8.7 \pm 1.2
Methylene Chloride	2.0 \pm 0.3	200 \pm 97	13 \pm 1.7	3.6 \pm 0.6	8.7 \pm 0.6
Tetrahydrofuran	5.5 \pm 0.4	124 \pm 25	29 \pm 3.6	3.3 \pm 1.2	5.6 \pm 4.9
Ethyl Acetate	3.8 \pm 0.4	186 \pm 90	25 \pm 23	5.7 \pm 1.2	10 \pm 4.4
Dimethyl Sulfoxide	^f	132 \pm 22	31 \pm 11	ND ^g	8.0 \pm 1.7
					ND ^h

^a5g compost extracted overnight with 20 mL solvent.^bn = 10 for explosives.^cTotal extractable matter is weighed residue from composite of 10 replicates.^dn = 8 with 2 highest data points deleted.^en = 2 with highest data point deleted.^fSolvent not evaporated.^gND = not detected

TABLE 2-7. COMPARISON OF SOLVENT EXTRACTION METHODS FOR COMPOSTS

Compost	Extraction Method	Avg \pm STD, $\mu\text{g/g}$ (n = 3 unless otherwise indicated)				
		RDX	HMX	2-A-46 DNT	4-A-26 DNT	TNT
Mesophilic	CRREL, 1g/25ml, Bath	157 \pm 74	27 \pm 8.7	ND	6 \pm 1	13 \pm 14
	ORNL, 5g/30ml, Bath	141 \pm 32	35 \pm 9.0	3.6 \pm 0	10 \pm 0	318 \pm 470
	ORNL, 5g/20ml, Bath ^b	129 \pm 34	35 \pm 5.3	3.7 \pm 0.4	8 \pm 0.4	47 ^a
	ORNL, 30g/100ml Probe ^a	170	37	3.5	8.2	487 \pm 1080
	SW-846, 30g/100mlx3, Probe	273 \pm 68	52 \pm 9.5	5.1 \pm 0.2	12.7 \pm 1.5	89 \pm 68 ^c
Thermophilic	ORNL, 90g/300mlx3, Probe ^d	226 \pm 34	51 \pm 3.9	6.0 \pm 0.4	16 \pm 0.5	473
	ORNL, 5g/20 mL, Bath ^b	23 \pm 8.4	6.0 \pm 3.2	1.8 \pm 0.4	3.6 \pm 0.3	179 \pm 430
	ORNL, 30g/100mL Probe ^a	23	5.3	1.3	3.7	34 \pm 24 ^c
	SW-846, 30g/100mLx3, Probe	13 \pm 13	ND	ND	3.5 \pm 0.7	483
	ORNL, 90g/300mLx3, Probe ^d	80 \pm 81 40 \pm 17 ^c	9.3 \pm 9.2	2 \pm 0	6.8 \pm 0.8	28 \pm 27
						158 \pm 158

^an = 2^bn = 10^cn = 8, 2 highest data points deleted^dBatch 2 of compost, n = 4^en = 3, highest data point deleted

2.3.3. Preparation of Organic Solvent Extracts

A preliminary small-scale extract was prepared from both composts to provide material for range-finding in the toxicological assays, and to estimate the mass of extractable matter needed for the full toxicological tests. Ca. 30 g of each compost were extracted with 100 mL of acetonitrile using a probe-type ultrasonicator. Three batches of each compost were extracted (total of 90 g). The solvent was removed using rotary evaporation, and the residues were resuspended in DMSO for toxicity testing.

On the basis of the preliminary toxicity tests, it was estimated that ca. 2 g of extractable matter were needed for the full battery of toxicity tests and chemical characterization. The 55 gallon drums of compost were resampled to provide enough compost for the extractions. The drums were rotated for several hours and the upper portion was mixed with a shovel before sampling. Ca. 1 kg aliquots were sieved through 10 mm and 3.35 mm screens and were mixed further by tumbling and stirring to promote homogeneity. Subaliquots of 450 g of the mesophilic compost and 900 g of the thermophilic compost were extracted 90 g per batch, using three 300 mL volumes of acetonitrile and ultrasonic extraction with a probe-type sonicator, after EPA SW-846 method 3550. This solvent was removed by rotary evaporation. For the ethyl acetate, a total of 180 and 360 g of compost (respectively) were used, and the same extraction protocol was followed. The residues from both sets of extractions were redissolved in DMSO. Table 2-8 summarizes the preparation parameters. Chemical characterization of the DMSO-reconstituted extracts included semivolatile organic compounds, PCBs/pesticides, and explosive compounds and TNT metabolites. Toxicity tests consisted of fathead minnow larvae (acetonitrile extract only) and *Ceriodaphnia dubia* freshwater toxicity and the Ames microbial mutagenicity test.

2.3.4. Chemical Characterization of Organic Solvent Extracts

The acetonitrile and ethyl acetate extracts generated from the mesophilic and thermophilic composts were analyzed for EPA semivolatile organic and PCB/pesticide Target Compound List species. For this characterization, aliquots of the extracts were solvent exchanged from DMSO into methylene chloride (semivolatile organics) or hexane (PCBs/pesticides) at a concentration of ca. 4 mg of extractable residue/mL by azeotropic distillation and redissolution. The analyses were performed by EPA Contract Laboratory Program GC-MS (semivolatile organics) and GC with electron capture detection (PCBs/pesticides) procedures. None of the TCL semivolatile organics were detected with a reporting limit of 10 to 50 mg/L, none of the pesticides with a reporting limit of 200 µg/L, and none of the PCBs with a reporting limit of 20 mg/L. As expected, TNT was detected as a TIC in the semivolatile organics analysis. Explosives and TNT metabolites analyses performed on the DMSO solutions by HPLC are included in Table 2-9. The explosives and metabolites concentrations were much higher than those in the aqueous leachates. There may have been an interference in the HPLC peak for HMX (note that the apparent HMX concentration extracted by ethyl acetate from the thermophilic compost was higher than that from the mesophilic compost) but the spectra did not disprove the apparent identification.

Qualitative gas chromatographic scans were made of the solvent extracts to supplement the GC-MS analyses. Ten mL aliquots of the preliminary acetonitrile extracts of the mesophilic and thermophilic composts were taken to dryness under a gentle stream of nitrogen, and were extracted with 1 mL of acetonitrile. Not all of the residue redissolved, even after warming the flask, suggesting the presence of inorganic or high molecular weight matter. Two µL injections were made by the solvent flush technique into a capillary column gas chromatograph equipped with a splitless injection port, a fused silica capillary column (30 m x 0.25 mm ID x 0.25 µm film) with DB-5 bonded phase, and a flame ionization detector. The column temperature program was 100°C (5 min. isothermal hold) to 300°C at 8°C/min with a 30 min. hold at 300°C, the injector and detectors were

TABLE 2-8. PARAMETERS OF ORGANIC SOLVENT EXTRACTS FOR CHEMICAL AND TOXICITY TESTING.

Parameter	Acetonitrile Extract			Ethyl Acetate Extract	
	Mesophilic	Thermophilic	Mesophilic	Thermophilic	
Mass of Compost Extracted, g	450	900	180	360	
Extractable Matter, g	1.33	2.03	0.85	0.92	
Volume of DMSO for Reconstitution, mL	135	270	54	108	
Concentration in DMSO					
mg Residue/mL	9.9	7.5	15.7	8.5	
g Compost/mL	3.3	3.3	3.3	3.3	

TABLE 2-9. EXPLOSIVE COMPOUND AND TNT METABOLITE ANALYSIS OF ORGANIC SOLVENT EXTRACTS AND HYDROLYZED BOUND FRACTION

Solvent Extract/Compost or Hydrolyzed Bound Fraction/ Compost	Concentration ^a , µg/mL				
	RDX	HMX	TNT	4-A-2, 6-DNT	2-A-4, 6-DNT
Ethyl Acetate/Mesophilic	1900	26	515	14	13
Ethyl Acetate/Thermophilic	750	125	50	6	4
Acetonitrile/Mesophilic	1550	210	550	13	12
Acetonitrile/Thermophilic	710	119	20	3	-
Hydrolyzed Mesophilic	3	-	-	101 ^b	-
Hydrolyzed Thermophilic	4	-	-	120 ^b	-
Hydrolysis Blank	-	-	-	-	-

^a " = not detected (ca. 0.3 µg/mL)

^b HPLC peak observed at correct retention time window and spectra did not allow elimination.

maintained at 270°C and 300°C (respectively), and the helium carrier gas flow rate was 1 mL/min. The extracts were then derivatized by adding 1 mL of N-methyl-N-trimethylsilyltrifluoroacetamide and letting the vial set at room temperature overnight. A 3 μ L aliquot was analyzed similarly. The chromatograms are shown in Figure 2-1. Very little gas chromatographable matter is visualized by the direct analysis of the solvent extracts (Figure 2-1 A and B). The mesophilic compost extract exhibits ca. 3-fold more peak area per mg of extractable matter than does the thermophilic compost extract. The main peaks are TNT and the monoaminodinitrotoluenes and some higher molecular weight material toward the end of the chromatogram. This confirms the small number of compounds detected by the GC-MS. In contrast, the derivatized extracts (Figure 2-1 C and D) show 4 to 5 times more detectable matter, indicating that the bulk of the organic matter is polar in nature. This suggests that the technique of derivatization followed by gas or supercritical fluid chromatography would be a powerful analytical technique to analyzing compost extracts and leachates.

2.4. Preparation and Chemical Characterization of Hydrolyzed Bound Fraction

Aliquots of the ethyl acetate-extracted composts were hydrolyzed to prepare samples for tests of toxicity associated with a bound, but chemically labile form of explosives transformation product. For example, transformation products bound through hydrolyzable amide bonds would be hydrolysed back to the original amines while products bound through nonhydrolyzable ether or heterocyclic linkages would not be liberated. The solvent-extracted composts remaining from the studies described in section 2.3. were hydrolyzed because all the unbound explosives compounds and their metabolites had been removed, and explosives-related compounds (if any) could be attributed to their bound forms.

Fifty g of the ethyl acetate-extracted mesophilic and thermophilic composts were refluxed for 6 hrs with 250 mL of 6 N hydrochloric acid following earlier work (11) showing the release of organic compounds from humic acids by boiling acid. The hydrolyzate was filtered off and extracted three times with 250 mL of methylene chloride, made alkaline (pH 11) with 50% sodium hydroxide solution, and re-extracted with methylene chloride. The methylene chloride extracts were combined and taken to dryness, and the residue was weighed and redissolved in dimethylsulfoxide (DMSO). A blank (no compost) and a positive control (50 g of ethyl acetate-extracted thermophilic compost spiked with 50 mg each of 4-amino-2,6-dinitrotoluene and 2,4-diamino-6-nitrotoluene) also were treated similarly. The only noteworthy observation from the preparation was that a precipitate formed upon addition of the sodium hydroxide to the hydrolyzate. The precipitate probably was composed of humic acids. It was filtered out and rinsed with methylene chloride to recover any products. Table 2-10 lists the preparation parameters for the experiment. The DMSO-reconstituted residues were subjected to the Ceriodaphnia dubia toxicity test and the Ames bacterial mutagenicity test.

Characterization was limited to HPLC analysis of explosives and TNT metabolites. The results are included in Table 2-9. HPLC peaks corresponding to a trace of RDX and 4-A-2,6-DNT were observed in the final extracts. The spectra of the peak for the latter would not allow conclusive confirmation or elimination of the peak identity, so the identification must be considered as tentative at the present. The spikes were not recovered in the spiked positive control sample. It was found in a subsequent experiment that the acidic refluxing of the spikes in the presence of the compost caused them to decompose. Future work must include more gentle hydrolysis conditions.

FIGURE 2-1. CAPILLARY COLUMN GC OF CRUDE AND DERIVATIZED COMPOST EXTRACTS. ELUTION OF TNT IS LABELED. BRACKETS SHOW ELUTION RANGE OF METABOLITES (CRUDE) AND TNT PLUS METABOLITES (TMS-DERIVATIZED).

ORNL-DWG 90-2208

CAPILLARY GC OF CRUDE AND DERIVATIZED COMPOST EXTRACTS




TABLE 2-10. PARAMETERS FOR PREPARATION OF HYDROLYZED
BOUND FRACTIONS FOR CHEMICAL AND TOXICITY TESTING

Parameter	Results for Sample		
	Mesophilic Compost	Thermophilic Compost	Blank
Mass of Compost, g	50	50	-
Mass of Bound Fraction Residue, mg	270	219	1.5
Volume of DMSO for Bound Fraction, mL	15	15	1.5
Concentrations in DMSO: mg Bound Fxn/mL DMSO	18.0	14.6	0.1
g of Compost/mL DMSO	3.3	3.3	-

2.5. Preparation of Particle-bound Organics Extract from CCLT Leachate

Our previous work suggested that organic compounds leached from the composts by the aqueous leaching tests could be adsorbed on particles. Organic compounds on particles would not be available for mutagenicity testing when the leachate is sterilized by filtration before application to the agar plate. Similarly, submicron-sized particles would not be ingested by the freshwater organisms unless they were attached to food particles. To investigate this, samples were prepared to test for particle bound mutagenicity. Four L of CCLT leachate were generated from 200 g of the mesophilic compost using the procedure we described in section 2.2.2. The CCLT leachate (3.9 L) was centrifuged for 30 min at 2900 rpm (rotor radius of 19 cm) to isolate 701 mg of particles. The particles were extracted with DMSO and were subjected to Ames mutagenicity testing. HPLC analysis of the particle extract revealed RDX (70 $\mu\text{g/g}$ of particles), 2-A-4,6-DNT (120 $\mu\text{g/g}$), and 4-A-2,6-DNT (40 $\mu\text{g/g}$). It is not clear why TNT and HMX (which were present in the leachate) were not detected.

2.6. Analysis of Mesophilic Compost Wood Chips for Explosive Compounds and TNT Metabolites

Wood chips were screened out of the composts during the preparation for regulatory leaching or organic solvent extraction, and it is possible that they could have absorbed explosive compounds and biotransformation products. This could occur while the chips were insulating the piles or (more

an HPLC analysis of the wood chips screened out of the mesophilic compost. The results listed in Table 2-11 show definite sorption of explosive compounds and metabolites by the wood chips. The observation that the concentration ratios of the compounds in the wood chips to those in the compost are reasonably consistent for the compounds (except for TNT) suggests that the sorption occurred post-composting (i.e., while stored in the drum) and not during the composting. The latter could create an additional waste problem at the composting field site.

TABLE 2-11. CHARACTERIZATION OF MESOPHILIC COMPOST WOOD CHIPS FOR EXPLOSIVES AND BIOTRANSFORMATION PRODUCTS

Compound	Concentration, $\mu\text{g/g}$	Ratio, chips/compost
RDX	68 ± 11	0.29
HMX	20 ± 1.5	0.39
2-A-4,6-DNT	2.7 ± 0.6	0.45
4-A-2,6-DNT	9.0 ± 1.0	0.56
TNT	25 ± 8.7	0.04

2.7. Conclusions

The chemical analyses demonstrate that the greater biotransformation of explosives achieved with the thermophilic composting condition versus the mesophilic composting condition results in lower concentrations of explosives and metabolites in the aqueous leachates and organic solvent extracts of the composts. The main compounds determined in the leachates and extracts were RDX, HMX, TNT, 2-A-4,6-DNT, and 4-A-2,6-DNT. Substantial percentages of these compounds in the composts can be leached by the CCLT or TCLP. GC-MS, GC-ECD, and ICP analyses of leachates and extracts do not reveal any EPA TCL constituents at significant concentrations, but the presence of these compounds would be site-specific and not necessarily associated with the explosives wastes which were composted. Additional polar organic compounds in the extracts can be visualized by the technique of chemical derivatization and GC, but at the present they cannot be assigned to explosives compounds or their transformation products because of the lack of a compost blank.

CHAPTER 3. AQUATIC TOXICITY TESTS

3.1. Introduction

The U.S. Environmental Protection Agency (EPA) and regulatory agencies in many states are requiring, with increasing frequency, the use of bioassays to (a) determine whether or not treated and nontreated wastewaters are suitable for release to the environment, (b) document water quality in streams that receive runoff or point-source discharges of treated or nontreated wastes, and (c) identify the toxic constituent(s) that reduce the biological quality of problematic wastewaters. Although many types of toxicity tests can provide meaningful information, "mini-chronic" 7-d static-renewal tests that use the survival and growth of fathead minnow larvae (*Pimephales promelas*) and the survival and fecundity of *Ceriodaphnia dubia* to quantify toxicity have become widely accepted and are now the preferred systems for assessing the biological quality of effluents and freshwater ambient receiving systems by many states. Recent studies show how tests with these two species can be applied to (a) toxicity assessments for effluents and ambient waters (cf. references 12-14) and (b) wastewater treatment process optimization efforts (15,16).

3.2. Methods

The present study was designed to provide information about (a) the chemical constituents present in composted high explosives, and (b) the toxicity and potential mutagenicity of materials extracted from the compost. Both aqueous and organic-solvent extracts of the compost were tested for toxicity and mutagenicity. The toxicity evaluations, which are discussed in this section, were made using fathead minnow larvae and *Ceriodaphnia*; the procedures used for these two tests closely followed EPA method 1000.0 (for the fathead minnow larvae survival and growth test) and EPA method 1002.0 (for the *Ceriodaphnia dubia* survival and reproduction test); EPA methods 1000.0 and 1002.0 are described by Horning and Weber (17). Some latitude is permitted in the use of these two test procedures. The type of dilution water, for example, can be varied according to the objectives of the study (reference 17, p. 17). In tests reported here, controls and test solutions were prepared using mineral water (Perrier^R) diluted to 10% of full-strength with deionized distilled water. Batches of water prepared for the tests were degassed by gentle aeration for 24 h at room temperature before use. Our experience has shown that diluted mineral water (DMW) prepared in this manner is satisfactory for both *Ceriodaphnia* and fathead minnow larvae. In the case of the *Ceriodaphnia* test, EPA method 1002.0 recommends the use of a mixture of fermented trout chow, cerophyll, and yeast (TCY). However, various studies, including many conducted in our laboratory, have shown that additions of algae (e.g., *Ankistrodesmus falcatus*) to TCY can increase *Ceriodaphnia* fecundity and decrease variability of fecundity among replicates. The net effect of such food amendments is more statistical power and a greater ability to quantify toxicity. Thus, the *Ceriodaphnia* used in all tests reported here were fed with *Ankistrodesmus*-augmented TCY. Standard operating procedures detailing the method for preparing algal-amended TCY, and all other procedures of the tests with both species, are available upon request (18). *Ceriodaphnia* are much more sensitive than fathead minnow larvae to a wide variety of effluents (cf. references 14 and 19). Additionally, *Ceriodaphnia* tests can be conducted using less than 20% of the amount of test solution that is needed for the fathead minnow test, and have been shown to be more cost-effective than the minnow test (20). For these reasons, we used the *Ceriodaphnia* test rather than the minnow test for many of the assessments reported here.

In some instances, the amount of material available for testing was small. Some of the organic-solvent extracts of the compost, for example, when brought to dryness, contained a few hundred milligrams of residue. These residue samples were analyzed both chemically and biologically (i.e.,

for toxicity and mutagenicity). When such small quantities of residue were available, it was sometimes necessary to scale down even the Ceriodaphnia test system. Normally, the Ceriodaphnia test is conducted by exposing single neonates to 17 mL volumes of test solution. Pre-rinsed 20 mL polystyrene microbeakers are used in this situation; 10 replicates are typically used to provide the necessary statistical power for accurate toxicity estimates. We used side-by-side comparisons of two test procedures (single Ceriodaphnia in 2.5 mL volumes of test solution in borosilicate glass test tubes, versus single Ceriodaphnia in 17 mL volumes of test solution in polystyrene microbeakers) to verify that a scaled down daphnid test system could be used to provide accurate estimates of toxicity.

Quality assurance/quality control (QA/QC) is an important aspect of toxicity testing if the results of the tests are to be used for regulatory purposes. Reports in Appendix A-5 show the kinds of QA/QC information available for each of the toxicity tests reported here; more complete information regarding the QA/QC for any of the tests described in the following sections is available upon request.

3.3. Observations

The toxicity tests included preliminary and final assessments of CCLT leachates of mesophilic and thermophilic composts (Tables 3-1 through 3-3), preliminary tests with acetonitrile extracts of mesophilic and thermophilic composts (Table 3-4), final tests of acetonitrile and ethyl acetate extracts of mesophilic and thermophilic composts (Tables 3-5 and 3-6), and a test of an acid-hydrolyzed extract of the bound fractions from mesophilic and thermophilic composts (Table 3-7). The use of controls and blanks among the various experiments differed according to objectives specific to the different tests, and thus are described, when appropriate, in the context of the individual tests below.

3.3.1. Tests of Preliminary CCLT Leachates

Toxicity of the CCLT leachates from both compost types was evaluated using Ceriodaphnia. Four concentrations (30%, 20%, 10%, and 5% of full-strength) of each leachate were tested to estimate toxicity. This test was conducted in 2.5-mL test chambers; two concentrations (20% and 5%) of each leachate type were also tested using 17-mL test chambers. The data for these tests are shown in Table 3-1.

Survival and reproduction of Ceriodaphnia was high both in the controls and in all tested concentrations of the CCLT leachates, and there was no difference in survival or reproduction attributable to the type of test system (i.e., 2.5-mL vs. 17-mL). Thus, the tests did not detect acute or chronic toxicity in either leachate up to highest concentration tested (30% of full-strength).

3.3.2. Tests of Final CCLT Leachates

Toxicity of the CCLT leachates prepared from the mesophilic and thermophilic composts was evaluated using both Ceriodaphnia and fathead minnow larvae. Each leachate was tested at four concentrations (40%, 60%, 80%, and 100% of full-strength) with each species. The synthetic acid rainwater (CCLT blank) used to prepare the leachates was also tested (at 40% and 60% of full-strength) to ensure that leachate toxicity, if detectable, could be attributed to materials originating from the compost rather than from the water used to prepare the leachates. The results of the tests are summarized in Table 3-2.

TABLE 3-1. RESULTS OF CERIODAPHNIA TOXICITY TESTS OF PRELIMINARY CCLT LEACHATES IN 2.5 ML (UPPER) AND 17-ML (LOWER) TEST CHAMBERS

Sample	Concentration (%)	Survival ^a (%)	Fecundity ^b (offspring per female)
Control ^c	100	90	18.7 ± 1.3
Thermophilic	5	100	22.2 ± 1.8
Thermophilic	10	100	21.4 ± 2.0
Thermophilic	20	100	22.0 ± 2.1
Thermophilic	30	90	23.0 ± 2.5
Mesophilic	5	90	22.1 ± 3.1
Mesophilic	10	100	22.2 ± 1.6
Mesophilic	20	90	20.6 ± 1.4
Mesophilic	30	90	20.2 ± 1.6

17-ML TEST CHAMBERS

Sample	Concentration (%)	Survival ^a (%)	Fecundity ^b (offspring per female)
Control ^c	100	100	21.2 ± 1.6
Thermophilic	5	100	19.7 ± 2.2
Thermophilic	20	100	20.2 ± 3.1
Mesophilic	5	100	19.9 ± 2.6
Mesophilic	20	90	17.8 ± 3.4

^aTen replicates were used for each treatment. Thus, 90% survival (e.g., for the control) means that nine of ten animals survived all 7 d.

^bMean ± 1 SD, calculated using only females that survived the full 7-d test period, and their offspring. n=10.

^cDiluted mineral water.

TABLE 3-2. RESULTS OF CERIODAPHNIA AND FATHEAD MINNOW LARVAE TOXICITY TESTS OF CCLT LEACHATES PREPARED FROM THERMOPHILIC AND MESOPHILIC COMPOSTS

Sample Type and Concentration	Fathead minnow larvae		Ceriodaphnia	
	Survival ^a (%)	Growth ^b (mg/fish \pm 1 SD)	Survival ^c (%)	Fecundity ^d (offspring per female \pm 1 SD)
Control ^e	97.5	0.36 \pm 0.02	91.6	20.6 \pm 4.3
Thermophilic				
40%	100.0	0.47 \pm 0.03	91.6	26.3 \pm 8.0
60%	93.0	0.41 \pm 0.04	100.0	21.3 \pm 7.4
80%	90.0	0.41 \pm 0.05	66.7	10.3 \pm 4.4
100%	66.7	0.52 \pm 0.05	91.6	6.4 \pm 3.9
Mesophilic				
40%	73.3	0.40 \pm 0.04	83.3	12.6 \pm 6.2
60%	26.7	0.30 \pm 0.08	58.3	1.9 \pm 2.7
80%	0.0	---	91.6	1.3 \pm 4.2
100%	0.0	---	0.0	---
Artificial Rain				
40%	93.3	0.41 \pm 0.08	91.6	19.7 \pm 5.4
60%	96.7	0.45 \pm 0.04	91.6	17.4 \pm 3.0

^aBased on 3 replicates (10 fish per replicate).

^bMean mg/fish, based on 3 replicates (10 fish per replicate). Growth is calculated as change in dry weight over the 7-d test period.

^cTwelve replicates (1 neonate per replicate) were used.

^dFecundity (mean \pm 1 SD) is calculated using only females that survived all 7 d of the test, and their offspring.

^eControl was diluted mineral water.

TABLE 3-3. SUMMARY OF RESULTS FOR CHEMICAL ANALYSES^a OF CONTROL WATER, SYNTHETIC RAINWATER, AND MESOPHILIC AND THERMOPHILIC LEACHATES PERFORMED IN CONJUNCTION WITH TOXICITY TESTS

Sample	pH	Cond. (μ S/cm)	Alkalinity (mg/L)	Hardness (mg/L)	Nitrate (mg N/L)	Phosphate (mg P/L)
Control ^b	7.7	86	30.1	41.1	0.505	0.009
Art. Rain.	5.8	4	1.5	< 2.0	0.026	0.012
Meso. Leach.	7.8	329	78.0	40.0	2.620	7.650
Therm. Leach.	7.9	310	78.0	40.0	0.726	6.025

^aData for pH and conductivity are means for 7 daily measurements; data for alkalinity, hardness, nitrate and phosphate are for single measurements made on the first day of the toxicity tests.

^bDiluted mineral water (1:9, v:v, Perrier and deionized distilled water).

TABLE 3-4. CERIODAPHNIA SURVIVAL AND FECUNDITY IN VARIOUS CONCENTRATIONS OF ACETONITRILE EXTRACTS OF THERMOPHILIC (T) OR MESOPHILIC (M) COMPOSTS CONTAMINATED WITH EXPLOSIVES

Compost type	Final concentration ^a (mg of dried residue per L)	Survival (%)	Fecundity ^b (mean \pm SD)
Control	----	90	20.1 \pm 2.5
M	12	80	0
M	24	0	---
M	36	0	---
M	48	0	---
M	72	0	---
T	12	90	16.8 \pm 2.0
T	24	40	0
T	36	10	0
T	48	0	---
T	72	0	---

^aDried residues of each extract were reconstituted in dimethyl sulfoxide (DMSO) before being tested.

^bCalculated using only females that survived all 7 d, and their offspring.

TABLE 3-5. RESULTS OF CERIODAPHNIA TOXICITY TESTS OF PURE DIMETHYLSULFOXIDE (DMSO), ACETONITRILE EXTRACTS^a OF THERMOPHILIC (T-A) AND MESOPHILIC COMPOST, AND ETHYL ACETATE EXTRACTS OF THERMOPHILIC(T-E) AND MESOPHILIC (M-E) COMPOST.

Sample	Conc. ^b (mg/L)	No. of reps.	No. of animals surviving for 7 d	Mean no. of offspring per female (\pm S)
Control		10	10	21.5 \pm 2.4
DMSO	2 mL/L	10	9	21.7 \pm 4.6
DMSO	5.5 mL/L	10	9	9.6 \pm 3.5
T-A	7.6	10	8	4.3 \pm 2.4
T-A	15.2	10	0	--- \pm ---
T-A	27.8	10	0	--- \pm ---
T-A	50.6	10	0	--- \pm ---
T-A	76.0	10	0	--- \pm ---
M-A	6.0	10	0	--- \pm ---
M-A	12.0	10	0	--- \pm ---
M-A	22.0	10	0	--- \pm ---
M-A	39.9	10	0	--- \pm ---
M-A	60.0	10	0	--- \pm ---
T-E	6.0	10	10	18.7 \pm 2.3
T-E	11.9	10	9	10.1 \pm 3.7
T-E	22.1	10	8	1.8 \pm 1.3
T-E	40.0	10	0	--- \pm ---
T-E	59.9	10	0	--- \pm ---
M-E	6.0	10	10	13.7 \pm 2.7
M-E	11.9	10	4	8.3 \pm 2.1
M-E	22.0	10	0	--- \pm ---
M-E	40.0	10	0	--- \pm ---
M-E	59.9	10	0	--- \pm ---

^aThe dried residues of all solvent extracts were reconstituted into pure DMSO before being tested.

^bConcentration of total extractables.

TABLE 3-6. RESULTS OF FATHEAD MINNOW TOXICITY TESTS OF ACETONITRILE EXTRACTS^a OF MESOPHILIC (M-A) AND THERMOPHILIC (T-A) COMPOSTS

Sample	Conc. ^b (mg/L)	No. of reps.	Mean survival (percent)	Mean growth (mg/fish \pm SD)
Control		3	100.0	0.33 \pm 0.05
T-A	7.6	3	86.7	0.29 \pm 0.03
T-A	15.2	3	76.7	0.24 \pm 0.01
T-A	27.8	3	30.0	0.18 \pm 0.04
T-A	50.6	3	0	--- \pm ---
T-A	76.0	3	0	--- \pm ---
M-A	6.0	3	70.0	0.30 \pm 0.08
M-A	12.0	3	0	--- \pm ---
M-A	22.0	3	0	--- \pm ---
M-A	39.9	3	0	--- \pm ---
M-A	60.0	3	0	--- \pm ---

^aThe dried residues of the solvent extracts were reconstituted into pure DMSO before being tested.

^bConcentration of total extractables.

Data in Table 3-2 showed that the mesophilic leachate was about twice as toxic as the leachate prepared from the thermophilic compost: fecundity of Ceriodaphnia in the 80% concentration of the thermophilic leachate, for example, was similar to fecundity of Ceriodaphnia in the 40% concentration of the mesophilic leachate. The two species used in the toxicity assessments appeared to be about equally sensitive to mesophilic leachate: in each case, this leachate's no-observed-effect concentration was about 40% of full-strength. The synthetic rainwater was not toxic to either species at the highest tested concentration. The full Toxicity Test Reports are included in Appendix A-5.

In conjunction with the toxicity assessments, samples of the CCLT leachates, the synthetic rainwater, and control water were analyzed for alkalinity, hardness, nitrate nitrogen, and total soluble phosphorus using EPA methods. The results of these analyses, plus those for pH and conductivity, are given in Table 3-3.

The greater nitrate content of the mesophilic leachate appears to correlate with its greater explosives content (see Table 2-3).

3.3.3. Tests of Preliminary Acetonitrile Extracts

Acetonitrile extracts of the thermophilic and mesophilic composts were tested for toxicity to Ceriodaphnia only. The dried residues of the two extracts were reconstituted in dimethyl sulfoxide (DMSO) before being tested. Each DMSO preparation contained about 15 mg of dried residue per

TABLE 3-7. SURVIVAL AND FECUNDITY OF CERIODAPHNIA TESTED WITH THE ACID-HYDROLYZED EXTRACTS^a OF THERMOPHILIC (T) AND MESOPHILIC (M) COMPOST

Sample	DMSO extract ^b (μ L/200 mL)	Residue tested (mg/L)	No. of reps.	No. of animals surviving for 7 d	Mean no. of offspring per female (SD) ^c
Neg. control	----	0.00	10	10	19.8 3.8
T	23.5	1.72	10	10	23.0 2.8
T	47.0	3.43	10	10	24.6 3.9
T	94.0	6.86	10	10	23.3 3.1
T	176.0	12.85	10	10	25.5 2.2
T	353.0	25.77	10	10	17.6 3.2 *
M	23.5	2.12	10	9	24.7 1.5
M	47.0	4.24	10	9	24.6 2.2
M	94.0	8.46	10	10	25.1 2.0
M	176.0	15.84	10	10	23.7 3.2
M	353.0	31.77	10	10	17.1 3.8 *
Blank	23.5	0.01	10	10	22.1 1.8
Blank	47.0	0.02	10	10	19.9 2.1 *
Blank	94.0	0.05	10	10	23.2 3.9
Blank	176.0	0.09	10	10	22.0 2.5
Blank	353.0	0.18	10	10	24.2 3.2

^aThe residues of the extracts were reconstituted into DMSO before being tested.

^bConcentrations of dried residue (in mg/mL of DMSO) for the thermophilic and mesophilic samples were 14.6 and 18.0, respectively; the concentration of dried residue in the positive control was 0.1 mg/mL of DMSO.

^cIn each residue test, asterisks show concentrations in which fecundity values were significantly ($p < 0.01$) lower than other concentrations.

mL. Five concentrations of each DMSO solution were tested. These tests included a negative control (i.e., diluted mineral water). The pure DMSO was also tested for toxicity to fathead minnow larvae. Only two concentrations (2.0 and 4.0 ppm DMSO by volume) and a control (no DMSO) were used in this test; the higher concentration was selected because it was equal to the highest concentration of DMSO to which Ceriodaphnia were exposed in the test described above. Survival of the fish in the controls and in the two test solutions >96% in each case. One-way analysis of variance did not reveal significant differences in growth of the fish among the three treatments ($p = 0.64$; $F = 0.47$, d.f. 2,6). The test results that are summarized in Table 3-4 showed that the acetonitrile-soluble residue extracted from the mesophilic compost was about twice as toxic (on a per-unit mass basis) as the residue extracted from the thermophilic compost. These findings were in good general agreement with the relative toxicities for the two CCLT leachates. Additionally, the toxicity of pure DMSO was moderately low for both Ceriodaphnia and the fathead minnows, which justifies its use as a carrier solvent for explosives residues in future tests.

3.3.4. Tests of Final Organic Solvent Extracts

Ceriodaphnia and fathead minnow larvae were both used to quantify the toxicity of the DMSO-reconstituted solutions of acetonitrile and ethyl acetate extracts of both coextractant types with both types of compost. Only fathead minnow larvae were used to test for toxicity of the acetonitrile extracts of the thermophilic and mesophilic composts. The results of the tests with these two species are summarized in Tables 3-5 and 3-6.

For both compost types, the residues from the acetonitrile extracts proved to be considerably more toxic than residues from the corresponding ethyl acetate extracts, based on the results of the Ceriodaphnia tests (Table 3-5). Additionally, for a given type of extractant, residues of the mesophilic compost were more toxic than those of the thermophilic compost (Table 3-5). Finally, a comparison of the results from Tables 3-5 and 3-6 suggested that the Ceriodaphnia were about two to three times more sensitive than the fathead minnows to the acetonitrile-extracted residues of each compost type. The definitive tests of the acetonitrile extracts (Table 3-5) showed that these residues were considerably more toxic than those evaluated in the preliminary tests (Table 3-4). This suggests that an additional toxic species may have been extracted from the compost. The extraction used to produce residues for the definitive tests was performed upon a separate sampling of the composts from the drums. Thus, it is also possible that inhomogeneities in the compost, or changes in the composts themselves, may have contributed to the observed differences.

3.3.5. Tests of The Hydrolyzed Bound Fraction from Composts

In these tests, three samples were evaluated for acute and chronic toxicity to Ceriodaphnia. One was an acid-hydrolyzed extract of the bound fraction from thermophilic compost, one was an acid-hydrolyzed extract of the bound fraction from mesophilic compost, and the third was a blank (i.e., a neutralized sample of the same acidic solution that was used to hydrolyze the residues of the two types of compost). A negative control (i.e., diluted mineral water) was also included with the test. The results of these tests are summarized in Table 3-7.

Survival of Ceriodaphnia in each case was high ($\geq 90\%$). The highest tested concentration of hydrolyzed residue from each compost type slightly depressed Ceriodaphnia reproduction. This reduction was evident based on within-residue statistical analysis (SAS General Linear Models Procedure, followed by a Least Significant Difference test; $p < 0.001$ for both cases, with $F_{4,45} = 9.97$ for the thermophilic material and $F_{4,45} = 13.05$ for the mesophilic material). The overall amount of variance explained by concentration in these analyses, however, was low in both cases (47.0% and 53.7%, respectively, for acid-hydrolyzed residues of the thermophilic and mesophilic composts). Using the same type of analysis, one concentration of positive control residue also reduced

reproduction significantly ($p = 0.006$). However, in this instance the overall percentage of variance explained by concentration was quite small (27.0%).

For interpretative purposes, it is useful to point out that EPA guidelines indicate that a test in which fecundity of the control equals or exceeds 15 offspring per female can be judged to be acceptable. The average fecundity of Ceriodaphnia exceeded 15 even in the highest tested concentration of hydrolyzed residue from each compost type (Table 3-7). Thus, the evidence for toxicity was largely statistical.

In the tests described above, little reduction in fecundity was noted even at concentrations as high as 20-25 mg/L for either hydrolyzed residue. In contrast, acetonitrile extracts of the two compost types tested earlier proved to be strongly toxic at 20-25 mg/L (Tables 3-4, 3-5 and 3-6). Thus, acid hydrolysis did not appear to release toxic amounts of substances that were bound to compost particles. Either the hydrolysis conditions were not strong enough to promote the release of such materials, or the concentration of bound toxic materials available to be released by acidic conditions was very low.

3.4. Conclusions

The overall conclusions to be drawn from the results of the toxicity tests described above are as follows: First, the CCLT leachates of the thermophilic compost show very little evidence for acute or chronic toxicity to either Ceriodaphnia or fathead minnows, but CCLT leachates of mesophilic compost were moderately toxic (Tables 3-1, 3-2). There was also not much evidence for toxicity associated with bound constituents for either compost type (cf. Table 3-7). Strong organic solvent (i.e., acetonitrile, ethyl acetate) extracts of both compost types were quite toxic, but those from the thermophilic compost were less toxic than those from the mesophilic compost. The toxicity of acetonitrile extracts were also greater than those of ethyl acetate. This pattern was concordant with the notion that most of the toxicity in the reconstituted residues of the solvent extracts could be accounted for by explosives and/or their degradation products. Thus, the EPA-approved toxicity tests conducted with Ceriodaphnia and fathead minnow larvae suggest that composting could be an effective means to lower toxicity, as well as the concentration, of high explosives such as TNT, RDX, and HMX.

The approximate toxic threshold for each type of leachate for both compost types was estimated from their capacity to reduce fecundity of Ceriodaphnia. These values, expressed as the concentration of residue (in mg/L) needed to show evidence of toxicity, could be determined only as 'less than' the lowest tested concentration (e.g., in the case of acetonitrile extracts of both compost types; Table 3-4). The toxicity threshold concentrations were then converted to toxic units (TUs; the reciprocal of the threshold toxicity concentration, expressed as g/L). The TUs, in turn, were used to estimate the intrinsic toxicity of each compost sample by correcting for (a) the amount of solvent (CCLT solution, acetonitrile, or ethyl acetate) used to prepare the extract, and (b) the mass of compost extracted. The results of these calculations are shown in Table 3-8.

The information shown in Table 3-4 shows that the mesophilic compost was 2 to 3 times more toxic than the thermophilic compost. However, within a particular type of compost, there appeared to be little difference in toxicity that could be attributed to the type of solvent. Thus, the CCLT procedure for estimating compost toxicity seems to provide toxicity estimates as good as those obtained using stronger solvents, and thus there does not appear to be any toxicity that the CCLT leaching procedure misses.

TABLE 3-8. ESTIMATED TOXICITY (TOXIC UNITS/GRAM) OF THERMOPHILIC AND MESOPHILIC COMPOST BASED ON RESULTS OF CERIODAPHNIA TESTS OF EXTRACTED MATERIAL OBTAINED USING AQUEOUS (CCLT) AND ORGANIC-SOLVENT (ACETONITRILE AND ETHYL ACETATE) EXTRACTANTS

Compost type	Solvent type	Toxic units/gram
Thermophilic	CCLT (aqueous)	0.03
Mesophilic	CCLT (aqueous)	0.11
Thermophilic	ethyl acetate	0.05
Mesophilic	ethyl acetate	0.10
Thermophilic	acetonitrile	> 0.04
Mesophilic	acetonitrile	> 0.05

CHAPTER 4. AMES TESTS

4.1. Introduction

In 1975 Bruce Ames described a technique (21) utilizing bacteria as a screening device for chemical mutagens. Test data on more than 5000 chemicals have been described since the original publication, establishing the Ames test as a major assay in detection of mutagenicity. The test measures back-mutation to histidine independence of mutant strains in the *his* operon of *Salmonella typhimurium* that can be reverted by base-pair substitutions (TA-100) or frameshift mutations (TA-98). These are the two most widely-used tester strains for mutagenicity assay. By inoculating a histidine deficient medium with the tester strains and various concentrations of the test material, revertant (mutated) colonies can be counted and the mutagenicity of the test material quantitated. Considering the universal acceptance of the Ames test and the need to test the toxicological characteristics of products resulting from composting of explosives wastes we undertook extensive Ames testing of the aqueous leachates and organic solvent extracts of the mesophilic and thermophilic composts.

4.2. Materials and Methods

The Ames assay is performed by mixing the following components in a sterile test tube placed in a 43°C heating block.

- (a) 2.0 mL of 0.6% agar containing 0.6% NaCl, 0.05 mM biotin, and 0.05 mM histidine,
- (b) 0.1 mL of freshly grown indicator organisms (10^9 organisms/mL) TA-98 or TA-100, and (if metabolic activation is used) 0.5 mL of S-9 solution,
- (c) appropriate amount of test material.

The contents of the tube are gently vortexed and poured and evenly distributed over the entire surface of a minimal glucose agar plate. Duplicate plates are run for each dilution of test substance. The top agar is allowed to solidify and then is incubated for 48 hours at 37°C. The number of *his*⁺ revertant colonies can then be counted. Spontaneous, negative (solvent), and positive controls (sodium azide and nitrofluorene) are run concurrently with each experiment.

The top agar is made by adding 6 g of Difco-Bacto agar and 5 g NaCl to 1 liter of distilled water, which is autoclaved and stored in 100-mL bottles at room temperature. Before use, the agar is melted (in an autoclave or microwave) and 10 mL of a sterile solution of 0.5 mM L-histidine-HCl, 0.5 mM biotin is added to the 100 mL of molten agar and mixed thoroughly.

Vogel-Bonner medium E with 2 percent glucose and 1.5 percent Bacto-Difco agar is used as the minimal medium and is prepared as follows:

Vogel-Bonner Salts (50X)	
Warm Distilled Water	670 mL
Magnesium Sulfate ($\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$)	10 g
Citric Acid Monohydrate	100 g
Potassium Hydrogen Phosphate (K_2HPO_4)	500 g
Sodium Ammonium Hydrogen Phosphate ($\text{NaH}_2\text{NH}_4\text{PO}_4 \cdot 4\text{H}_2\text{O}$)	175 g

The above salts are added to the warm water (45°C) in the specified order. Each salt is completely dissolved before the next is added. When the salts are all dissolved, the solution is cooled to room temperature. About 5 mL of chloroform is added to the solution and stored in a capped bottle at room temperature.

Dissolve 15 g of Difco-Bacto agar in 1 liter of water by autoclaving. Cool to about 60 to 70°C and add 20 mL of 50X V-B salt solution and 50 mL of sterile 40 percent glucose solution. Mix thoroughly and dispense into 100 x 15 mm plastic petri dishes (25 mL per plate).

The indicator *Salmonella* strains are kept frozen in nutrient broth supplemented with 10% sterile glycerol at -80°C in 1 mL aliquots, each of which contains about 10⁹ cells. For each experiment, the 1 mL frozen aliquots are allowed to thaw at room temperature and inoculated into 30 ml of nutrient broth. The cultures are grown at 37°C, unshaken for 4 hours, then gently shaken (100 rpm) for 11 to 14 hours. Histidine dependency is checked for each strain whenever experiments are performed.

The S-9 preparation was a rat liver S-9 with Aroclor activation, obtained from Litton Bionetics (Oklahoma City, OK). It was diluted 0.04 ml to 0.5 mL with salt solution before addition with the tester strains.

4.3. Observations

Preparatory to testing of compost extracts for mutagenic activity, the genotypes of the *Salmonella* tester strains were confirmed as was their response to known mutagens, known explosives, and some explosive derivatives.

4.3.1. Confirmation of Strains

The Bacterial tester strains TA-98 and TA-100 used in the test procedure have mutations in the *rfa* and *uvrB* genes. They also contain the R-factor plasmid pKM101. The genotypes of the tester strains were confirmed by evaluating their sensitivity to crystal violet and to UV light and to resistance to ampicillin. Both strains were killed by exposure to crystal violet and UV irradiation but were unharmed by ampicillin, thus confirming their genotype.

The known mutagens, nitrofluorene, acetylaminofluorene, and sodium azide, were then tested for their mutagenic effects on the tester strains with and without metabolic activation (rat liver microsomal fraction S-9). The effects of the known mutagens are shown in Table 4-1. As with the crystal violet, UV light and ampicillin, the tester strains showed good response to the known mutagens. Having verified the genomic integrity and the expected reversion in response to known mutagens, tests of known explosives and their derivatives were undertaken and compared with extracts of the composted explosives.

4.3.2. Tests of Explosive Compounds and TNT Metabolites

We initially tested four of the explosives known or suspected to be present in the compost piles, i.e., TNT, tetryl, RDX, and HMX. Only TNT and tetryl were found to be mutagenic, with the latter more mutagenic and toxic than the former (Appendix Table 6A-1). The addition of a rat liver metabolic activation system had no effect on the mutagenicity of the two compounds. Based on the slope of the plot of the number of revertants induced versus mg of test compound, it is estimated that for TA-98, tetryl induced 8,200 revertants per mg (rev/mg) and for TA-100 the value is 23,000 rev/mg. The corresponding values for TNT are 5,400 and 6,600 rev/mg, respectively.

TABLE 4-1. RESULTS FROM AMES TESTING OF KNOWN MUTAGENS

Sample	TA98 ^c revertants/plate		TA100 ^c revertants/plate	
	-S9	+S9	-S9	+S9
1. Control	24	28	117	111
2. Nitrofluorene ^a	874	NT	NT	NT
3. Acetylaminofluorene ^a	NT	533	NT	227
4. Sodium Azide ^a	NT ^b	NT	586	NT

^aPositive controls. Nitrofluorene (10 µg/plate). Acetylaminofluorene (10 µg/plate). Sodium azide (2 µg/plate).

^bNT - not tested

^c*Salmonella* overnight culture - 2×10^9 cells/ml. Ampicillin resistant, UV sensitive, crystal violet sensitive.

In addition to TNT and tetryl, we also tested biotransformation products of TNT. The results are summarized in Table 4-2 and are detailed in Appendix Table 6A-2. On a weight basis, TNT was more mutagenic than the two monoamino metabolites for both strains of bacteria. The relative sensitivity of the TA-98 strain to TNT and its metabolites was TNT > 4-A-2,6-DNT > 2-A-4,6-DNT, while for TA-100 the relative sensitivity was TNT > 2-A-4,6-DNT > 4-A-2,6-DNT. HMX and RDX do not show detectable mutagenicity, but tetryl was the most mutagenic of the explosives and metabolites tested.

Finally, two other diaminonitrotoluenes were tested to complete our investigation into the potential changes in toxicity associated with biotransformation of TNT. The specific mutagenic activities for 2,4-DA-6-NT and 2,6-DA-4-NT) are included in Table 4-2, and the data are listed in Table 6A-3. It is apparent that the mutagenicity was diminished as nitro groups were reduced to amino groups, and the activity is highly isomer-specific. The 2,4-DA-6-NT does not exhibit detectable mutagenicity, while 2,6-DA-4-NT is approximately one-tenth as mutagenic as TNT.

4.3.3. Tests of CCLT Leachates and Suspended Particles

Having confirmed the genome and mutagenic reactivity of the *Salmonella* tester strains and their reaction to explosives and their biotransformation products known or suspected to be in the test compost, we began analysis of the leachates. Testing of the preliminary CCLT leachates of the mesophilic and thermophilic composts showed little mutagenic activity (Appendix Table 6A-4). Subsequently, we retested larger volumes of fresh CCLT leachates from the final leaching of the mesophilic and thermophilic composts and found similar results. Only the mesophilic compost leachate tested in bacterial strain TA-98 showed a positive, albeit slight mutagenic response up to the maximum allowable dose in the test protocol (see Table 6A-5). The test was performed without the addition of a rat liver metabolic activation system (S-9). The response for the mesophilic

TABLE 4-2. SUMMARY OF AMES MUTAGENICITY TESTS FOR EXPLOSIVES AND METABOLITES

Compound	Specific Activity ^a , rev/mg	
	TA-98	TA-100
HMX	ND	ND
RDX	ND	ND
Tetryl	8,200	23,000
TNT	5,400	6,600
4-A-2,6-DNT	1,000	500
2-A-4,6-DNT	500	2,300
2,4-DA-6-NT	ND	ND
2,6-DA-4,NT	325	750

^aWithout S-9 metabolic activation

leachate, ca. 59 rev/mL with the TA-98 strain, was only slightly above background. This corresponds to ca. 60 rev/mg of extractable matter and 1,200 rev/g of leached compost.

The mutagenicity calculated for the CCLT leachate from the concentrations of the explosive compounds and TNT metabolites and their specific mutagenicities mathematically accounts for 37% of the observed mutagenicity (TA-98 strain), with TNT constituting 36% of the activity. The observed TNT metabolites accounted for ca. 1%. Although this suggests that removal of TNT would effectively detoxify the compost and leachates, 37% of the observed activity cannot be attributed to TNT and its metabolites because there was no blank compost without explosives. In addition, this mathematical exercise does not take into account potential synergistic and antagonistic interactions among and with the explosives compounds and their metabolites. The presence of other mutagens in the CCLT leachate is suggested by its greater response with the TA-98 strain while the pure explosives and TNT metabolites generally caused a greater response with the TA-100 strain (Table 4-2).

Because the CCLT leachates did not elicit a clear mutagenic response even when tested at the maximum dose levels tolerated by the assay system, we concentrated the CCLT leachates 100-fold by flash evaporating 100 mL to dryness and then extracting the residue with 1 mL of DMSO. The results for these mutagenicity tests are summarized in Table 4-3, and the full data set is included in Table 6A-6. The specific activity in revertants per mL, calculated per mL of original CCLT leachate (i.e., corrected for the concentration factor) was approximately 700 and 150 for the mesophilic

leachate in the TA-98 and TA-100 strains (respectively), and 170 and 40 (respectively) for the thermophilic leachate. Corresponding calculations for revertants per mg of extracted matter and revertants per g of leached compost also are shown in Table 4-3. In all cases, the TA-98 strain was more sensitive than TA-100 to the mutagens in the extracts. The mesophilic leachate is 2 to 4-fold more mutagenic than the thermophilic leachate, which was in agreement with the

TABLE 4-3. SUMMARY OF RESULTS FOR AMES MUTAGENICITY TEST OF THE DMSO CONCENTRATE OF THE MESOPHILIC AND THERMOPHILIC COMPOST CCLT LEACHATES

Sample	Specific Activity ^a , rev.			
	TA-98		TA-100	
	Per mg of Extractables	Per g of Compost	Per mg of Extractables	Per g of Compost
DMSO Concentrates of Dried CCLT Leachates:				
Mesophilic Compost	760	14,000	160	3,000
Thermophilic Compost	150	3,400	34	800

^aWithout S-9 metabolic activation

Ceriodaphnia and fathead minnow larvae tests. Also, the specific activities were ca. 10-fold higher than those measured on the original leachates. Two explanations are possible. Because the leachate was concentrated 100-fold for these tests, much greater doses were applied to the plates and it is possible that different regions in the dose-response curve are being compared. In general, the overall mutagenicity of a complex mixture (such as compost leachates) depends upon the different mutagens present in the mixture. Because the mutagenic response of each individual mutagen is different at different dose ranges, the comparison of the overall mutagenicity data of complex mixtures at various dose ranges is easily misleading.

It also is possible that these higher activities were contributed by particle-associated mutagens which the DMSO extracted from the dried residue. Particles were filtered out of the CCLT leachates before the direct Ames testing, because of the need to remove interfering bacteria. Subsequently another CCLT leachate of mesophilic compost was prepared and tested before and after centrifuging to remove the particles (section 2.5). The particles resulting from centrifugation were extracted with DMSO and the DMSO extract also was tested. The results of these tests are shown in Table 4-4 (full data set in Table 6A-7). Although the experiment confirms the association of mutagenicity with the particulate fraction, the total leachate activity is ca. 10-fold greater than that of the particles alone. This experiment suggests that the particles do not harbor large amounts of toxicity unavailable to the toxicity tests. Because the mutagenicity of the leachate is very low and slightly above background, we do not feel that the apparent differences between the uncentrifuged and centrifuged leachate samples are significant. The differences between the absolute activities determined for this small-volume leachate (4 L of leachate from 200 g of compost) and the larger generation of leachate (28 L generated from 1400 g of compost) discussed above are attributed to compost inhomogeneity (see section 2.2.5.).

TABLE 4-4. AMES TEST OF MESOPHILIC CCLT LEACHATES^a, BEFORE AND AFTER CENTRIFUGATION, AND LEACHATE SOLIDS EXTRACTED WITH DMSO

Sample	Specific Activity ^a , rev.			
	TA-98		TA-100	
Mesophilic CCLT Leachates:	Per mL of Compost Leachate ^c	Per g of Compost ^d	Per mL of Compost Leachate	Per g of Compost ^d
Before Centrifugation	400	8,000	300	6,000
After Centrifugation	200	4,000	100	2,000
Leachate Solids	35	700	15	300
Mesophilic CCLT ^e	59	1,200	ND	ND

^a4 L of Mesophilic CCLT Leachate prepared from 200 g of compost

^bWithout S-9 metabolic activation

^cActivity was calculated based on 1 mL of compost leachate

^dActivity was calculated based on 1 g of compost

^eTest of final CCLT leachate (pg. 42 for comparison)

4.3.4. Tests of Organic Solvent Extracts

Preliminary acetonitrile extracts of the composts were tested for rangefinding purposes. The acetonitrile extracts were dried and reconstituted with DMSO. The results of Ames testing are summarized in Table 4-5 and are fully listed in Table 6A-8. On the basis of revertants per mg of extractable matter or per g of compost extracted, the specific activities are greater than those measured directly upon the CCLT leachates, but are comparable with those measured upon the concentrated CCLT leachate residues. This undoubtedly is due partly to the better solubilities of mutagens in acetonitrile versus the acidified water in the CCLT and the great dilution of mutagens in the CCLT. It is also possible that the different sample vehicles (DMSO versus acidified water) applied to the plates contributed to the differences in observed mutagenic activity, or that inhibitors present in the aqueous leachates were not extracted with the organic solvents.

In this experiment, unlike the CCLT leachates, the mesophilic compost extract was only slightly more mutagenic than was the thermophilic compost extract. As with the CCLT leachates, however, the TA-98 strain was more sensitive than TA-100. Only the mesophilic compost extract was tested with the S-9 rat liver metabolic activation system, and no increase in activity was observed (data not shown). Comparison of the actual mutagenicity with that predicted from the concentrations and specific mutagenicities of TNT and its metabolites mathematically accounted for 73% of the actual activity found with the TA-100 strain for the extract of the mesophilic compost and only 19% for the thermophilic compost. The mathematical accounting for the activity found using the TA-100 strain was about two-thirds less. Again, this activity cannot be confidently assigned to TNT and its metabolites.

TABLE 4-5. AMES TEST OF PRELIMINARY ACETONITRILE EXTRACTS OF COMPOST

Compost	Specific Activity ^a , rev.			
	TA-98		TA-100	
	Per mg of Extractables	Per g of Compost	Per mg of Extractables	Per g of Compost
Mesophilic	3,400	11,200	1,300	4,300
Thermophilic	2,500	4,500	1,000	1,800

^aWithout S-9 Activation

In two subsequent experiments both acetonitrile and ethyl acetate were tested for their ability to extract mutagens from mesophilic and thermophilic composts. The acetonitrile and ethyl acetate extracts of the composts were tested for bacterial mutagenicity with strains TA-98 and TA-100. Activation with rat liver S-9 was not included because our previous experience with both aqueous leachates and with the preliminary acetonitrile extracts showed no enhancement of mutagenicity, indicating the presence of direct-acting mutagens. The data for the extracts are summarized in Table 4-6, and the full data sets are listed in Appendix Tables 6A-9 - 6A-11. As before, the organic solvent extracts exhibited considerable mutagenicity. It is interesting to note that the mutagenic activities found in the acetonitrile and ethyl acetate extracts were comparable with each other and also there was no significant difference between the extracts from the mesophilic and thermophilic composts. The former observation suggests that all available mutagenicity has been extracted. The reason for the latter is not clear, but may be related to compost inhomogeneity.

4.3.5. Tests of Hydrolyzed Bound Fraction

The hydrolyzed bound fraction sample (see section 2.4) was tested to determine if the hydrolysis conditions liberated mutagenic compounds. The data listed in Appendix Table 6A-12 show that no mutagenicity was detected in the acid-hydrolyzed bound fraction of either compost. At this stage it cannot be concluded that hydrolyzable mutagens are absent because the hydrolysis conditions were subsequently found (see section 2.4) to be too severe for the survival of mutagenic 4-A-2,6-DNT and 2,4-DA-6-NT. The latter are potential hydrolysis products of a bound fraction with acid-labile bonds. More gentle hydrolysis conditions are needed.

4.4 Discussion and Conclusions

Not surprisingly, tetryl, TNT and some known TNT derivatives shown or thought to be present in the compost were found to be mutagenic. It is important to note that the mutagenicity of TNT is diminished as nitro groups are reduced to amino groups, and that the activity is highly isomer-specific. In general, greater biotransformation results in lesser mutagenicity. The CCLT leachates

TABLE 4-6. AMES TEST RESULT SUMMARY FOR FINAL ORGANIC SOLVENT EXTRACTS OF COMPOSTS

Compost/solvent ^a	Specific Activity ^a , rev.			
	TA-98		TA-100	
	Per mg of Extractables	Per g of Compost	Per mg of Extractables	Per g of Compost
Mesophilic/MeCN	6,400	19,000	3,600	10,600
Mesophilic/EtoAc	4,300	20,000	2,200	7,400
Thermophilic/MeCN	10,000	22,400	2,500	5,900
Thermophilic/EtoAc	6,600 ^c	16,800 ^c	5,100	13,000

^aMeCN = Acetonitrile, EtoAc = EthylAcetate

^bAverage value based on duplicate data

^cSingle value

of mesophilic but not thermophilic compost showed mutagenic activity some of which could be attributed to TNT. Comparing the predicted activities of the extracts or leachates of the composts (calculated from the concentrations of the explosives or metabolites and their specific mutagenicities) with the actual activities would suggest that TNT contributes substantially to their activities. For example, ca. 37% of the mutagenicity of the mesophilic compost CCLT leachate is mathematically accounted by TNT alone. However, it is premature to attribute 37% of the activity to TNT because of potential synergistic or antagonistic effects, and the possible presence of other mutagens which have not been identified. While CCLT extraction originally indicated mutagenic activity was found only in the mesophilic compost, the extractions with organic solvents indicated mutagenic activity remaining in both mesophilic and thermophilic composts. The mutagenic activity determined in the 100-fold concentrated CCLT residue also showed that the high dilution of the CCLT makes detection of mutagenicity difficult. In the absence of the starting material (i.e., contaminated sediment) for comparison, the degree to which composting reduced the mutagenic activity of the starting material could not be ascertained. The lack of a compost blank (from sediment without explosives) prevents assigning any of the observed activity to explosives or their biotransformation products.

To place these findings in perspective, Table 4-7 compares the mutagenicities of the mesophilic compost extracts and leachates with those of a coal combustion stack ash extract, a petroleum crude oil, crude shale oil, a coal-derived fuel oil blend, and pure TNT. The compost is more mutagenic than the stack ash, comparable to the petroleum crude oil, and much less active than the shale oil and coal-derived fuel oil blend and pure TNT. Although the oils and fuels required S-9 metabolic activation for the mutagenicity test (and therefore represent predominantly direct-acting mutagens) and the data for the compost leachates or extracts were obtained without metabolic activation, the incorporation of S-9 into the test for the compost did not increase the activity.

TABLE 4-7. COMPARISON OF MUTAGENIC ACTIVITIES

Material	Mutagenic Activity with Strain TA-98, rev/g ^a
CCLT Leachate of Mesophilic Compost	1,200
100-Fold DMSO Condensate of CCLT Leachate of Mesophilic Compost	14,000
Acetonitrile Extract of Mesophilic Compost	19,000
Methylene Chloride Extract of Coal Combustion Stack Ash ^b	61
Wilmington Crude Oil ^c	5,000 ^d
Paraho-SOHIO Crude Shale Oil ^c	390,000 ^d
SRC-II coal-Derived Fuel Oil Blend ^c	1,100,000 ^d
TNT	5,400,000

^aData are for TA-98 without S-9 metabolic activation unless otherwise indicated.

^bReference 22. ^cReference 23. ^dS-9 metabolic activation used.

CHAPTER 5.0 CONCLUSIONS

The major conclusions from this characterization of the mesophilic and thermophilic composts from the LAAP field composting experiment are as follows:

- (a) The major explosives-related compounds determined in the leachates and organic solvent extracts of the composts were HMX, RDX, TNT, and two TNT biotransformation products, 2-A-4,6-DNT, and 4-A-2,6-DNT.
- (b) Compost heterogeneity considerably complicates chemical analysis and toxicity testing, and requires that relatively large masses of sample or large numbers of samples be tested.
- (c) The composted product of explosives processing wastes contains low levels of leachable toxic compounds. The leachates from the mesophilic compost were slightly more toxic than those from the thermophilic compost. The toxicity cannot be attributed to explosives or their biotransformation products without tests of a suitable blank compost. Reduction in toxicity of explosives processing wastes by composting cannot be proved without tests of the starting material. The effects of the 10 month interval between composting and characterization are unknown.
- (d) The CCLT does not appear to miss large reservoirs of aquatic organism toxicity or bacterial mutagenicity which are available to organic solvent extraction, but the high dilution of the leached material in the CCLT protocol makes determination of mutagenic activity difficult without preconcentration of the CCLT leachate.
- (e) Particles which pass with the CCLT leachate through the EPA pressure filtering protocol do not appear to harbor significant amounts of organic solvent extractable mutagenicity.
- (f) Liberation of a bound fraction of explosives transformation products will require a gentler hydrolysis than refluxing 6N hydrochloric acid to allow potential mutagens to survive.

ACKNOWLEDGEMENTS

This work would not have been possible without the assistance and input of many people. The authors gratefully acknowledge J. E. Caton, W. M. Caidwell, G. S. Fleming, R. M. Edwards, and E. T. Maestas of the Analytical Chemistry Division, L. A. Kszos, L. F. Wicker, P. W. Braden, R. D. Bailey, G. J. Haynes, G. W. Morris, J. Richmond, and L. S. Ewald of the Environmental Sciences Division, and K. S. Ironside of the Health and Safety Research Division.

REFERENCES

1. Bartell, R. Introduction/Overview 1989. In Proceeding of the Workshop on Composting of Explosives Contaminated Soils. USATHAMA Report No. CETHA-TS-SR-892276, p.D-2, D-3 U.S. Army Toxic and Hazardous Materials Agency, Edgewood, MD.
2. Isbister, J. D., R. C. Doyle, and J. K. Kitchens. 1982. Composting of Explosives. U. S. Army Report DRXTH-TE: Atlantic Research Corporation, Alexandria, VA.
3. Doyle, R. C., J. D. Isbister, G. L. Anspach, and J. F. Kitchens. 1986. Composting Explosives/Organics-Contaminated Soils. U. S. Army Report AMXTH-TE-CR-86077. Atlantic Research Corporation, Alexandria, VA.
4. Williams, R. T., P. S. Ziegenfuss, and P. J. Marks. 1988. Task Order -8 Field Demonstration - Composting of Explosives-Contaminated Sediments at the Louisiana Army ammunition Plant (LAAP). U. S. Army Report AMXTH-IR-TE-88242.
5. Synthetic Precipitation Leach Test (also referred to as the Clean Closure Leach Test), SW-846 Draft Method 1312, obtained from the U. S. EPA Office of Solid Waste.
6. Test Methods for Evaluating Solid Waste, SW-846, U. S. Environmental Protection Agency, Office Of Solid Waste Emergency Response, Washington, D. C. November, 1986.
7. U. S. EPA Contract Laboratory Program, Statement of Work for Organic Analysis, Multi-Media, Multi-Concentration, SOW No. 787, Rev: 1/87,2/87, 7/87, Contract IFB WA-87K236, IFB WA-87K237, and IFB WA-87K238.
8. Jenkins, T. F., and C., L. Grant, 1987. Comparison of Extraction Techniques for Munitions Residues in Soil. Anal. Chem. 59:1326-1331.
9. Jenkins, T. F., and M. E. Walsh. 1987. Development of an Analytical Method for Explosives Residues in Soil. CRREL Report No. 87-7, U. S. Army Cold Regions Research and Engineering Laboratory, Hanover, NH.
10. Toxicity Characteristic Leaching Procedure, SW-846 Method 1310.
11. Cheshire, M. V., P. A. Cranwell, C. P. Falshaw, A. J. Floyd, and R. D. Haworth. 1967. Humic Acid -II. Structure of Humic Acids. Tetrahedron. 23: 1669-1682.
12. Burton, G. A., D. Nimmo, D. Murphey and F. Payne. 1987. Stream Profile Determinations using Microbial Activity Assays and Ceriodaphnia. Environ. Toxicol. Chem. 6:505-513.
13. Nimmo, D. W. R., D. Link, L. P. Parrish, G. J. Rodriguez, W. Wuerthele, and P. H. Davies. 1989. Comparison of On-Site and Laboratory Toxicity Tests: Derivation of Site-Specific Criteria for Un-Ionized Ammonia in a Colorado Transitional Stream. Environ. Toxicol. Chem. 8:1177-1189.
14. Stewart, A. J., L. A. Kszos, B. C. Harvey, L. F. Wicker, G. J. Haynes and R. D. Bailey. 1990. Ambient Toxicity Dynamics: Assessments using Ceriodaphnia dubia and Fathead Minnow (Pimephales promelas) Larvae in Short-Term Tests. Environ. Toxicol. Chem. 9: (in press).

REFERENCES (Cont'd)

15. Taylor, P. A., K. Zabelsky, R. H. Kingrea, and A. J. Stewart. 1987. Optimization of a Biological/Chemical Wastewater Treatment Process using Biototoxicity Tests and Chemical Analysis. Y/DZ-260. Y-12 Plant, Oak Ridge, TN.
16. Taylor, P. A., A. J. Stewart, and L. Holt. 1988. Toxicity of Common Salts to Three Biototoxicity Test Organisms. Y/DZ-420, Y-12 Plant, Oak Ridge, TN.
17. Horning, W. B., II and C. I. Weber. 1985. Short-Term Methods for Estimating the Chronic Toxicity of Effluents and Receiving Waters to Freshwater Organisms. EPA/600/4-85/014. U.S. Environmental Protection Agency Environmental Monitoring and Support Laboratory, Cincinnati, OH.
18. Kszos, L. A., A. J. Stewart, L. F. Wicker, and G. M. Logsdon. 1989. Environmental Sciences Division Toxicology Laboratory Standard Operating Procedures. ORNL/TM-11194, Oak Ridge National Laboratory, Oak Ridge, TN.
19. Loar, J. M., Ed. 1990. Environmental Sciences Division. Third Annual Report on the ORNL Biological Monitoring and Abatement Program. ORNL/TM. Oak Ridge National Laboratory, Oak Ridge, TN.
20. Kszos, L. A. and A. J. Stewart. 1990. Effort-Allocation Analysis of the 7-d Fathead Minnow (Pimephales promelas) and Ceriodaphnia dubia Toxicity Tests. Environ. Toxicol. Chem. (submitted).
21. Ames, B. N., J. McCann, and E. Yamasaki. 1975. Methods for Detecting Carcinogens and Mutagens in the Salmonella/Mammalian-Microsome Mutagenicity Test. Mutation Research. 31: 347-364.
22. Griest, W. H. , B. A. Tomkins, and J. E. Caton. 1986. Development of Methods for Analyzing Organics in Fly Ash. EA-4792. The Electric Power Research Institute, Palo Alto, CA.
23. Guerin, M. R., I. B. Rubin, T. K. Rao, B. R. Clark, and J. L. Epler. 1981. Distribution of Mutagenic Activity in Petroleum and Petroleum Substitutes. Fuel 60: 282-289.

APPENDIX A-1
LIST OF ABBREVIATIONS

List of Abbreviations

<u>Abbreviation</u>	<u>Full Name</u>
Explosive Compounds and TNT Metabolites:	
2,6-DA-4-DT	2,6-Diamino-4-nitrotoluene
2,4-DA-6-DT	2,4-Diamino-6-nitrotoluene
2,4,6-TNBAIc	2,4,6-Trinitrobenzylalcohol
RDX	Hexahydro-1,3,5-trinitro-1,3,5-triazine
HMX	Octahydro-1,3,5,7-tetranitro-1,3,5,7-tetrazocine
1,3,5-TNB	1,3,5-Trinitrobenzene
TNB	1,3,5-Trinitrobenzene
1,3-DNB	1,3-Dinitrobenzene
2-A-4,6-DNT	2-Amino-4,6-dinitrotoluene
4-A-2,6-DNT	4-Amino-2,6-dinitrotoluene
2,6-DNT	2,6-Dinitrotoluene
2,4-DNT	2,4-Dinitrotoluene
TNT	1,3,5-Trinitrotoluene
Tetryl	N-methyl-N,2,4,6-Tetranitroaniline
4-OHA-2,6-DNT	4-Hydroxyamino-2,6-dinitrotoluene
Azoxydimer	2,2',6,6'-tetranitro-4,4'-azoxytoluene
Others:	
CCLT	Clean Closure Leach Test
CLP	EPA Contract Laboratory Program
EPA	U.S. Environmental Protection Agency
GC-MS	Gas chromatography-mass spectrometry
ICP	Inductively coupled plasma emission spectroscopy
L	Liter
m	Milli-
SVOA	Semivolatile organic compound analysis
TCLP	Toxicity characteristic leaching procedure
VOA	Volatile organic compound analysis
μ	Micro-

APPENDIX A-2
HPLC METHOD FOR EXPLOSIVES AND TNT METABOLITES

HPLC Analysis of Explosive Compounds and TNT Metabolites

Reverse Phase Separation on Octadecylsilane Column

The aqueous leachates were analyzed using a reverse phase separation on an octyldecylsilane (ODS) phase. This procedure is capable of resolving TNT, tetryl, HMX, RDX, monoaminodinitrotoluenes, diaminonitrotoluenes, trinitrobenzyl alcohol, and the azoxydimer. However, the dinitrotoluenes are resolved with difficulty, mono and dinitrobenzene are not separated, and the trinitrobenzoic acid elutes very early and is difficult to separate from the solvent peak.

A 250 μ L sample of aqueous leachate or standard is injected onto a 150 X 4.6 mm Zorbax ODS column and then eluted at a flow rate of 1 mL/min. with the following solvent program: 0 to 8 min., 20% aqueous acetonitrile; 8 to 18 min., an aqueous acetonitrile gradient increasing in acetonitrile concentration from 20% to 30%; 18 to 38 min., an aqueous acetonitrile gradient from 30% acetonitrile to 45% acetonitrile; 38 to 50 min., an aqueous acetonitrile gradient from 45% acetonitrile to 75% acetonitrile; 50 to 60 min., 75% aqueous acetonitrile; 60 to 70 min., a reverse acetonitrile gradient going from 75% acetonitrile to 20% acetonitrile; and 70 to 80 min., equilibration of the column with 20% acetonitrile in preparation for the next injection. This method yields the chromatogram shown in Fig. A2-1. Several chromatographic parameters are summarized in Table A2-1. Abbreviations used in the figure are listed in Appendix A-1. Although the total run time is 80 min., the *k*-values listed in Table A2-1 indicate good separations and the peak shape is quite good as indicated by the peak asymmetry. Further examination of this data does indicate that 1,3-dinitrobenzene and 1,3,5-trinitrobenzene are not separated adequately for individual quantitation. In addition 2,4-dinitrotoluene and 2,6-dinitrotoluene are not adequately separated in Figure 2A-1; however, as indicated by the *k*-values in Table 2A-1, the dinitrotoluenes can be separately quantitated under more ideal conditions, (new column). The one potential metabolite that can not be determined with this procedure is trinitrobenzoic acid which is essentially not retained by the column, (*k*=0.23). Even a much weaker eluent, (10% or less acetonitrile) will not cause trinitrobenzoic to be significantly retained.

The method described above yields very good results for most metabolites even though a sample that is 10 to 100 times the ideal sample volume is injected. Sensitivities for clean standards range from approximately 0.042 ppm in the injected solution (corresponding to 11 ng injected) for 2-amino-4,6-dinitrotoluene to 0.53 ppm (133 ng) for 4-hydroxyamino-2,6-dinitrotoluene using a 250 μ L injection volume and a diode array detector and a wavelength of 254 nm for quantitation. The typical sensitivity is represented by the 0.064 ppm (16 ng) for TNT. Quantitation is achieved by external standardization with 9 concentrations of authentic standards ranging from 0.1 to 4 μ g/mL. Peak areas are used for calculations and the spectra of the peaks are examined to confirm identifications.

FIGURE 2-A1. SEPARATION OF EXPLOSIVE COMPOUNDS AND TNT METABOLITES ON OCTADECYLSILANE REVERSE PHASE COLUMN. SAMPLE WAS 250 μ L OF A 1 PPM STANDARD MIXTURE.

ORNL-DWG 90-2207

REVERSE PHASE HPLC OF STANDARDS




Table 2A-1. Chromatographic Parameters for Explosives and TNT Metabolites on Zorbax Column with Acetonitrile Gradient

<u>COMPOUND</u>	<u>RET. TIME, MIN</u>	<u>k¹</u>	<u>WIDTH²</u>
1,3,5-TNBA	1.61	0.23	0.19
2,4-DA-6-NT	6.93	4.28	0.28
2,6-DA-4NT	7.87	4.99	0.30
TNBAIc	18.57	13.1	0.36
RDX	19.59	13.9	0.33
HMX	21.34	15.2	0.35
1,3-DNB	24.82	17.90	0.35
1,3,5-TNB	24.94	17.99	0.35
2-A-4,6-DNT	30.11	21.9	0.32
4-A-2,6-NT	30.76	22.4	0.32
2,6-DNT	33.18	24.2	0.35
2,4-DNT	33.45	24.5	0.30
2,4,6-TNT	35.28	25.9	0.35
TETRYL	36.29	26.6	0.34
4-OHA-2,6-DNT	37.43	27.5	0.37
Azoxydimer	51.92	38.54	0.23

¹ The k-value is the corrected retention time divided by the time required for an unretained compound to travel through the system.

² Width is the peak width at the inflection points expressed in minutes.

Anion Exchange/Reverse Phase Separation on Mixed Mode Column

This procedure improved upon the separation of explosives and metabolites, particularly to allow the trinitrobenzoic acid to be determined. It was used beginning with the organic solvent extracts of the composts. The separation procedure has been developed using a mixed-mode HPLC column in which the phase bonded to the silica surface contains both an octyldecylsilane (reversed-phase function) and a secondary amine (anion exchange function) incorporated into a single ligand in a 1:1 ratio. The separation that can be achieved on this 150 mm X 4.6 mm i.d. column is illustrated in Figure 2A-2. Because of the dual nature of this separating media a mixture of compounds with a wide range of polarities such as those anticipated for TNT and its metabolites can be fractionated in a reasonable run time. The method developed for this separation utilizes a gradient built from three different eluting solvents: A. an aqueous solution containing phosphate that has been adjusted to pH 5.1 with the final concentration of phosphate being 0.015 M in a 10:90 water:methanol solvent, (the sources of phosphate are potassium dihydrogen phosphate and dipotassium hydrogen phosphate in the appropriate amounts to achieve the desired pH and concentration); B. methanol; and C. acetonitrile. The gradient program for this separation starts with equilibration at 72% Solvent A and 28% acetonitrile and elution with this mixture for 1 min after sample injection. From 1 to 5 min a linear gradient reduces solvent A from 72 to 68% and increases acetonitrile from 28 to 32%. Isocratic elution conditions are then maintained for the time interval from 5 to 14 minutes. Between 14 and 20 min a linear gradient reduces solvent A by 4% per minute and increases both methanol and acetonitrile by 2% per minute so that at 20 min the eluting solution is 44% solvent A, 12% methanol and 44% acetonitrile. From 20 to 26 min a linear gradient reduces solvent A from 44 to 2% and increases methanol from 12 to 54% while maintaining the acetonitrile at 44%. Isocratic conditions are then maintained from 26 to 33 min. Then between 33 and 38 min a very steep gradient restores the eluting solution to its starting conditions after which the column is reequilibrated for 7 min. This solvent program is summarized in Table 2A-2. The incorporation of both acetonitrile and methanol in building the gradient appear to be necessary. Peak shape in the early portion of the chromatogram is much better when acetonitrile is the major portion of the organic portion of the eluting solution. However, if acetonitrile is used exclusively in the more rapidly changing portions of the gradient, (after 14 min), large momentary pressure fluctuations are observed. Some of these fluctuations are sufficient to terminate the operation of the more sensitive HPLC instrumentation systems. It is believed that these pressure fluctuations may result from a momentary precipitation of phosphate in a capillary or near a frit. At any rate the substitution of methanol resolves this problem. However, for many HPLC systems where transfer lines are 0.25 mm or greater and frits porosity is 5 microns, it is quite probable that the gradient profile could utilize acetonitrile instead of methanol.

The separation on the mixed-mode column allows trinitrobenzoic acid to be determined along with the other anticipated metabolites. The various chromatographic parameters are listed in Table 2A-3, using the above solvent program, a 50 μ L injection volume, and a wavelength of 254 nm for quantitation. Quantitation is achieved as described above for the reverse phase separation. The detection limits have been determined according to the procedure discussed by Hubaux and Vos, [*Anal. Chem.* 42, 849 (1970)]. Limits are presented for two different confidence levels. The results listed in Table 2A-2 and the chromatogram shown in Figure 2A-2 were generated from sample injection volumes of 50 μ L; and thus they represent a realistic expectation for this method. More ideal results have been generated by injecting standards in volumes of 10 μ L or less. In summary, the method described above yields very good results for most metabolites even though a sample that is 5 or more times the ideal sample volume is injected.

Table 2A-2. Gradient Program for HPLC Method on Anion/C18 Column

Solvent A: 0.015 M Potassium Phosphate at pH 5.1 in 10:90
methanol:water.

Solvent B: Methanol

Solvent C: Acetonitrile

<u>Time</u>	<u>% A</u>	<u>% B</u>	<u>% C</u>
1	72	0	28
5	68	0	32
14	68	0	32
20	44	12	44
26	2	54	44
33	2	54	44
38	72	0	28
45	72	0	28

FIGURE 2A-2. CHROMATOGRAM SHOWING SEPARATION OF EXPLOSIVES AND TNT METABOLITES ON ANION/C18 MIXED-MODE COLUMN. SAMPLE WAS 50 μ L OF A 1 PPM STANDARD MIXTURE.

ORNL-DWG 90-6906

MIXED MODE HPLC OF STANDARDS




Table 2A-3. Chromatographic Parameters for Explosives and TNT Metabolites on Mixed Mode Anion/C18 column

<u>COMPOUND</u>	<u>RET. TIME, MIN.</u>	<u>K¹</u>	<u>SYM² MIN.</u>	<u>DET. 95, PPM</u>	<u>LIMIT³ 99, PPM</u>
1,3,5-TNBA	34.0	17.3	1.3	0.28	0.40
2,4-DA-6-NT	5.7	2.1	1.1	0.36	0.48
2,6-DA-4-NT	4.8	1.6	1.3	0.12	0.16
2,4,6-TNBAIc	7.2	13.3	0.84	0.52	0.72
RDX	11.0	4.9	0.72	0.41	0.56
HMX	14.8	7.0	1.2	0.36	0.54
1,3-DNB	10.1	4.5	1.2	0.20	0.28
1,3,5-TNB	8.3	3.5	1.2	0.20	0.28
2-A-4,6-DNT	19.1	9.3	1.1	0.60	0.88
4-A-2,6-DNT	18.2	8.8	0.93	0.24	0.36
2,6-DNT	14.4	6.8	1.1	0.60	0.80
2,4-DNT	14.5	6.8	1.1	0.48	0.64
2,4,6-TNT	12.8	5.9	1.1	0.20	0.40
4-OHA-2,6-DNT	20.2	9.9	1.2	0.8	1.2 ⁴
Azoxydimer	26.6	13.3	1.3	4.00	2.8

¹The K-value, (capacity ratio), is the corrected retention time divided by the time required for an unretained compound to travel through the system.

²SYM is the total peak area after the apex divided by the total peak area before the apex.

³DET. LIMIT is the detection limit in PPM calculated for the 95 % and 99 % confidence levels according the methods suggested by Hubaux and Vos, [*Anal. Chem.* **42**, 849 (1970)].

⁴The detection limits listed for 4-hydroxy-2,6-dinitrotoluene must be considered approximations because this compound has limited stability under the chromatographic conditions employed.

APPENDIX A-3
DATA REPORTS FOR ORGANICS AND METALS
ANALYSES OF TCLP LEACHATES

WATER SEMI-VOLATILE SURROGATE RECOVERY

Name: OAK RIDGE NATIONAL LAB

Contract: NR

Code: OR

Case No.: C705

SAS No.: 92741

SDG No.: C705

EPA	S1	S2	S3	S4	S5	S6	OTHER	TOT
SAMPLE NO.	(NBZ)	(FBB)	(TPH)	(PHL)	(2FP)	(TBP)		OUT
01	890605-119	44	81	89	25	35	49	0
02	890602-166	44	82	93	27	37	56	0
03	890602-168	47	81	92	23	27	24	0
04	890602-170	46	84	87	21	25	40	0
05	890605-120	59	84	87	26	32	52	0
06	890605-121	59	84	82	21	28	55	0
07	890622-041	55	82	91	28	37	53	0
08	890524-072	57	84	78	28	38	57	0
09	890524-073	55	82	83	32	45	61	0
10	890524-074	55	82	83	32	45	61	0
11	890524-075	55	82	83	32	45	61	0
12	890524-076	55	82	83	32	45	61	0
13	890524-077	55	82	83	32	45	61	0
14	890524-078	55	82	83	32	45	61	0
15								
16								
17								
18								
19								
20								
21								
22								
23								
24								
25								
26								
27								
28								
29								
30								

GSF
2/15/90

QC LIMITS
 S1 (NBZ) = Nitrobenzene-d5 (35-114)
 S2 (FBB) = 2-Fluorobiphenyl (43-116)
 S3 (TPH) = Terphenyl-d14 (33-141)
 S4 (PHL) = Phenol-d5 (10-94)
 S5 (2FP) = 2-Fluorophenol (21-100)
 S6 (TBP) = 2,4,6-Trisubstituted Phenol (10-123)

* Column to be used to flag recovery values
 * Values outside of contract required QC limits
 D Surrogates diluted out

1 of 1

FORM 11 SU-1

1/870R

Best Available Copy

CSF
2/1/89

WATER SAMPLING/ANALYTICAL REPORT - SPILL IN THE SPILL EXPLORATE RECORD

1. Name of the Spill: SPILL IN THE SPILL EXPLORATE RECORD
 2. Date of Spill: 2/1/89
 3. Time of Spill: 10:00 AM
 4. Location of Spill: SPILL IN THE SPILL EXPLORATE RECORD
 5. Name of Spill: SPILL IN THE SPILL EXPLORATE RECORD

COMPOUND	SPILL ADDED 100 L	CONCENTRATION 100 L	CONCENTRATION 100 L	CONCENTRATION 100 L	CONCENTRATION 100 L
Phenol	10.00	1.0	10.00	1.0	10.00
1-Chlorophenol	10.00	1.0	10.00	1.0	10.00
2-Chlorophenol	10.00	1.0	10.00	1.0	10.00
3-Chlorophenol	10.00	1.0	10.00	1.0	10.00
4-Chlorophenol	10.00	1.0	10.00	1.0	10.00
5-Chlorophenol	10.00	1.0	10.00	1.0	10.00
6-Chlorophenol	10.00	1.0	10.00	1.0	10.00
7-Chlorophenol	10.00	1.0	10.00	1.0	10.00
8-Chlorophenol	10.00	1.0	10.00	1.0	10.00
9-Chlorophenol	10.00	1.0	10.00	1.0	10.00
10-Chlorophenol	10.00	1.0	10.00	1.0	10.00

COMPOUND	SPILL ADDED 100 L	CONCENTRATION 100 L	CONCENTRATION 100 L	CONCENTRATION 100 L	CONCENTRATION 100 L
Phenol	10.00	1.0	10.00	1.0	10.00
1-Chlorophenol	10.00	1.0	10.00	1.0	10.00
2-Chlorophenol	10.00	1.0	10.00	1.0	10.00
3-Chlorophenol	10.00	1.0	10.00	1.0	10.00
4-Chlorophenol	10.00	1.0	10.00	1.0	10.00
5-Chlorophenol	10.00	1.0	10.00	1.0	10.00
6-Chlorophenol	10.00	1.0	10.00	1.0	10.00
7-Chlorophenol	10.00	1.0	10.00	1.0	10.00
8-Chlorophenol	10.00	1.0	10.00	1.0	10.00
9-Chlorophenol	10.00	1.0	10.00	1.0	10.00
10-Chlorophenol	10.00	1.0	10.00	1.0	10.00

1. Name of the Spill: SPILL IN THE SPILL EXPLORATE RECORD
 2. Date of Spill: 2/1/89
 3. Time of Spill: 10:00 AM
 4. Location of Spill: SPILL IN THE SPILL EXPLORATE RECORD
 5. Name of Spill: SPILL IN THE SPILL EXPLORATE RECORD

Best Available Copy

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

Request Number: OAL92741

TCLP Leachate
18BL351 of
Mesophilic Compost
(August 1) WAG
11-10-90

Procedure Number: 8270

Matrix: TCLP EXTRACT

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/7/24/89

Lab Sample ID: 890524-072

Sample wt/vol: 900 ML

Lab File ID: DC1476

Date Sampled: 19-May-1989

Date Received 24-May-1989 09:10

% Moisture: not dec.

dec:

Date Analyzed: 5-Jul-1989

Material Description ME COMPOST

Date of Report: 14-JUL-89

CAS NO.	COMPOUND	CONCENTRATION UNITS:		2
		(UG/L or UG/KG)	UG/L	
108-95-2	phenol	11.00	U	
111-44-4	bis(2-chloroethyl)ether	11.00	U	
95-57-8	2-chlorophenol	11.00	U	
541-73-1	1,3-dichlorobenzene	11.00	U	
106-46-7	1,4-dichlorobenzene	11.00	U	
100-51-6	benzyl alcohol	11.00	U	
95-50-1	1,2-dichlorobenzene	11.00	U	
95-48-7	2-methylphenol	11.00	U	
108-60-1	bis(2-chloroisopropyl)ether	11.00	U	
106-44-5	4-methylphenol	11.00	U	
621-64-7	n-nitroso-di-n-propylamine	11.00	U	
67-72-1	hexachloroethane	11.00	U	
98-95-3	nitrobenzene	11.00	U	
78-59-1	isophorone	11.00	U	
88-75-5	2-nitrophenol	11.00	U	
105-67-9	2,4-dimethylphenol	11.00	U	
65-85-0	benzoic acid	56.00	U	
111-91-1	bis(2-chloroethoxy)methane	11.00	U	
120-83-2	2,4-dichlorophenol	11.00	U	
120-82-1	1,2,4-trichlorobenzene	11.00	U	
91-20-3	naphthalene	11.00	U	
106-47-8	4-chloroaniline	11.00	U	
87-68-3	hexachlorobutadiene	11.00	U	
59-50-7	4-chloro-3-methylphenol	11.00	U	
91-57-6	2-methylnaphthalene	11.00	U	
77-47-4	hexachlorocyclopentadiene	11.00	U	
88-06-2	2,4,6-trichlorophenol	11.00	U	
95-95-4	2,4,5-trichlorophenol	56.00	U	
91-58-7	2-chloronaphthalene	11.00	U	
88-74-4	2-nitroaniline	56.00	U	
131-11-3	dimethylphthalate	11.00	U	
208-96-8	acenaphthylene	11.00	U	
606-20-2	2,6-dinitrotoluene	11.00	U	

REVIEWED

By: *df kwanter*
Date: 8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

Request Number: OAL92741

1BBL3S1

Procedure Number: 8270

Matrix: TCLP EXTRACT

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/7/24/89

Lab Sample ID: 890524-072

Sample wt/vol: 900 ML

Lab File ID: >C1476

Date Sampled: 19-May-1989

Date Received 24-May-1989 09:10

% Moisture: not dec.

dec:

Date Analyzed: 5-Jul-1989

Material Description ME COMPOST

Date of Report: 14-JUL-89

CAS NO.	COMPOUND	CONCENTRATION UNITS: (UG/L or UG/KG) UG/L	2
99-09-2	3-nitroaniline	56.00	U
83-32-9	acenaphthene	11.00	U
51-28-5	2,4-dinitrophenol	56.00	U
100-02-7	4-nitrophenol	56.00	U
132-64-9	dibenzofuran	11.00	U
121-14-2	2,4-dinitrotoluene	11.00	U
84-66-2	diethylphthalate	11.00	U
7005-72-3	4-chlorophenyl-phenylether	11.00	U
86-73-7	fluorene	11.00	U
100-01-6	4-nitroaniline	56.00	U
534-52-1	4,6-dinitro-2-methylphenol	56.00	U
86-30-6	n-nitrosodiphenylamine (1)	11.00	U
101-55-3	4-bromophenyl-phenylether	11.00	U
118-74-1	hexachlorobenzene	11.00	U
87-86-5	pentachlorophenol	56.00	U
85-01-8	phenanthrene	11.00	U
120-12-7	anthracene	11.00	U
84-74-2	di-n-butylphthalate	11.00	U
206-44-0	fluoranthene	11.00	U
129-00-0	pyrene	11.00	U
85-68-7	butylbenzylphthalate	11.00	U
91-94-1	3,3'-dichlorobenzidine	22.00	U
56-55-3	benzo(a)anthracene	11.00	U
218-01-9	chrysene	11.00	U
117-81-7	bis(2-ethylhexyl)phthalate	11.00	U
117-84-0	di-n-octylphthalate	11.00	U
205-99-2	benzo(b)fluoranthene	11.00	U
207-08-9	benzo(k)fluoranthene	11.00	U
50-32-8	benzo(a)pyrene	11.00	U
193-39-5	indeno(1,2,3-cd)pyrene	11.00	U
53-70-3	dibenz(a,h)anthracene	11.00	U
191-24-2	benzo(g,h,i)perylene	11.00	U

(1) - Cannot be separated from Diphenylamine

By

Date

8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

Request Number: OAL92741

1BBL3S1

Procedure Number: 8270

Matrix: TCLP EXTRACT

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/7/24/89

Lab Sample ID: 890524-072

Sample Wt/vol: 900 ML

Lab File ID: >C1476

Date Sampled: 19-May-1989

Date Received 24-May-1989 09:10

% Moisture: not dec. dec:

Date Analyzed: 5-Jul-1989

Material Description HE COMPOST

Date of Report: 14-JUL-89

Number TICs found: 5

CONCENTRATION UNITS:
(UG/L or UG/KG) UG/L

CAS NO.	COMPOUND NAME	RT	CONC	Q
1.	TRINITROTOLUENE	24.27	850	J
2.	UNKNOWN	26.46	13	J
3.	UNKNOWN	28.21	24	J
4.	UNKNOWN	28.46	10	J
5.	UNKNOWN	43.82	10	J

REVIEWED

J. Wacht
8/8/89

Semi-volatile Organic Analysis Data

EPA SAMPLE NO.

Request Number: OAL92741

Procedure Number: 8270

Matrix: TCLP EXTRACT

TCLP Leachate
2BBL352 of
Municipal Compost
(Aliquot 2) WNG
11-38-90

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/7/24/89

Lab Sample ID: 890524-073

Sample wt/vol: 900 ML

Lab File ID: >C1477

Date Sampled: 19-May-1989

Date Received 24-May-1989 09:10

% Moisture: not dec.

dec:

Date Analyzed: 5-JUL-1989

Material Description ME COMPOST

Date of Report: 14-JUL-89

CAS NO.	COMPOUND	CONCENTRATION UNITS:		Q
		(UG/L or UG/KG)	UG/L	
108-95-2	phenol	11.00	U	
111-44-4	bis(2-chloroethyl)ether	11.00	U	
95-57-8	2-chlorophenol	11.00	U	
541-73-1	1,3-dichlorobenzene	11.00	U	
106-46-7	1,4-dichlorobenzene	11.00	U	
100-51-6	benzyl alcohol	11.00	U	
95-50-1	1,2-dichlorobenzene	11.00	U	
95-48-7	2-methylphenol	11.00	U	
108-60-1	bis(2-chloroisopropyl)ether	11.00	U	
106-44-5	4-methylphenol	11.00	U	
621-64-7	n-nitroso-di-n-propylamine	11.00	U	
67-72-1	hexachloroethane	11.00	U	
98-95-3	nitrobenzene	11.00	U	
78-59-1	isophorone	11.00	U	
88-75-5	2-nitrophenol	11.00	U	
105-67-9	2,4-dimethylphenol	11.00	U	
65-85-0	benzoic acid	56.00	U	
111-91-1	bis(2-chloroethoxy)methane	11.00	U	
120-83-2	2,4-dichlorophenol	11.00	U	
120-82-1	1,2,4-trichlorobenzene	11.00	U	
91-20-3	naphthalene	11.00	U	
106-47-8	4-chloroaniline	11.00	U	
87-68-3	hexachlorobutadiene	11.00	U	
59-50-7	4-chloro-3-methylphenol	11.00	U	
91-57-6	2-methylnaphthalene	11.00	U	
77-47-4	hexachlorocyclopentadiene	11.00	U	
88-06-2	2,4,6-trichlorophenol	11.00	U	
95-95-4	2,4,5-trichlorophenol	56.00	U	
91-58-7	2-chloronaphthalene	11.00	U	
88-74-4	2-nitroaniline	56.00	U	
131-11-3	dimethylphthalate	11.00	U	
208-96-8	acenaphthylene	11.00	U	
606-20-2	2,6-dinitrotoluene	11.00	U	

EVENED

Date

8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

Request Number: OAL92741

2BBL3S2

Procedure Number: 8270

Matrix: TCLP EXTRACT

Series: Frequency:

Charge Number: 33900213

Customer Name: GRIEST/7/24/89

Lab Sample ID: 890524-073

Sample wt/vol: 900 ML

Lab File ID: >C1477

Date Sampled: 19-May-1989

Date Received 24-May-1989 09:10

% Moisture: not dec. dec:

Date Analyzed: 5-Jul-1989

Material Description ME COMPOST

Date of Report: 14-JUL-89

CAS NO.	COMPOUND	CONCENTRATION UNITS:		Q
		(UG/L or UG/KG)	UG/L	
99-09-2-----	3-nitroaniline	56.00	U	
83-32-9-----	acenaphthene	11.00	U	
51-28-5-----	2,4-dinitrophenol	56.00	U	
100-02-7-----	4-nitrophenol	56.00	U	
132-64-9-----	dibenzofuran	11.00	U	
121-14-2-----	2,4-dinitrotoluene	11.00	U	
84-66-2-----	diethylphthalate	11.00	U	
7005-72-3-----	4-chlorophenyl-phenylether	11.00	U	
86-73-7-----	fluorene	11.00	U	
100-01-6-----	4-nitroaniline	56.00	U	
534-52-1-----	4,6-dinitro-2-methylphenol	56.00	U	
86-30-6-----	n-nitrosodiphenylamine (1)	11.00	U	
101-55-3-----	4-bromophenyl-phenylether	11.00	U	
118-74-1-----	hexachlorobenzene	11.00	U	
87-86-5-----	pentachlorophenol	56.00	U	
85-01-8-----	phenanthrene	11.00	U	
120-12-7-----	anthracene	11.00	U	
84-74-2-----	di-n-butylphthalate	11.00	U	
206-44-0-----	fluoranthene	11.00	U	
129-00-0-----	pyrene	11.00	U	
85-68-7-----	butylbenzylphthalate	11.00	U	
91-94-1-----	3,3'-dichlorobenzidine	22.00	U	
56-55-3-----	benzo(a)anthracene	11.00	U	
218-01-9-----	chrysene	11.00	U	
117-81-7-----	bis(2-ethylhexyl)phthalate	11.00	U	
117-84-0-----	di-n-octylphthalate	11.00	U	
205-99-2-----	benzo(b)fluoranthene	11.00	U	
207-08-9-----	benzo(k)fluoranthene	11.00	U	
50-32-8-----	benzo(a)pyrene	11.00	U	
193-39-5-----	indeno(1,2,3-cd)pyrene	11.00	U	
53-70-3-----	dibenz(a,h)anthracene	11.00	U	
191-24-2-----	benzo(g,h,i)perylene	11.00	U	

(1) - Cannot be separated from Diphenylamine

8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO

Request Number: 0AL92741

2BBL352

Procedure Number: 8270

Matrix: TCLP EXTRACT

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/7/24/89

Lab Sample ID: 890524-073

Sample wt/vol: 900 ML

Lab File ID: >C1477

Date Sampled: 19-May-1989

Date Received 24-May-1989 09:10

% Moisture: not dec.

dec:

Date Analyzed: 5-Jul-1989

Material Description HE COMPOST

Date of Report: 14-JUL-89

Number TICs found: 4

CONCENTRATION UNITS:
(UG/L or UG/KG) UG/L

CAS NO.	COMPOUND NAME	RT	CONC	Q
1.	TRINITROTOLUENE	24.31	1200	J
2.	UNKNOWN	28.20	22	J
3.	UNKNOWN	28.46	11	J
4.	UNKNOWN	43.83	14	J

REVIEWED

J. Wachtel
Date 8/8/89

Pesticide Organics Analysis Data Sheet

EPA SAMPLE NO.

Request Number: OAL92741

TCLP Leachate
2BBL3S2 of
Mesophilic Compost
(Aliquot 2)WYG
1-28-90

Procedure Number: 8270

Matrix: TCLP EXTRACT

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/7/24/89

Lab Sample ID: 890524-073

Sample wt/vol: 100 ML

Lab File ID: NR

Date Sampled: 19-May-1989

Date Received 24-May-1989 09:10

% Moisture: not dec. dec:

Date Analyzed: 16-Jun-1989

Material Description ME COMPOST

Date of Report: 14-JUL-89

CAS NO.	COMPOUND	CONCENTRATION UNITS:	
		(ug/L or ug/Kg) UG/L	Q
319-84-6	ALPHA-BHC	0.50	U
319-85-7	BETA-BHC	0.50	U
319-86-8	DELTA-BHC	0.50	U
58-89-9	GAMMA-BHC (LINDANE)	0.50	U
76-44-8	HEPTACHLOR	0.50	U
309-00-2	ALDRIN	0.50	U
1024-57-3	HEPTACHLOR EPOXIDE	0.50	U
959-98-8	ENDOSULFAM I	0.50	U
60-57-1	DIELDRIN	1.00	U
72-55-9	4,4'-DDE	1.00	U
72-20-8	ENDRIN	1.00	U
33213-65-9	ENDOSULFAM II	1.00	U
72-54-8	4,4'-DDD	1.00	U
1031-07-8	ENDOSULFAM SULFATE	1.00	U
50-29-3	4,4'-DDT	1.00	U
72-43-5	METHOXYCHLOR	5.00	U
53494-70-5	ENDRIN KETONE	1.00	U
5103-71-9	ALPHA CHLORDANE	5.00	U
5103-74-2	GAMMA CHLORDANE	5.00	U
8001-35-2	TOXAPHENE	10.00	U
12674-11-2	AROCLOR-1016	5.00	U
11104-28-2	AROCLOR-1221	5.00	U
11141-16-5	AROCLOR-1232	5.00	U
53469-21-9	AROCLOR-1242	5.00	U
12672-29-6	AROCLOR-1248	5.00	U
11097-69-1	AROCLOR-1254	10.00	U
11096-84-5	AROCLOR-1260	10.00	U

RECEIVED

JH Washite
8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

Request Number: OAL92741

TCLP Leachate
2BBL3S3 of
Mesophilic Compost
(Matrix Spike)WAG
1-30-90

Procedure Number: 8270

Matrix: TCLP EXTRACT

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/7/24/89

Lab Sample ID: 890524-074

Sample wt/vol: 900 ML

Lab File ID: >C1478

Date Sampled: 19-May-1989

Date Received 24-May-1989 09:10

% Moisture: not dec.

dec:

Date Analyzed: 6-Jul-1989

Material Description HE COMPOST

Date of Report: 14-JUL-89

CAS NO.	COMPOUND	CONCENTRATION UNITS:		Q
		(UG/L or UG/KG)	UG/L	
108-95-2	phenol	72.00	MS	
111-44-4	bis(2-chloroethyl)ether	11.00	U	
95-57-8	2-chlorophenol	87.00	MS	
541-73-1	1,3-dichlorobenzene	11.00	U	
106-46-7	1,4-dichlorobenzene	48.00	MS	
100-51-6	benzyl alcohol	11.00	U	
95-50-1	1,2-dichlorobenzene	11.00	U	
95-48-7	2-methylphenol	11.00	U	
108-60-1	bis(2-chloroisopropyl)ether	11.00	U	
106-44-5	4-methylphenol	11.00	U	
621-64-7	n-nitroso-di-n-propylamine	38.00	MS	
67-72-1	hexachloroethane	11.00	U	
98-95-3	nitrobenzene	11.00	U	
78-59-1	isophorone	11.00	U	
88-75-5	2-nitrophenol	7.00	J	
105-67-9	2,4-dimethylphenol	11.00	U	
65-85-0	benzoic acid	56.00	U	
111-91-1	bis(2-chloroethoxy)methane	11.00	U	
120-83-2	2,4-dichlorophenol	11.00	U	
120-82-1	1,2,4-trichlorobenzene	53.00	MS	
91-20-3	naphthalene	11.00	U	
106-47-8	4-chloroaniline	11.00	U	
87-68-3	hexachlorobutadiene	11.00	U	
59-50-7	4-chloro-3-methylphenol	60.00	MS	
91-57-6	2-methylnaphthalene	11.00	U	
77-47-4	hexachlorocyclopentadiene	11.00	U	
88-06-2	2,4,6-trichlorophenol	11.00	U	
95-95-4	2,4,5-trichlorophenol	56.00	U	
91-58-7	2-chloronaphthalene	11.00	U	
88-74-4	2-nitroaniline	56.00	U	
131-11-3	dimethylphthalate	11.00	U	
208-96-8	acenaphthylene	11.00	U	
606-20-2	2,6-dinitrotoluene	11.00	U	

REVIEWED

B. J. W. Richter
8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

Request Number: OAL92741

2BBL3S3

Procedure Number: 8270

Matrix: TCLP EXTRACT

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/7/24/89

Lab Sample ID: 890524-074

Sample Wt/vol: 900 ML

Lab File ID: DC1478

Date Sampled: 19-May-1989

Date Received: 24-May-1989 09:10

% Moisture: not dec.

dec:

Date Analyzed: 6-Jul-1989

Material Description: HE COMPOST

Date of Report: 14-JUL-89

CAS NO.	COMPOUND	CONCENTRATION UNITS:		Q
		(UG/L or UG/KG)	UG/L	
99-09-2	3-nitroaniline	56.00	U	
83-32-9	acenaphthene	57.00	MS	
51-28-5	2,4-dinitrophenol	56.00	U	
100-02-7	4-nitrophenol	58.00	MS	
132-64-9	dibenzofuran	11.00	U	
121-14-2	2,4-dinitrotoluene	44.00	MS	
84-66-2	diethylphthalate	11.00	U	
7005-72-3	4-chlorophenyl-phenylether	11.00	U	
86-73-7	fluorene	11.00	U	
100-01-6	4-nitroaniline	56.00	U	
534-52-1	4,6-dinitro-2-methylphenol	56.00	U	
86-30-6	n-nitrosodiphenylamine (1)	11.00	U	
101-55-3	4-bromophenyl-phenylether	11.00	U	
118-74-1	hexachlorobenzene	11.00	U	
87-86-5	pentachlorophenol	91.00	MS	
85-01-8	phenanthrene	11.00	U	
120-12-7	anthracene	11.00	U	
84-74-2	di-n-butylphthalate	11.00	U	
206-44-0	fluoranthene	11.00	U	
129-00-0	pyrene	55.00	MS	
85-68-7	butylbenzylphthalate	11.00	U	
91-94-1	3,3'-dichlorobenzidine	22.00	U	
56-55-3	benzo(a)anthracene	11.00	U	
218-01-9	chrysene	11.00	U	
117-81-7	bis(2-ethylhexyl)phthalate	11.00	U	
117-84-0	di-n-octylphthalate	11.00	U	
205-99-2	benzo(b)fluoranthene	11.00	U	
207-08-9	benzo(k)fluoranthene	11.00	U	
50-32-8	benzo(a)pyrene	11.00	U	
193-39-5	indeno(1,2,3-cd)pyrene	11.00	U	
53-70-3	dibenz(a,h)anthracene	11.00	U	
191-24-2	benzo(g,h,i)perylene	11.00	U	

(1) - Cannot be separated from Diphenylamine

B.

8/8/89

Pesticide Organics Analysis Data Sheet

EPA SAMPLE NO.

Request Number: OAL92741

Mesophilic Compost
2BB13S3
TCLP leachate
(Matrix Spike)W H G
1-28-90

Procedure Number: 8270

Matrix: TCLP EXTRACT

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/7/24/89

Lab Sample ID: 890524-074

Sample wt/vol: 100 ML

Lab File ID: NR

Date Sampled: 19-May-1989

Date Received 24-May-1989 09:10

% Moisture: not dec.

dec:

Date Analyzed: 16-Jun-1989

Material Description ME COMPOST

Date of Report: 14-JUL-89

CAS NO.	COMPOUND	CONCENTRATION UNITS: (ug/L or ug/Kg) UG/L	Q
319-84-6	ALPHA-BHC	0.50	U
319-85-7	BETA-BHC	0.50	U
319-86-8	DELTA-BHC	0.50	U
58-89-9	GAMMA-BHC (LINDANE)	3.55	
76-44-8	HEPTACHLOR	1.84	
309-00-2	ALDRIN	1.96	
1024-57-3	HEPTACHLOR EPOXIDE	0.50	U
959-98-8	ENDOSULFAM I	0.50	U
60-57-1	DIELDRIN	6.35	
72-55-9	4,4'-DDE	1.00	U
72-20-8	ENDRIN	6.23	
33213-65-9	ENDOSULFAM II	1.00	U
72-54-8	4,4'-DDD	1.00	U
1031-07-8	ENDOSULFAM SULFATE	1.00	U
50-29-3	4,4'-DDT	10.02	
72-43-5	METHOXYCHLOR	5.00	U
53494-70-5	ENDRIN KETONE	1.00	U
5103-71-9	ALPHA CHLORDANE	5.00	U
5103-74-2	GAMMA CHLORDANE	5.00	U
8001-35-2	TOXAPHENE	10.00	U
12674-11-2	AROCLOR-1016	5.00	U
11104-28-2	AROCLOR-1221	5.00	U
11141-16-5	AROCLOR-1232	5.00	U
53469-21-9	AROCLOR-1242	5.00	U
12672-29-6	AROCLOR-1248	5.00	U
11097-69-1	AROCLOR-1254	10.00	U
11096-82-5	AROCLOR-1260	10.00	U

REVIEWED

J. L. W. H. G.
5/28/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

Request Number: 0AL92741

Procedure Number: 8270

Matrix: TCLP EXTRACT

TCLP Leachate
28BL354 of
Mesophilic Compost
(matrix spike duplicate)

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/7/24/89

Lab Sample ID: 890524-075

Sample wt/vol: 900 ML

Lab File ID: >C1479

Date Sampled: 19-May-1989

Date Received 24-May-1989 09:10

% Moisture: not dec.

dec:

Date Analyzed: 6-Jul-1989

Material Description ME COMPOST

Date of Report: 14-JUL-89

CAS NO.	COMPOUND	CONCENTRATION UNITS: (UG/L or UG/KG) UG/L	Q
108-95-2	phenol	87.00	MS
111-44-4	bis(2-chloroethyl)ether	11.00	U
95-57-8	2-chlorophenol	86.00	MS
541-73-1	1,3-dichlorobenzene	11.00	U
106-46-7	1,4-dichlorobenzene	42.00	MS
100-51-6	benzyl alcohol	11.00	U
95-50-1	1,2-dichlorobenzene	11.00	U
95-48-7	2-methylphenol	11.00	U
108-60-1	bis(2-chloroisopropyl)ether	11.00	U
106-44-5	4-methylphenol	11.00	U
621-64-7	n-nitroso-di-n-propylamine	48.00	MS
67-72-1	hexachloroethane	11.00	U
98-95-3	nitrobenzene	11.00	U
78-59-1	isophorone	11.00	U
88-75-5	2-nitrophenol	11.00	U
105-67-9	2,4-dimethylphenol	11.00	U
65-85-0	benzoic acid	56.00	U
111-91-1	bis(2-chloroethoxy)methane	11.00	U
120-83-2	2,4-dichlorophenol	11.00	U
120-82-1	1,2,4-trichlorobenzene	47.00	MS
91-20-3	naphthalene	11.00	U
106-47-8	4-chloroaniline	11.00	U
87-68-3	hexachlorobutadiene	11.00	U
59-50-7	4-chloro-3-methylphenol	71.00	MS
91-57-6	2-methylnaphthalene	11.00	U
77-47-4	hexachlorocyclopentadiene	11.00	U
88-06-2	2,4,6-trichlorophenol	11.00	U
95-95-4	2,4,5-trichlorophenol	56.00	U
91-58-7	2-chloronaphthalene	11.00	U
88-74-4	2-nitroaniline	56.00	U
131-11-3	dimethylphthalate	11.00	U
208-96-8	acenaphthylene	11.00	U
606-20-2	2,6-dinitrotoluene	11.00	U

REVIEWED

J. Wachter
Date 8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

Request Number: OAL92741

2BBL3S4

Procedure Number: 8270

Matrix: TCLP EXTRACT

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/7/24/89

Lab Sample ID: 890524-075

Sample wt/vol: 900 ML

Lab File ID: >C1479

Date Sampled: 19-May-1989

Date Received 24-May-1989 09:10

% Moisture: not dec.

dec:

Date Analyzed: 6-Jul-1989

Material Description HE COMPOST

Date of Report: 14-JUL-89

CAS NO.	COMPOUND	CONCENTRATION UNITS:		Q
		(UG/L or UG/KG)	UG/L	
99-09-2	3-nitroaniline	56.00	U	
83-32-9	acenaphthene	79.00	MS	
51-28-5	2,4-dinitrophenol	56.00	U	
100-02-7	4-nitrophenol	56.00	U	
132-64-9	dibenzofuran	11.00	U	
121-14-2	2,4-dinitrotoluene	53.00	MS	
84-66-2	diethylphthalate	11.00	U	
7005-72-3	4-chlorophenyl-phenylether	11.00	U	
86-73-7	fluorene	11.00	U	
100-01-6	4-nitroaniline	56.00	U	
534-52-1	4,6-dinitro-2-methylphenol	56.00	U	
86-30-6	n-nitrosodiphenylamine (1)	11.00	U	
101-55-3	4-bromophenyl-phenylether	11.00	U	
118-74-1	hexachlorobenzene	11.00	U	
87-86-5	pentachlorophenol	82.00	MS	
85-01-8	phenanthrene	11.00	U	
120-12-7	anthracene	11.00	U	
84-74-2	di-n-butylphthalate	11.00	U	
206-44-0	fluoranthene	11.00	U	
129-00-0	pyrene	60.00	MS	
85-68-7	butylbenzylphthalate	11.00	U	
91-94-1	3,3'-dichlorobenzidine	22.00	U	
56-55-3	benzo(a)anthracene	11.00	U	
218-01-9	chrysene	11.00	U	
117-81-7	bis(2-ethylhexyl)phthalate	11.00	U	
117-84-0	di-n-octylphthalate	11.00	U	
205-99-2	benzo(b)fluoranthene	11.00	U	
207-08-9	benzo(k)fluoranthene	11.00	U	
50-32-8	benzo(a)pyrene	11.00	U	
193-39-5	indeno(1,2,3-cd)pyrene	11.00	U	
53-70-3	dibenz(a,h)anthracene	11.00	U	
191-24-2	benzo(g,h,i)perylene	11.00	U	

(1) - Cannot be separated from Diphenylamine

Date

8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

Request Number: OAL92741

Procedure Number: 8270

Matrix: TCLP EXTRACT

TCLP Leachate
18BL451 of
Thermophilic Compost
(August 0)WNG
1-27-90

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/7/24/89

Lab Sample ID: 890524-076

Sample wt/vol: 900 ML

Lab File ID: >C1480

Date Sampled: 19-May-1989

Date Received 24-May-1989 09:10

% Moisture: not dec.

dec:

Date Analyzed: 6-Jul-1989

Material Description HE COMPOST

Date of Report: 14-JUL-89

CAS NO.	COMPOUND	CONCENTRATION UNITS:		2
		(UG/L or UG/KG)	UG/L	
108-95-2	phenol	11.00	U	
111-44-4	bis(2-chloroethyl)ether	11.00	U	
95-57-8	2-chlorophenol	11.00	U	
541-73-1	1,3-dichlorobenzene	11.00	U	
106-46-7	1,4-dichlorobenzene	11.00	U	
100-51-6	benzyl alcohol	11.00	U	
95-50-1	1,2-dichlorobenzene	11.00	U	
95-48-7	2-methylphenol	11.00	U	
108-60-1	bis(2-chloroisopropyl)ether	11.00	U	
106-44-5	4-methylphenol	11.00	U	
621-64-7	n-nitroso-di-n-propylamine	11.00	U	
67-72-1	hexachloroethane	11.00	U	
98-95-3	nitrobenzene	11.00	U	
78-59-1	isophorone	11.00	U	
88-75-5	2-nitrophenol	11.00	U	
105-67-9	2,4-dimethylphenol	11.00	U	
65-85-0	benzoic acid	56.00	U	
111-91-1	bis(2-chloroethoxy)methane	11.00	U	
120-83-2	2,4-dichlorophenol	11.00	U	
120-82-1	1,2,4-trichlorobenzene	11.00	U	
91-20-3	naphthalene	11.00	U	
106-47-8	4-chloroaniline	11.00	U	
87-68-3	hexachlorobutadiene	11.00	U	
59-50-7	4-chloro-3-methylphenol	11.00	U	
91-57-6	2-methylnaphthalene	11.00	U	
77-47-4	hexachlorocyclopentadiene	11.00	U	
88-06-2	2,4,6-trichlorophenol	11.00	U	
95-95-4	2,4,5-trichlorophenol	56.00	U	
91-58-7	2-chloronaphthalene	11.00	U	
88-74-4	2-nitroaniline	56.00	U	
131-11-3	dimethylphthalate	11.00	U	
208-96-8	acenaphthylene	11.00	U	
606-20-2	2,6-dinitrotoluene	11.00	U	

REVIEWED

By

Date

J. L. Wachtel
8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

Request Number: OAL92741

188L451

Procedure Number: 8270

Matrix: TCLP EXTRACT

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/7/24/89

Lab Sample ID: 890524-076

Sample wt/vol: 900 ML

Lab File ID: >C1480

Date Sampled: 19-May-1989

Date Received 24-May-1989 09:10

% Moisture: not dec.

dec:

Date Analyzed: 6-Jul-1989

Material Description HE COMPOST

Date of Report: 14-JUL-89

CAS NO.	COMPOUND	CONCENTRATION UNITS:		Q
		(UG/L or UG/KG)	UG/L	
99-09-2	3-nitroaniline	56.00	U	
83-32-9	acenaphthene	11.00	U	
51-28-5	2,4-dinitrophenol	56.00	U	
100-02-7	4-nitrophenol	56.00	U	
132-64-9	dibenzofuran	11.00	U	
121-14-2	2,4-dinitrotoluene	11.00	U	
84-66-2	diethylphthalate	11.00	U	
7005-72-3	4-chlorophenyl-phenylether	11.00	U	
86-73-7	fluorene	11.00	U	
100-01-6	4-nitroaniline	56.00	U	
534-52-1	4,6-dinitro-2-methylphenol	56.00	U	
86-30-6	n-nitrosodiphenylamine (1)	11.00	U	
101-55-3	4-bromophenyl-phenylether	11.00	U	
118-74-1	hexachlorobenzene	11.00	U	
87-86-5	pentachlorophenol	56.00	U	
85-01-8	phenanthrene	11.00	U	
120-12-7	anthracene	11.00	U	
84-74-2	di-n-butylphthalate	4.00	J	
206-44-0	fluoranthene	11.00	U	
129-00-0	pyrene	11.00	U	
85-68-7	butylbenzylphthalate	11.00	U	
91-94-1	3,3'-dichlorobenzidine	22.00	U	
56-55-3	benzo(a)anthracene	11.00	U	
218-01-9	chrysene	11.00	U	
117-81-7	bis(2-ethylhexyl)phthalate	11.00	U	
117-84-0	di-n-octylphthalate	11.00	U	
205-99-2	benzo(b)fluoranthene	11.00	U	
207-08-9	benzo(k)fluoranthene	11.00	U	
50-32-8	benzo(a)pyrene	11.00	U	
193-39-5	indeno(1,2,3-cd)pyrene	11.00	U	
53-70-3	dibenz(a,h)anthracene	11.00	U	
191-24-2	benzo(g,h,i)perylene	11.00	U	

(1) - Cannot be separated from Diphenylamine

REVIEWED

S. *Wacht*
Date 8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

Request Number: OAL92741

188L451

Procedure Number: 8270

Matrix: TCLP EXTRACT

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/7/24/89

Lab Sample ID: 890524-076

Sample wt/vol: 900 ML

Lab File ID: >C1480

Date Sampled: 19-May-1989

Date Received 24-May-1989 09:10

% Moisture: not dec.

dec:

Date Analyzed: 6-Jul-1989

Material Description HE COMPOST

Date of Report: 14-JUL-89

Number TICs found: 5

CONCENTRATION UNITS:
(UG/L or UG/KG) UG/L

CAS NO.	COMPOUND NAME	RT	CONC	Q
1.	UNKNOWN	7.36	14	J
2.	TRINITROTOLUENE	24.26	260	J
3.	UNKNOWN	28.46	16	J
4.	UNKNOWN HYDROCARBON	40.67	28	J
5.	UNKNOWN	43.84	51	J

RECEIVED

J. Wachter
8/8/89

Pesticide Organics Analysis Data Sheet

EPA SAMPLE NO.

Request Number: OAL92741

Procedure Number: 8270

Matrix: TCLP EXTRACT

TCLP Leachate
18BL4S1 at
Thermophilic Compost
(Aliquot 2)
WAG
1-28-90

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/7/24/89

Lab Sample ID: 890524-076

Sample Wt/vol: 100 ML

Lab File ID: NR

Date Sampled: 19-May-1989

Date Received 24-May-1989 09:10

% Moisture: not dec.

dec:

Date Analyzed: 16-Jun-1989

Material Description HE COMPOST

Date of Report: 14-JUL-89

CAS NO.	COMPOUND	CONCENTRATION UNITS:		Q
		(ug/L or ug/Kg) UG/L		
319-84-6	ALPHA-BHC	0.50	U	
319-85-7	BETA-BHC	0.50	U	
319-86-8	DELTA-BHC	0.50	U	
58-89-9	GAMMA-BHC (LINDANE)	0.50	U	
76-44-8	HEPTACHLOR	0.50	U	
309-00-2	ALDRIN	0.50	U	
1024-57-3	HEPTACHLOR EPOXIDE	0.50	U	
959-98-8	ENDOSULFAM I	0.50	U	
60-57-1	DIELDRIN	1.00	U	
72-55-9	4,4'-DDE	1.00	U	
72-20-8	ENDRIN	1.00	U	
33213-65-9	ENDOSULFAM II	1.00	U	
72-54-8	4,4'-DDD	1.00	U	
1031-07-8	ENDOSULFAM SULFATE	1.00	U	
50-29-3	4,4'-DDT	1.00	U	
72-43-5	METHOXYCHLOR	5.00	U	
53494-70-5	ENDRIN KETONE	1.00	U	
5103-71-9	ALPHA CHLORDANE	5.00	U	
5103-74-2	GAMMA CHLORDANE	5.00	U	
8001-35-2	TOXAPHENE	10.00	U	
12674-11-2	AROCLOR-1016	5.00	U	
11104-28-2	AROCLOR-1221	5.00	U	
11141-16-5	AROCLOR-1232	5.00	U	
53469-21-9	AROCLOR-1242	5.00	U	
12672-29-6	AROCLOR-1248	5.00	U	
11097-69-1	AROCLOR-1254	10.00	U	
11096-82-5	AROCLOR-1260	10.00	U	

E. J. WAGNER

J. J. WAGNER
8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO

Request Number: OAL92741

Procedure Number: 8270

Matrix: TCLP EXTRACT

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/7/24/89

Lab Sample ID: 890524-077

Sample wt/vol: 900 ML

Lab File ID: DC1481

Date Sampled: 19-May-1989

Date Received 24-May-1989 09:10

% Moisture: not dec.

dec:

Date Analyzed: 6-Jul-1989

Material Description ME COMPOST

Date of Report: 14-JUL-89

CAS NO.	COMPOUND	CONCENTRATION UNITS		Q
		(UG/L or UG/KG)	UG/L	
108-95-2	phenol	11.00	U	
111-44-4	bis(2-chloroethyl)ether	11.00	U	
95-57-8	2-chlorophenol	11.00	U	
541-73-1	1,3-dichlorobenzene	11.00	U	
106-46-7	1,4-dichlorobenzene	11.00	U	
100-51-6	benzyl alcohol	11.00	U	
95-50-1	1,2-dichlorobenzene	11.00	U	
95-48-7	2-methylphenol	11.00	U	
108-60-1	bis(2-chloroisopropyl)ether	11.00	U	
106-44-5	4-methylphenol	11.00	U	
621-64-7	n-nitroso-di-n-propylamine	11.00	U	
67-72-1	hexachloroethane	11.00	U	
98-95-3	nitrobenzene	11.00	U	
78-59-1	isophorone	11.00	U	
88-75-5	2-nitrophenol	11.00	U	
105-67-9	2,4-dimethylphenol	11.00	U	
65-85-0	benzoic acid	56.00	U	
111-91-1	bis(2-chloroethoxy)methane	11.00	U	
120-83-2	2,4-dichlorophenol	11.00	U	
120-82-1	1,2,4-trichlorobenzene	11.00	U	
91-20-3	naphthalene	11.00	U	
106-47-8	4-chloroaniline	11.00	U	
87-68-3	hexachlorobutadiene	11.00	U	
59-50-7	4-chloro-3-methylphenol	11.00	U	
91-57-6	2-methylnaphthalene	11.00	U	
77-47-4	hexachlorocyclopentadiene	11.00	U	
88-06-2	2,4,6-trichlorophenol	11.00	U	
95-95-4	2,4,5-trichlorophenol	56.00	U	
91-58-7	2-chloronaphthalene	11.00	U	
88-74-4	2-nitroaniline	56.00	U	
131-11-3	dimethylphthalate	11.00	U	
208-96-8	acenaphthylene	11.00	U	
606-20-2	2,6-dinitrotoluene	11.00	U	

Date

L. J. Lachter
8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

Request Number: OAL92741

1BBL4S2

Procedure Number: 8270

Matrix: TCLP EXTRACT

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/7/24/89

Lab Sample ID: 890524-077

Sample wt/vol: 900 ML

Lab File ID: >C1481

Date Sampled: 19-May-1989

Date Received 24-May-1989 09:10

% Moisture: not dec.

dec:

Date Analyzed: 6-Jul-1989

Material Description HE COMPOST

Date of Report: 14-JUL-89

CAS NO.	COMPOUND	CONCENTRATION UNITS: (UG/L or UG/KG) UG/L	2
99-09-2	3-nitroaniline	56.00	U
83-32-9	acenaphthene	11.00	U
51-28-5	2,4-dinitrophenol	56.00	U
100-02-7	4-nitrophenol	56.00	U
132-64-9	dibenzofuran	11.00	U
121-14-2	2,4-dinitrotoluene	11.00	U
84-66-2	diethylphthalate	11.00	U
7005-72-3	4-chlorophenyl-phenylether	11.00	U
86-73-7	fluorene	11.00	U
100-01-6	4-nitroaniline	56.00	U
534-52-1	4,6-dinitro-2-methylphenol	56.00	U
86-30-6	n-nitrosodiphenylamine (1)	11.00	U
101-55-3	4-bromophenyl-phenylether	11.00	U
118-74-1	hexachlorobenzene	11.00	U
87-86-5	pentachlorophenol	56.00	U
85-01-8	phenanthrene	11.00	U
120-12-7	anthracene	11.00	U
84-74-2	di-n-butylphthalate	8.00	J
206-44-0	fluoranthene	11.00	U
129-00-0	pyrene	11.00	U
85-68-7	butylbenzylphthalate	11.00	U
91-94-1	3,3'-dichlorobenzidine	22.00	U
56-55-3	benzo(a)anthracene	11.00	U
218-01-9	chrysene	11.00	U
117-81-7	bis(2-ethylhexyl)phthalate	11.00	U
117-84-0	di-n-octylphthalate	11.00	U
205-99-2	benzo(b)fluoranthene	11.00	U
207-08-9	benzo(k)fluoranthene	11.00	U
50-32-8	benzo(a)pyrene	11.00	U
193-39-5	indeno(1,2,3-cd)pyrene	11.00	U
53-70-3	dibenz(a,h)anthracene	11.00	U
191-24-2	benzo(g,h,i)perylene	11.00	U

(1) - Cannot be separated from Diphenylamine

E. S. S. S.

H. W. W. W.
8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

Request Number: OAL92741

1BBL4S2

Procedure Number: 8270

Matrix: TCLP EXTRACT

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/7/24/89

Lab Sample ID: 890524-077

Sample Wt/vol: 900 ML

Lab File ID: PC1481

Date Sampled: 19-May-1989

Date Received 24-May-1989 09:10

% Moisture: not dec. dec:

Date Analyzed: 6-Jul-1989

Material Description HE COMPOST

Date of Report: 14-JUL-89

Number TICs found: 2

CONCENTRATION UNITS:
(UG/L or UG/KG) UG/L

CAS NO.	COMPOUND NAME	RT	CONC	Q
1.	UNKNOWN	7.37	14	J
2.	TRINITROTOLUENE	24.26	240	J

REVIEWED

Jf Wachte
8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

Request Number: OAL92741

TCLP Blank
BLK WME
1-24-90

Procedure Number: 8270

Matrix: TCLP EXTRACT

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/7/24/89

Lab Sample ID: 890524-078

Sample wt/vol: 900 ML

Lab File ID: DC1482

Date Sampled: 19-May-1989

Date Received 24-May-1989 09:10

% Moisture: not dec.

dec:

Date Analyzed: 6-Jul-1989

Material Description HE COMPOST

Date of Report: 14-JUL-89

CAS NO.	COMPOUND	CONCENTRATION UNITS		Q
		(UG/L or UG/KG)	UG/L	
108-95-2	phenol	11.00	U	
111-44-4	bis(2-chloroethyl)ether	11.00	U	
95-57-8	2-chlorophenol	11.00	U	
541-73-1	1,3-dichlorobenzene	11.00	U	
106-46-7	1,4-dichlorobenzene	11.00	U	
100-51-6	benzyl alcohol	11.00	U	
95-50-1	1,2-dichlorobenzene	11.00	U	
95-48-7	2-methylphenol	11.00	U	
108-60-1	bis(2-chloroisopropyl)ether	11.00	U	
106-44-5	4-methylphenol	11.00	U	
621-64-7	n-nitroso-di-n-propylamine	11.00	U	
67-72-1	hexachloroethane	11.00	U	
98-95-3	nitrobenzene	11.00	U	
78-59-1	isophorone	11.00	U	
88-75-5	2-nitrophenol	11.00	U	
105-67-9	2,4-dimethylphenol	11.00	U	
65-85-0	benzoic acid	56.00	U	
111-91-1	bis(2-chloroethoxy)methane	11.00	U	
120-83-2	2,4-dichlorophenol	11.00	U	
120-82-1	1,2,4-trichlorobenzene	11.00	U	
91-20-3	naphthalene	11.00	U	
106-47-8	4-chloroaniline	11.00	U	
87-68-3	hexachlorobutadiene	11.00	U	
59-50-7	4-chloro-3-methylphenol	11.00	U	
91-57-6	2-methylnaphthalene	11.00	U	
77-47-4	hexachlorocyclopentadiene	11.00	U	
88-06-2	2,4,6-trichlorophenol	11.00	U	
95-95-4	2,4,5-trichlorophenol	56.00	U	
91-58-7	2-chloronaphthalene	11.00	U	
88-74-4	2-nitroaniline	56.00	U	
131-11-3	dimethylphthalate	11.00	U	
208-96-8	acenaphthylene	11.00	U	
606-20-2	2,6-dinitrotoluene	11.00	U	

RECEIVED

J. J. Wachte
8/08/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

Request Number: OAL92741

BLK

Procedure Number: 8270

Matrix: TCLP EXTRACT

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/7/24/89

Lab Sample ID: 890524-078

Sample wt/vol: 900 ML

Lab File ID: >C1482

Date Sampled: 19-May-1989

Date Received 24-May-1989 09:10

% Moisture: not dec.

dec:

Date Analyzed: 6-Jul-1989

Material Description NE COMPOST

Date of Report: 14-JUL-89

CAS NO.	COMPOUND	CONCENTRATION UNITS:		Q
		(UG/L or UG/KG)	UG/L	
99-09-2	3-nitroaniline	56.00	U	
83-32-9	acenaphthene	11.00	U	
51-28-5	2,4-dinitrophenol	56.00	U	
100-04-7	4-nitrophenol	56.00	U	
132-64-9	dibenzofuran	11.00	U	
121-14-2	2,4-dinitrotoluene	11.00	U	
84-66-2	diethylphthalate	11.00	U	
7005-72-3	4-chlorophenyl-phenylether	11.00	U	
86-73-7	fluorene	11.00	U	
100-01-6	4-nitroaniline	56.00	U	
534-52-1	4,6-dinitro-2-methylphenol	56.00	U	
86-30-6	n-nitrosodiphenylamine (1)	11.00	U	
101-55-3	4-bromophenyl-phenylether	11.00	U	
118-74-1	hexachlorobenzene	11.00	U	
87-86-5	pentachlorophenol	56.00	U	
85-01-8	phenanthrene	11.00	U	
120-12-7	anthracene	11.00	U	
84-74-2	di-n-butylphthalate	11.00	U	
206-44-0	fluoranthene	11.00	U	
129-00-0	pyrene	11.00	U	
85-68-7	butylbenzylphthalate	11.00	U	
91-94-1	3,3'-dichlorobenzidine	22.00	U	
56-55-3	benzo(a)anthracene	11.00	U	
218-01-9	chrysene	11.00	U	
117-81-7	bis(2-ethylhexyl)phthalate	11.00	U	
117-84-0	di-n-octylphthalate	11.00	U	
205-99-2	benzo(b)fluoranthene	11.00	U	
207-08-9	benzo(k)fluoranthene	11.00	U	
50-32-8	benzo(a)pyrene	11.00	U	
193-39-5	indeno(1,2,3-cd)pyrene	11.00	U	
53-70-3	dibenz(a,h)anthracene	11.00	U	
191-24-2	benzo(g,h,i)perylene	11.00	U	

(1) - Cannot be separated from Diphenylamine

Date

8/08/89

Semi-volatile Organic Analysis Data

EPA SAMPLE NO.

Request Number: OAL92741

BLK

Procedure Number: 8270

Matrix: TCLP EXTRACT

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/7/24/89

Lab Sample ID: 890524-078

Sample wt/vol: 900 ML

Lab File ID: >C1482

Date Sampled: 19-May-1989

Date Received 24-May-1989 09:10

X Moisture: not dec.

dec:

Date Analyzed: 6-Jul-1989

Material Description HE COMPOST

Date of Report: 14-JUL-89

Number TICs found: 4

CONCENTRATION UNITS:
(UG/L or UG/KG) UG/L

CAS NO.	COMPOUND NAME	RT	CONC	Q
1.	UNKNOWN	7.40	24	J
2.	UNKNOWN	28.46	29	J
3.	UNKNOWN HYDROCARBON	40.68	70	J
4.	UNKNOWN	43.84	140	J

REVIEWED

by J. Wachtel
Date 8/8/89

1D

PESTICIDE ORGANICS ANALYSIS DATA SHEET

LAB NAME: OAK RIDGE NATIONAL LAB
 LAB CODE: ORNL CASE NO: NA
 MATRIX: WATER
 SAMPLE WT/VOL: 100 ML
 LEVEL: LOW
 % MOISTURE:
 EXTRACTION METHOD: SEPF
 GPC CLEANUP: N pH: 5.3

EPA SAMPLE NO.: 2BBL3S2
 SAS NO: NA SDG NO: 2741R
 LAB SAMPLE ID: 890524-073
 LAB FILE ID:
 DATE RECEIVED: 05/24/89
 DATE EXTRACTED: 06/06/89
 DATE ANALYZED: 06/16/89
 DILUTION FACTOR: 1

CAS NO	COMPOUND	CONCENTRATION (UG/L)	UNITS Q
319-84-6	ALPHA-BHC	0.50	U
319-85-7	BETA-BHC	0.50	U
319-86-8	DELTA-BHC	0.50	U
58-89-9	GAMMA-BHC (LINDANE)	0.50	U
76-44-8	HEPTACHLOR	0.50	U
309-86-8	ALDRIN	0.50	U
1024-57-3	HEPTACHLOR EPOXIDE	0.50	U
959-98-8	ENDOSULFAN I	0.50	U
60-57-1	DIELDRIN	1.00	U
72-55-9	4,4'-DDE	1.00	U
77-20-8	ENDRIN	1.00	U
33213 65-9	ENDOSULFAN II	1.00	U
72-54-8	4,4'-DDD	1.00	U
1031-07-8	ENDOSULFAN SULFATE	1.00	U
50-29-3	4,4'-DDT	1.00	U
72-43-5	METHOXYCHLOR	5.0	U
53494-70-5	ENDRIN KETONE	1.00	U
5103-71-9	ALPHA-CHLORDANE	5.0	U
5103-74-2	GAMMA-CHLORDANE	5.0	U
8001-35-2	TOXAPHENE	10.0	U
1267-11-2	AROCLOR-1016	5.0	U
11104-28-2	ARO CLOR-1221	5.0	U
11141-16-5	AROCLOR-1232	5.0	U
53469-21-9	AROCLOR-1242	5.0	U
12672-29-6	AROCLOR 1248	5.0	U
11097-69-1	AROCLOR-1254	10.0	U
11096-82-5	AROCLOR-1260	10.0	U

FORM I PEST

1D

PESTICIDE ORGANICS ANALYSIS DATA SHEET

LAB NAME: OAK RIDGE NATIONAL LAB
 LAB CODE: ORNL CASE NO: NA
 MATRIX: WATER
 SAMPLE WT/VOL: 100 ML
 LEVEL: LOW
 % MOISTURE:
 EXTRACTION METHOD: SEPF
 GPC CLEANUP: N pH: 5.4

EPA SAMPLE NO.: 1BBL4S1
 SAS NO: NA SDG NO: 2741F
 LAB SAMPLE ID: 890524-077
 LAB FILE ID:
 DATE RECEIVED: 05/24/89
 DATE EXTRACTED: 06/06/89
 DATE ANALYZED: 06/16/89
 DILUTION FACTOR: 1

CAS NO	COMPOUND	CONCENTRATION (UG/L)	UNITS
319-84-6	ALPHA-BHC	0.50	U
319-85-7	BETA-BHC	0.50	U
319-86-8	DELTA-BHC	0.50	U
58-89-9	GAMMA-BHC (LINDANE)	0.50	U
76-44-8	HEPTACHLOR	0.50	U
309-86-8	ALDRIN	0.50	U
1024-57-3	HEPTACHLOR EPOXIDE	0.50	U
959-98-8	ENDOSULFAN I	0.50	U
60-57-1	DIELDRIN	1.00	U
72-55-9	4,4'-DDE	1.00	U
77-20-8	ENDRIN	1.00	U
33213-65-9	ENDOSULFAN II	1.00	U
72-54-8	4,4'-DDD	1.00	U
1031-07-8	ENDOSULFAN SULFATE	1.00	U
50-29-3	4,4'-DDT	1.00	U
72-43-5	METHOXYCHLOR	5.0	U
53494-70-5	ENDRIN KETONE	1.00	U
5103-71-9	ALPHA-CHLORDANE	5.0	U
5103-74-2	GAMMA-CHLORDANE	5.0	U
8001-35-2	TOXAPHENE	10.0	U
1267-11-2	AROCLOR-1016	5.0	U
11104-28-2	AROCLOR-1221	5.0	U
11141-16-5	AROCLOR-1232	5.0	U
53469-21-9	AROCLOR-1242	5.0	U
12672-29-6	AROCLOR 1248	5.0	U
11097-69-1	AROCLOR-1254	10.0	U
11096-82-5	AROCLOR-1260	10.0	U

FORM I PEST

2E

WATER PESTICIDE SURROGATE RECOVERY

LAB NAME: OAK RIDGE NATIONAL LABORATORY CONTRACT: NA
 LAB CODE: ORNL CASE NO.: NA SAS NO.:NA SDG NO.: 2741R

EPA		S1		OTHER	
SAMPLE NO.		(DCBP)	*		*
1	BLANK		94		
2	2BBL3S2		95		
3	2BBL3S3MS		79		
4	1BBL4S1		85		
5					
6					
7					
8					
9					
10					
11					
12					
13					
14					
15					
16					
17					
18					
19					
20					
21					
22					
23					
24					
25					
26					
27					
28					
29					
30					

ADVISORY QC LIMITS (20-150)

S1 (DBC)=Dibutylchloroendate
 * Column used to flag recovery values
 * Value outside QC limits
 + No surrogate detected in sample
 D Surrogates diluted out

* Value outside QC limits
 + No surrogate detected in sample

1D

PESTICIDE ORGANICS ANALYSIS DATA SHEET

LAB NAME: OAK RIDGE NATIONAL LAB EPA SAMPLE NO.: 2BRL3SDMS
 LAB CODE: ORNL CASE NO: NA SAS NO: NA SDG NO: 2741R
 MATRIX: WATER LAB SAMPLE ID: 890524-074
 SAMPLE WT/VOL: 100 ML LAB FILE ID:
 LEVEL: LOW DATE RECEIVED: 05/24/89
 % MOISTURE: DATE EXTRACTED: 06/06/89
 EXTRACTION METHOD: SEPF DATE ANALYZED: 06/16/89
 GPC CLEANUP: N pH: 5.5 DILUTION FACTOR: 1

CAS NO	COMPOUND	CONCENTRATION (UG/L)	UNITS	Q
319-84-6	ALPHA-BHC	0.50	U	
319-85-7	BETA-BHC	0.50	U	
319-86-8	DELTA-BHC	0.50	U	
58-89-9	GAMMA-BHC (LINDANE)	3.55		
76-44-8	HEPTACHLOR	1.84		
309-86-8	ALDRIN	1.96		
1024-57-3	HEPTACHLOR EPOXIDE	0.50	U	
959-98-8	ENDOSULFAN I	0.50	U	
60-57-1	DIELDRIN	6.35		
72-55-9	4,4'-DDE	1.00	U	
77-20-8	ENDRIN	6.23		
33213-65-9	ENDOSULFAN II	1.00	U	
72-54-8	4,4'-DDD	1.00	U	
1031-07-8	ENDOSULFAN SULFATE	1.00	U	
50-29-3	4,4'-DDT	10.02		
72-43-5	METHOXYCHLOR	5.0	U	
53494-70-5	ENDRIN KETONE	1.00	U	
5103-71-9	ALPHA-CHLORDANE	5.0	U	
5103-74-2	GAMMA-CHLORDANE	5.0	U	
8001-35-2	TOXAPHENE	10.0	U	
1267-11-2	AROCLOR-1016	5.0	U	
11104-28-2	AROCLOR-1221	5.0	U	
11141-16-5	AROCLOR-1232	5.0	U	
53469-21-9	AROCLOR-1242	5.0	U	
12672-29-6	AROCLOR 1248	5.0	U	
11097-69-1	AROCLOR-1254	10.0	U	
11096-82-5	AROCLOR-1260	10.0	U	

FORM I PEST

92

Analyst No.	Customers Id	Date/Time Sampled	Analysis	Result	Units	Procedure No	Completed
Frequency	Material Desc.						
1890524-038	1891451						
ITCLP	EXTRACT HE COMPOST						
	Thermophilic (No 1 Mixture)						
	Compost						
	W.H.G.						
	1-28-70						
			AG	< 1.0E-02	MG/L	EPA 200.7	6-Jun-1989
			AL	2.2E+00	MG/L	EPA 200.7	6-Jun-1989
			AS	< 1.0E-01	MG/L	EPA 200.7	6-Jun-1989
			B	5.9E-01	MG/L	EPA 200.7	6-Jun-1989
			BA	8.0E-01	MG/L	EPA 200.7	6-Jun-1989
			BE	< 8.0E-04	MG/L	EPA 200.7	6-Jun-1989
			CA	1.5E+02	MG/L	EPA 200.7	6-Jun-1989
			CD	< 4.0E-03	MG/L	EPA 200.7	6-Jun-1989
			CO	6.6E-03	MG/L	EPA 200.7	6-Jun-1989
			CR	2.0E-02	MG/L	EPA 200.7	6-Jun-1989
			CU	< 2.0E-02	MG/L	EPA 200.7	6-Jun-1989
			FE	8.5E-01	MG/L	EPA 200.7	6-Jun-1989
			GA	< 6.0E-01	MG/L	EPA 200.7	6-Jun-1989
			MG	2.4E+01	MG/L	EPA 200.7	6-Jun-1989
			MM	1.6E+00	MG/L	EPA 200.7	6-Jun-1989
			MO	< 8.0E-02	MG/L	EPA 200.7	6-Jun-1989
			NA	1.5E+03	MG/L	EPA 200.7	6-Jun-1989
			NI	< 1.2E-02	MG/L	EPA 200.7	6-Jun-1989
			P	1.9E+01	MG/L	EPA 200.7	6-Jun-1989
			PB	< 6.0E-02	MG/L	EPA 200.7	6-Jun-1989
			SB	< 8.0E-02	MG/L	EPA 200.7	6-Jun-1989
			SE	< 1.6E-01	MG/L	EPA 200.7	6-Jun-1989
			SI	4.9E+00	MG/L	EPA 200.7	6-Jun-1989
			SH	< 1.0E-01	MG/L	EPA 200.7	6-Jun-1989
			SR	6.7E-01	MG/L	EPA 200.7	6-Jun-1989
			TI	< 4.0E-02	MG/L	EPA 200.7	6-Jun-1989
			V	< 8.0E-03	MG/L	EPA 200.7	6-Jun-1989
			ZM	5.9E-01	MG/L	EPA 200.7	6-Jun-1989
			ZR	< 4.0E-02	MG/L	EPA 200.7	6-Jun-1989

Analy No.	Customers Id	Date/Time Sampled	Matrix	Material Desc.	Analysis	Result	Units	Procedure No	Completed
1890524-039	2BBL452								
ITCLP	EXTRACT HE COMPOST								
	Thermophilic (Mar 2 11:20 AM)								
	Compost								
	WUC								
	1-28-70								
			AG			< 1.0E-02	MG/L	EPA 200.7	6-Jun-1989
			AL			8.6E+00	MG/L	EPA 200.7	6-Jun-1989
			AS			< 1.0E-01	MG/L	EPA 200.7	6-Jun-1989
			B			6.3E-01	MG/L	EPA 200.7	6-Jun-1989
			BA			9.4E-01	MG/L	EPA 200.7	6-Jun-1989
			BE			< 8.0E-04	MG/L	EPA 200.7	6-Jun-1989
			CA			1.3E+02	MG/L	EPA 200.7	6-Jun-1989
			CD			4.5E-03	MG/L	EPA 200.7	6-Jun-1989
			CO			9.7E-03	MG/L	EPA 200.7	6-Jun-1989
			CR			2.9E-02	MG/L	EPA 200.7	6-Jun-1989
			CU			< 2.0E-02	MG/L	EPA 200.7	6-Jun-1989
			FE			3.6E+00	MG/L	EPA 200.7	6-Jun-1989
			GA			< 6.0E-01	MG/L	EPA 200.7	6-Jun-1989
			MG			2.2E+01	MG/L	EPA 200.7	6-Jun-1989
			MN			1.6E+00	MG/L	EPA 200.7	6-Jun-1989
			MO			< 8.0E-02	MG/L	EPA 200.7	6-Jun-1989
			NA			1.5E+03	MG/L	EPA 200.7	6-Jun-1989
			NI			2.9E-02	MG/L	EPA 200.7	6-Jun-1989
			P			2.0E+01	MG/L	EPA 200.7	6-Jun-1989
			PB			< 6.0E-02	MG/L	EPA 200.7	6-Jun-1989
			SB			< 1.6E-01	MG/L	EPA 200.7	6-Jun-1989
			SE			< 1.6E-01	MG/L	EPA 200.7	6-Jun-1989
			SI			< 1.0E-01	MG/L	EPA 200.7	6-Jun-1989
			SM			7.0E-01	MG/L	EPA 200.7	6-Jun-1989
			SR			1.5E-01	MG/L	EPA 200.7	6-Jun-1989
			TI			< 8.0E-03	MG/L	EPA 200.7	6-Jun-1989
			V			6.5E-01	MG/L	EPA 200.7	6-Jun-1989
			ZN			< 4.0E-02	MG/L	EPA 200.7	6-Jun-1989
			ZR						

APPENDIX A-4

DATA REPORTS FOR ORGANICS AND METALS
ANALYSES OF CCLT LEACHATES

>>> REQUEST FOR ANALYTICAL SERVICES <<<

ACD Request No
OAL 92805

Type of Collection		Sample Matrix		Sample ID (15)		Sample Date		Analysis (30)		Detection Limit		
Time	Location	W	A	T	E	R	mm	dd	yy	time	Method ID (15)	Detection Limit
3390002113	W. H. GRIEST											
<p>Comments: pH 5.0 water (acidified w/ H₂SO₄/HNO₃); Residue of Compost. Samples are in refrigerator, shelf #1 left side (our archival samples are in the right side) F-10.</p>												
890622 027	W. H. GRIEST											
890622 028	W. H. GRIEST											
890622 030	W. H. GRIEST											
890622 031	W. H. GRIEST											
890622 032	W. H. GRIEST											
890622 033	W. H. GRIEST											
890622 034	W. H. GRIEST											
890622 035	W. H. GRIEST											
890622 036	W. H. GRIEST											
890622 037	W. H. GRIEST											
890622 038	W. H. GRIEST											
890622 039	W. H. GRIEST											
890622 040	W. H. GRIEST											
890622 041	W. H. GRIEST											
890622 042	W. H. GRIEST											
890622 043	W. H. GRIEST											
890622 044	W. H. GRIEST											
890622 045	W. H. GRIEST											
890622 046	W. H. GRIEST											
890622 047	W. H. GRIEST											
890622 048	W. H. GRIEST											
890622 049	W. H. GRIEST											
890622 050	W. H. GRIEST											
890622 051	W. H. GRIEST											
890622 052	W. H. GRIEST											
890622 053	W. H. GRIEST											
890622 054	W. H. GRIEST											
890622 055	W. H. GRIEST											
890622 056	W. H. GRIEST											
890622 057	W. H. GRIEST											
890622 058	W. H. GRIEST											
890622 059	W. H. GRIEST											
890622 060	W. H. GRIEST											
890622 061	W. H. GRIEST											
890622 062	W. H. GRIEST											
890622 063	W. H. GRIEST											
890622 064	W. H. GRIEST											
890622 065	W. H. GRIEST											
890622 066	W. H. GRIEST											
890622 067	W. H. GRIEST											
890622 068	W. H. GRIEST											
890622 069	W. H. GRIEST											
890622 070	W. H. GRIEST											
890622 071	W. H. GRIEST											
890622 072	W. H. GRIEST											
890622 073	W. H. GRIEST											
890622 074	W. H. GRIEST											
890622 075	W. H. GRIEST											
890622 076	W. H. GRIEST											
890622 077	W. H. GRIEST											
890622 078	W. H. GRIEST											
890622 079	W. H. GRIEST											
890622 080	W. H. GRIEST											
890622 081	W. H. GRIEST											
890622 082	W. H. GRIEST											
890622 083	W. H. GRIEST											
890622 084	W. H. GRIEST											
890622 085	W. H. GRIEST											
890622 086	W. H. GRIEST											
890622 087	W. H. GRIEST											
890622 088	W. H. GRIEST											
890622 089	W. H. GRIEST											
890622 090	W. H. GRIEST											
890622 091	W. H. GRIEST											
890622 092	W. H. GRIEST											
890622 093	W. H. GRIEST											
890622 094	W. H. GRIEST											
890622 095	W. H. GRIEST											
890622 096	W. H. GRIEST											
890622 097	W. H. GRIEST											
890622 098	W. H. GRIEST											
890622 099	W. H. GRIEST											
890622 100	W. H. GRIEST											

W. H. GRIEST

RPD

W. H. GRIEST

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

MESO CCLT 1

Lab Name: Oak Ridge National Lab Contract: NR

Lab Code: Case no: ORNL SAS No: 92805 SDG No: C713

Matrix: (soil/water) WATER Lab Sample ID: 890622-027

Sample wt/vol: 900 ML Lab File ID: >C1498

Level: (low/med) LOW Date Received 22-Jun-1989

% Moisture: not dec. dec. Date Analyzed: 13-Jul-1989

Extraction: (Sepf/Cont/Sonc) CONT Date Extracted: 26-Jun-1989

GPC Cleanup: (Y/N) N pH: 7.79 Dilution Factor: 1.05

		CONCENTRATION UNITS:		
		(UG/L or UG/KG)	UG/L	2
CAS NO.	COMPOUND			
108-95-2	phenol	11.00	U	
111-44-4	bis(2-chloroethyl)ether	11.00	U	
95-57-8	2-chlorophenol	11.00	U	
541-73-1	1,3-dichlorobenzene	11.00	U	
106-46-7	1,4-dichlorobenzene	11.00	U	
100-51-6	benzyl alcohol	11.00	U	
95-50-1	1,2-dichlorobenzene	11.00	U	
95-48-7	2-methylphenol	11.00	U	
108-60-1	bis(2-chloroisopropyl)ether	11.00	U	
106-44-5	4-methylphenol	11.00	U	
621-64-7	n-nitroso-di-n-propylamine	11.00	U	
67-72-1	hexachloroethane	11.00	U	
98-95-3	nitrobenzene	11.00	U	
78-59-1	isophorone	11.00	U	
68-75-5	2-nitrophenol	11.00	U	
105-67-9	2,4-dimethylphenol	11.00	U	
65-85-0	benzoic acid	53.00	U	
111-91-1	bis(2-chloroethoxy)methane	11.00	U	
120-83-2	2,4-dichlorophenol	11.00	U	
120-82-1	1,2,4-trichlorobenzene	11.00	U	
91-20-3	naphthalene	11.00	U	
106-47-8	4-chloroaniline	11.00	U	
87-68-3	hexachlorobutadiene	11.00	U	
59-50-7	4-chloro-3-methylphenol	11.00	U	
91-57-6	2-methylnaphthalene	11.00	U	
77-47-4	hexachlorocyclopentadiene	11.00	U	
88-06-2	2,4,6-trichlorophenol	11.00	U	
95-95-4	2,4,5-trichlorophenol	53.00	U	
91-58-7	2-chloronaphthalene	11.00	U	
88-74-4	2-nitroaniline	53.00	U	
131-11-3	dimethylphthalate	11.00	U	
208-96-8	acenaphthylene	11.00	U	
606-20-2	2,6-dinitrotoluene	11.00	U	

REVIEWED

Date

8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

MESO CCLT 1

Lab Name: Oak Ridge National Lab Contract: NR
Lab Code: Case no: ORML SAS No: 92805 SDG No: C713
Matrix: (soil/water) WATER Lab Sample ID: 890622-027
Sample wt/vol: 900 ML Lab File ID: >C1498
Level: (low/med) LOW Date Received 22-Jun-1989
% Moisture: not dec. dec. Date Analyzed: 13-Jul-1989
Extraction: (Sepf/Cont/Sonc) CONT Date Extracted: 26-Jun-1989
GPC Cleanup: (Y/N) N pH: 7.79 Dilution Factor: 1.05

CONCENTRATION UNITS:
(UG/L or UG/KG) UG/L

CAS NO.	COMPOUND	2
99-09-2-----	3-nitroaniline	53.00 U
83-32-9-----	acenaphthene	11.00 U
51-28-5-----	2,4-dinitrophenol	53.00 U
100-02-7-----	4-nitrophenol	53.00 U
132-64-9-----	dibenzofuran	11.00 U
121-14-2-----	2,4-dinitrotoluene	11.00 U
84-66-2-----	diethylphthalate	11.00 U
7005-72-3-----	4-chlorophenyl-phenylether	11.00 U
86-73-7-----	fluorene	11.00 U
100-01-6-----	4-nitroaniline	53.00 U
534-52-1-----	4,6-dinitro-2-methylphenol	53.00 U
86-30-6-----	n-nitrosodiphenylamine (1)	11.00 U
101-55-3-----	4-bromophenyl-phenylether	11.00 U
118-74-1-----	hexachlorobenzene	11.00 U
87-86-5-----	pentachlorophenol	53.00 U
85-01-8-----	phenanthrene	11.00 U
120-12-7-----	anthracene	11.00 U
84-74-2-----	di-n-butylphthalate	11.00 U
206-44-0-----	fluoranthene	11.00 U
129-00-0-----	pyrene	11.00 U
65-68-7-----	butylbenzylphthalate	11.00 U
91-94-1-----	3,3'-dichlorobenzidine	21.00 U
56-55-3-----	benzo(a)anthracene	11.00 U
218-01-9-----	chrysene	11.00 U
117-81-7-----	bis(2-ethylhexyl)phthalate	11.00 U
117-84-0-----	di-n-octylphthalate	11.00 U
205-99-2-----	benzo(b)fluoranthene	11.00 U
207-08-9-----	benzo(k)fluoranthene	11.00 U
50-32-8-----	benzo(a)pyrene	11.00 U
193-39-5-----	indeno(1,2,3-cd)pyrene	11.00 U
53-70-3-----	dibenz(a,h)anthracene	11.00 U
191-24-2-----	benzo(g,h,i)perylene	11.00 U

(1) - Cannot be separated from Diphenylamine

By

Date

df Wachte
8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

Request Number: OAL92805

MESO CCLT 1

Procedure Number: 8270

Matrix: WATER

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/8/22/89

Lab Sample ID: 890622-027

Sample wt/vol: 900 ML

Lab File ID: >C1498

Date Sampled:

Date Received 22-Jun-1989 09:10

% Moisture: not dec.

dec:

Date Analyzed: 13-Jul-1989

Material Description

Date of Report: 20-JUL-89

Number TICs found: 9

CONCENTRATION UNITS:
(UG/L or UG/KG) UG/L

CAS NO.	COMPOUND NAME	RT	CONC	Q
1.	UNKNOWN	7.11	5	J
2.	UNKNOWN	7.84	3	J
3.	UNKNOWN	8.85	7	J
4.	UNKNOWN	21.05	2	J
5.	TRINITROTOLUENE	24.33	300	J
6.	UNKNOWN	26.58	21	J
7.	UNKNOWN	27.50	4	J
8.	UNKNOWN	28.20	4	J
9.	UNKNOWN	28.27	12	J

REVIEWED

d/wachte
8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

Lab Name: Oak Ridge National Lab Contract: NR
Lab Code: Case no: ORNL SAS No: 92805 SDG No: C713
Matrix: (soil/water) WATER Lab Sample ID: 890622-028
Sample wt/vol: 900 ML Lab File ID: >C1499
Level: (low/med) LOW Date Received 22-Jun-1989
% Moisture: not dec. dec. Date Analyzed: 13-Jul-1989
Extraction: (Sepf/Cont/Sonc) CONT Date Extracted: 26-Jun-1989
GPC Cleanup: (Y/N) N pH: 7.79 Dilution Factor: 1.05

MESO CCLT 2

CAS NO.	COMPOUND	CONCENTRATION UNITS: (UG/L or UG/KG) UG/L	Q
108-95-2	phenol	11.00	U
111-44-4	bis(2-chloroethyl)ether	11.00	U
95-57-8	2-chlorophenol	11.00	U
541-73-1	1,3-dichlorobenzene	11.00	U
106-46-7	1,4-dichlorobenzene	11.00	U
100-51-6	benzyl alcohol	11.00	U
95-50-1	1,2-dichlorobenzene	11.00	U
95-48-7	2-methylphenol	11.00	U
108-60-1	bis(2-chloroisopropyl)ether	11.00	U
106-44-5	4-methylphenol	11.00	U
621-64-7	n-nitroso-di-n-propylamine	11.00	U
67-72-1	hexachloroethane	11.00	U
98-95-3	nitrobenzene	11.00	U
78-59-1	isophorone	11.00	U
88-75-5	2-nitrophenol	11.00	U
105-67-9	2,4-dimethylphenol	11.00	U
65-85-0	benzoic acid	53.00	U
111-91-1	bis(2-chloroethoxy)methane	11.00	U
120-83-2	2,4-dichlorophenol	11.00	U
120-82-1	1,2,4-trichlorobenzene	11.00	U
91-20-3	naphthalene	11.00	U
106-47-8	4-chloroaniline	11.00	U
87-68-3	hexachlorobutadiene	11.00	U
59-50-7	4-chloro-3-methylphenol	11.00	U
91-57-6	2-methylnaphthalene	11.00	U
77-47-4	hexachlorocyclopentadiene	11.00	U
88-06-2	2,4,6-trichlorophenol	11.00	U
95-95-4	2,4,5-trichlorophenol	53.00	U
91-58-7	2-chloronaphthalene	11.00	U
88-74-4	2-nitroaniline	53.00	U
131-11-3	dimethylphthalate	11.00	U
208-96-8	acenaphthylene	11.00	U
606-20-2	2,6-dinitrotoluene	11.00	U

REVIEWED

B. *H. Wachte*
8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

MESO CCLT 2

Lab Name: Oak Ridge National Lab Contract: MR

Lab Code: Case no: ORNL SAS No: 92805 SDG No: C713

Matrix: (soil/water) WATER Lab Sample ID: 890622-028

Sample wt/vol: 900 ML Lab File ID: >C1499

Level: (low/med) LOW Date Received 22-Jun-1989

% Moisture: not dec. dec. Date Analyzed: 13-Jul-1989

Extraction: (Sapf/Cont/Sonc) CONT Date Extracted: 26-Jun-1989

GPC Cleanup: (Y/N) N pH: 7.79 Dilution Factor: 1.05

CAS NO.	COMPOUND	CONCENTRATION UNITS:	
		(UG/L or UG/KG) UG/L	Q
99-09-2-----	3-nitroaniline	53.00	U
83-32-9-----	acenaphthene	11.00	U
51-28-5-----	2,4-dinitrophenol	53.00	U
100-02-7-----	4-nitrophenol	53.00	U
132-64-9-----	dibenzofuran	11.00	U
121-14-2-----	2,4-dinitrotoluene	11.00	U
84-66-2-----	diethylphthalate	11.00	U
7005-72-3-----	4-chlorophenyl-phenylether	11.00	U
86-73-7-----	fluorene	11.00	U
100-01-6-----	4-nitroaniline	53.00	U
534-52-1-----	4,6-dinitro-2-methylphenol	53.00	U
86-30-6-----	n-nitrosodiphenylamine (1)	11.00	U
101-55-3-----	4-bromophenyl-phenylether	11.00	U
118-74-1-----	hexachlorobenzene	11.00	U
87-86-5-----	pentachlorophenol	53.00	U
85-01-8-----	phenanthrene	11.00	U
120-12-7-----	anthracene	11.00	U
84-74-2-----	di-n-butylphthalate	11.00	U
206-44-0-----	fluoranthene	11.00	U
129-00-0-----	pyrene	11.00	U
85-68-7-----	butylbenzylphthalate	11.00	U
91-94-1-----	3,3'-dichlorobenzidine	21.00	U
56-55-3-----	benzo(a)anthracene	11.00	U
218-01-9-----	chrysene	11.00	U
117-81-7-----	bis(2-ethylhexyl)phthalate	6.00	J
117-84-0-----	di-n-octylphthalate	11.00	U
205-99-2-----	benzo(b)fluoranthene	11.00	U
207-08-9-----	benzo(k)fluoranthene	11.00	U
50-32-8-----	benzo(a)pyrene	11.00	U
193-39-5-----	indeno(1,2,3-cd)pyrene	11.00	U
53-70-3-----	dibenz(a,h)anthracene	11.00	U
191-24-2-----	benzo(g,h,i)perylene	11.00	U

(1) - Cannot be separated from Diphenylamine

W. W. Wichter
8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

Request Number: OAL92805

MESO CCLT 2

Procedure Number: 8270

Matrix: WATER

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/8/22/89

Lab Sample ID: 890622-028

Sample wt/vol: 900 ML

Lab File ID: >C1499

Date Sampled:

Date Received 22-Jun-1989 09:10

% Moisture: not dec.

dec:

Date Analyzed: 13-Jul-1989

Material Description

Date of Report: 20-JUL-89

Number TICs found: 8

CONCENTRATION UNITS:
(UG/L or UG/KG) UG/L

CAS NO.	COMPOUND NAME	RT	CONC	Q
1.	UNKNOWN	7.09	30	J
2.	UNKNOWN	7.83	3	J
3.	UNKNOWN	8.83	11	J
4.	UNKNOWN	14.20	3	J
5.	TRINITROTOLUENE	24.31	350	J
6.	UNKNOWN	26.56	24	J
7.	UNKNOWN Amino dinitro toluene	28.18	4	J
8.	UNKNOWN	28.25	12	J

WtG
1-25-89

REVIEWED

H. Kachta
8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO

THERMO CCLT 1

Lab Name: Oak Ridge National Lab Contract: NR

Lab Code: Case no: ORNL SAS No: 92805 SDG No: C713

Matrix: (soil/water) WATER Lab Sample ID: 890622-029

Sample wt/vol: 900 ML Lab File ID: KC1500

Level: (low/med) LOW Date Received: 22-Jun-1989

% Moisture: not dec. dec. Date Analyzed: 13-Jul-1989

Extraction: (Sepf/Cont/Sonc) CONT Date Extracted: 26-Jun-1989

GPC Cleanup: (Y/N) X pH: 7.83 Dilution Factor: 1.05

CAS NO.	COMPOUND	CONCENTRATION UNITS:		Q
		(UG/L or UG/KG)	UG/L	
108-95-2	phenol	11.00	U	
111-44-4	bis(2-chloroethyl)ether	11.00	U	
95-57-8	2-chlorophenol	11.00	U	
541-73-1	1,3-dichlorobenzene	11.00	U	
106-46-7	1,4-dichlorobenzene	11.00	U	
100-51-6	benzyl alcohol	11.00	U	
95-50-1	1,2-dichlorobenzene	11.00	U	
95-48-7	2-methylphenol	11.00	U	
108-60-1	bis(2-chloroisopropyl)ether	11.00	U	
106-44-5	4-methylphenol	11.00	U	
621-64-7	n-nitroso-di-n-propylamine	11.00	U	
67-72-1	hexachloroethane	11.00	U	
98-95-3	nitrobenzene	11.00	U	
78-59-1	isophorone	11.00	U	
88-75-5	2-nitrophenol	11.00	U	
105-67-9	2,4-dimethylphenol	11.00	U	
65-85-0	benzoic acid	53.00	U	
111-91-1	bis(2-chloroethoxy)methane	11.00	U	
120-83-2	2,4-dichlorophenol	11.00	U	
120-82-1	1,2,4-trichlorobenzene	11.00	U	
91-20-3	naphthalene	11.00	U	
106-47-8	4-chloroaniline	11.00	U	
87-68-3	hexachlorobutadiene	11.00	U	
59-50-7	4-chloro-3-methylphenol	11.00	U	
91-57-6	2-methylnaphthalene	11.00	U	
77-47-4	hexachlorocyclopentadiene	11.00	U	
88-06-2	2,4,6-trichlorophenol	11.00	U	
95-95-4	2,4,5-trichlorophenol	53.00	U	
91-58-7	2-chloronaphthalene	11.00	U	
88-74-4	2-nitroaniline	53.00	U	
131-11-3	dimethylphthalate	11.00	U	
208-96-8	acenaphthylene	11.00	U	
606-20-2	2,6-dinitrotoluene	11.00	U	

REVIEWED

8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

THERMO CCLT 1

Lab Name: Oak Ridge National Lab Contract: NR

Lab Code: Case no: ORNL SAS No: 92805 SDG No: C713

Matrix: (soil/water) WATER Lab Sample ID: 890622-029

Sample wt/vol: 900 ML Lab File ID: >C1500

Level: (low/med) LOW Date Received 22-Jun-1989

% Moisture: not dec. dec. Date Analyzed: 13-Jul-1989

Extraction: (Sepi/Cont/Sonc) CONT Date Extracted: 26-Jun-1989

GPC Cleanup: (Y/N) N pH: 7.83 Dilution Factor: 1.05

CAS NO.	COMPOUND	CONCENTRATION UNITS:	
		(UG/L or UG/KG) UG/L	Q
99-09-2	3-nitroaniline	53.00	U
83-32-9	acenaphthene	11.00	U
51-28-5	2,4-dinitrophenol	53.00	U
100-02-7	4-nitrophenol	53.00	U
132-64-9	dibenzofuran	11.00	U
121-14-2	2,4-dinitrotoluene	11.00	U
84-66-2	diethylphthalate	11.00	U
7005-72-3	4-chlorophenyl-phenylether	11.00	U
86-73-7	fluorene	11.00	U
100-01-6	4-nitroaniline	53.00	U
534-52-1	4,6-dinitro-2-methylphenol	53.00	U
86-30-6	n-nitrosodiphenylamine (1)	11.00	U
101-55-3	4-bromophenyl-phenylether	11.00	U
118-74-1	hexachlorobenzene	11.00	U
87-86-5	pentachlorophenol	53.00	U
85-01-8	phenanthrene	11.00	U
120-12-7	anthracene	11.00	U
84-74-2	di-n-butylphthalate	11.00	U
206-44-0	fluoranthene	11.00	U
129-00-0	pyrene	11.00	U
85-68-7	butylbenzylphthalate	11.00	U
91-94-1	3,3'-dichlorobenzidine	21.00	U
56-55-3	benzo(a)anthracene	11.00	U
218-01-9	chrysene	11.00	U
117-81-7	bis(2-ethylhexyl)phthalate	11.00	U
117-84-0	di-n-octylphthalate	11.00	U
205-99-2	benzo(b)fluoranthene	11.00	U
207-08-9	benzo(k)fluoranthene	11.00	U
50-32-8	benzo(a)pyrene	11.00	U
193-39-5	indeno(1,2,3-cd)pyrene	11.00	U
53-70-3	dibenz(a,h)anthracene	11.00	U
191-24-2	benzo(g,h,i)perylene	REVIEWED 000	U

(1) - Cannot be separated from Diphenylamine

Date

8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

Request Number: OAL92805

THERMO CCLT 1

Procedure Number: 8270

Matrix: WATER

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/8/22/89

Lab Sample ID: 890622-029

Sample wt/vol: 900 ML

Lab File ID: >C1500

Date Sampled:

Date Received 22-Jun-1989 09:10

% Moisture: not dec.

dec:

Date Analyzed: 13-Jul-1989

Material Description

Date of Report: 20-JUL-89

Number TICs found: 5

CONCENTRATION UNITS:
(UG/L or UG/KG) UG/L

CAS NO.	COMPOUND NAME	RT	CONC	Q
1.	UNKNOWN	7.08	21	J
2.	UNKNOWN	7.83	2	J
3.	TRINITROTOLUENE	24.27	120	J
4.	UNKNOWN	26.54	7	J
5.	UNKNOWN Amad. nitrotoluene	27.46	4	J

WMO
1-28-70

REVIEWED

By H. Wachte
8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

THERMO CCLT 2

Lab Name: Oak Ridge National Lab Contract: MR
 Lab Code: Case no: ORNL SAS No: 92805 SDG No: C713
 Matrix: (soil/water) WATER Lab Sample ID: 890622-030
 Sample wt/vol: 900 ML Lab File ID: >C1501
 Level: (low/med) LOW Date Received 22-Jun-1989
 % Moisture: not dec. dec. Date Analyzed: 13-Jul-1989
 Extraction: (Sepi/Cont/Sonc) CONT Date Extracted: 26-Jun-1989
 GPC Cleanup: (Y/N) N pH: 7.91 Dilution Factor: 1.05

CAS NO.	COMPOUND	CONCENTRATION UNITS: (UG/L or UG/KG) UG/L	2
108-95-2	phenol	11.00	U
111-44-4	bis(2-chloroethyl)ether	11.00	U
95-57-8	2-chlorophenol	11.00	U
541-73-1	1,3-dichlorobenzene	11.00	U
106-46-7	1,4-dichlorobenzene	11.00	U
100-51-6	benzyl alcohol	11.00	U
95-50-1	1,2-dichlorobenzene	11.00	U
95-48-7	2-methylphenol	11.00	U
108-60-1	bis(2-chloroisopropyl)ether	11.00	U
106-44-5	4-methylphenol	11.00	U
621-64-7	n-nitroso-di-n-propylamine	11.00	U
67-72-1	hexachloroethane	11.00	U
98-95-3	nitrobenzene	11.00	U
78-59-1	isophorone	11.00	U
88-75-5	2-nitrophenol	11.00	U
105-67-9	2,4-dimethylphenol	11.00	U
65-85-0	benzoic acid	53.00	U
111-91-1	bis(2-chloroethoxy)methane	11.00	U
120-83-2	2,4-dichlorophenol	11.00	U
120-82-1	1,2,4-trichlorobenzene	11.00	U
91-20-3	naphthalene	11.00	U
106-47-8	4-chloroaniline	11.00	U
87-68-3	hexachlorobutadiene	11.00	U
59-50-7	4-chloro-3-methylphenol	11.00	U
91-57-6	2-methylnaphthalene	11.00	U
77-47-4	hexachlorocyclopentadiene	11.00	U
88-06-2	2,4,6-trichlorophenol	11.00	U
95-95-4	2,4,5-trichlorophenol	53.00	U
91-58-7	2-chloronaphthalene	11.00	U
88-74-4	2-nitroaniline	53.00	U
131-11-3	dimethylphthalate	11.00	U
208-96-8	acenaphthylene	11.00	U
606-20-2	2,6-dinitrotoluene	11.00	U

ff wachte
 Date 8/8/89

Semi-volatile Organic Analysis Data

EPA SAMPLE NO.

THERMO CCLT 2

Lab Name: Oak Ridge National Lab Contract: NR

Lab Code: Case no: ORNL SAS No: 92805 SDG No: C713

Matrix: (soil/water) WATER Lab Sample ID: 890622-030

Sample wt/vol: 900 ML Lab File ID: >C1501

Level: (low/med) LOW Date Received: 22-Jun-1989

% Moisture: not dec. dec. Date Analyzed: 13-Jul-1989

Extraction: (Sep/Cont/Sonc) COMT Date Extracted: 26-Jun-1989

GPC Cleanup: (Y/N) N pH: 7.91 Dilution Factor: 1.05

CAS NO.	COMPOUND	CONCENTRATION UNITS:	
		(UG/L or UG/KG) UG/L	Q
99-09-2-----	3-nitroaniline	53.00	U
83-32-9-----	acenaphthene	11.00	U
51-28-5-----	2,4-dinitrophenol	53.00	U
100-02-7-----	4-nitrophenol	53.00	U
132-64-9-----	dibenzofuran	11.00	U
121-14-2-----	2,4-dinitrotoluene	11.00	U
84-66-2-----	diethylphthalate	11.00	U
7005-72-3-----	4-chlorophenyl-phenylether	11.00	U
86-73-7-----	fluorene	11.00	U
100-01-6-----	4-nitroaniline	53.00	U
534-52-1-----	4,6-dinitro-2-methylphenol	53.00	U
86-30-6-----	n-nitrosodiphenylamine (1)	11.00	U
101-55-3-----	4-bromophenyl-phenylether	11.00	U
118-74-1-----	hexachlorobenzene	11.00	U
87-86-5-----	pentachlorophenol	53.00	U
85-01-8-----	phenanthrene	11.00	U
120-12-7-----	anthracene	11.00	U
84-74-2-----	di-n-butylphthalate	11.00	U
206-44-0-----	fluoranthene	11.00	U
129-00-0-----	pyrene	11.00	U
85-68-7-----	butylbenzylphthalate	11.00	U
91-94-1-----	3,3'-dichlorobenzidine	21.00	U
56-55-3-----	benzo(a)anthracene	11.00	U
218-01-9-----	chrysene	11.00	U
117-81-7-----	bis(2-ethylhexyl)phthalate	11.00	U
117-84-0-----	di-n-octylphthalate	11.00	U
205-99-2-----	benzo(b)fluoranthene	11.00	U
207-08-9-----	benzo(k)fluoranthene	11.00	U
50-32-8-----	benzo(a)pyrene	11.00	U
193-39-5-----	indeno(1,2,3-cd)pyrene	11.00	U
53-70-3-----	dibenz(a,h)anthracene	11.00	U
191-24-2-----	benzo(g,h,i)perylene	11.00	U

(1) - Cannot be separated from Diphenylamine

REVIEWED
Sf Wichter
 8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

Request Number: QAL92805

THERMO CCLT 2

Procedure Number: 8270

Matrix: WATER

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/8/22/89

Lab Sample ID: 890622-030

Sample wt/vol: 900 ML

Lab File ID: >C1501

Date Sampled:

Date Received 22-Jun-1989 09:10

% Moisture: not dec.

dec:

Date Analyzed: 13-Jul-1989

Material Description

Date of Report: 20-JUL-89

Number TICs found: 6

CONCENTRATION UNITS:
(UG/L or UG/KG) UG/L

CAS NO.	COMPOUND NAME	RT	CONC	Q
1.	UNKNOWN	7.09	40	J
2.	UNKNOWN	7.82	3	J
3.	TRINITROTOLUENE	24.27	85	J
4.	UNKNOWN	26.53	6	J
5.	UNKNOWN Aminodinitrotoluene	27.46	3	J
6.	UNKNOWN	29.06	3	J

W 46
1-28.50

REVIEWED

H. W. Richter
8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

BLANK CCLT

Lab Name: Oak Ridge National Lab Contract: MR

Lab Code: Case no: ORNL SAS No: 92805 SDG No: C713

Matrix: (soil/water) WATER Lab Sample ID: 890622-031

Sample wt/vol: 900 ML Lab File ID: >C1502

Level: (low/med) LOW Date Received 22-Jun-1989

% Moisture: not dec. dec. Date Analyzed: 13-Jul-1989

Extraction: (Sepf/Cont/Sonc) COMT Date Extracted: 26-Jun-1989

GPC Cleanup: (Y/N) N pH: 7.21 Dilution Factor: 1.05

CAS NO.	COMPOUND	CONCENTRATION UNITS:	
		(UG/L or UG/KG) UG/L	Q
108-95-2	phenol	11.00	U
111-44-4	bis(2-chloroethyl)ether	11.00	U
95-57-8	2-chlorophenol	11.00	U
541-73-1	1,3-dichlorobenzene	11.00	U
106-46-7	1,4-dichlorobenzene	11.00	U
100-51-6	benzyl alcohol	11.00	U
95-50-1	1,2-dichlorobenzene	11.00	U
95-48-7	2-methylphenol	11.00	U
108-60-1	bis(2-chloroisopropyl)ether	11.00	U
106-44-5	4-methylphenol	11.00	U
621-64-7	n-nitroso-di-n-propylamine	11.00	U
67-72-1	hexachloroethane	11.00	U
98-95-3	nitrobenzene	11.00	U
78-59-1	isophorone	11.00	U
88-75-5	2-nitrophenol	11.00	U
105-67-9	2,4-dimethylphenol	11.00	U
65-85-0	benzoic acid	53.00	U
111-91-1	bis(2-chloroethoxy)methane	11.00	U
120-83-2	2,4-dichlorophenol	11.00	U
120-82-1	1,2,4-trichlorobenzene	11.00	U
91-20-3	naphthalene	11.00	U
106-47-8	4-chloroaniline	11.00	U
87-68-3	hexachlorobutadiene	11.00	U
59-50-7	4-chloro-3-methylphenol	11.00	U
91-57-6	2-methylnaphthalene	11.00	U
77-47-4	hexachlorocyclopentadiene	11.00	U
88-06-2	2,4,6-trichlorophenol	11.00	U
95-95-4	2,4,5-trichlorophenol	53.00	U
91-58-7	2-chloronaphthalene	11.00	U
88-74-4	2-nitroaniline	53.00	U
131-11-3	dimethylphthalate	11.00	U
208-96-8	acenaphthylene	11.00	U
606-20-2	2,6-dinitrotoluene	11.00	U

REVIEWED

By

Date

Semi-volatile Organic Analysis Data

EPA SAMPLE NO.

Lab Name: Oak Ridge National Lab Contract: NR
Lab Code: Case no: ORNL SAS No: 92805 SDG No: C713
Matrix: (soil/water) WATER Lab Sample ID: 890622-031
Sample wt/vol: 900 ML Lab File ID: >C1502
Level: (low/med) LOW Date Received 22-Jun-1989
% Moisture: not dec. dec. Date Analyzed: 13-Jul-1989
Extraction: (Sepf/Cont/Sonc) CONT Date Extracted: 26-Jun-1989
GPC Cleanup: (Y/N) N pH: 7.21 Dilution Factor: 1.05

BLANK CCLT

CAS NO.	COMPOUND	CONCENTRATION UNITS:		Q
		(UG/L or UG/KG)	UG/L	
99-09-2	3-nitroaniline	53.00	U	
83-32-9	acenaphthene	11.00	U	
51-28-5	2,4-dinitrophenol	53.00	U	
100-02-7	4-nitrophenol	53.00	U	
132-64-9	dibenzofuran	11.00	U	
121-14-2	2,4-dinitrotoluene	11.00	U	
84-66-2	diethylphthalate	11.00	U	
7005-72-3	4-chlorophenyl-phenylether	11.00	U	
86-73-7	fluorene	11.00	U	
100-01-6	4-nitroaniline	53.00	U	
534-52-1	4,6-dinitro-2-methylphenol	53.00	U	
86-30-6	n-nitrosodiphenylamine (1)	11.00	U	
101-55-3	4-bromophenyl-phenylether	11.00	U	
118-74-1	hexachlorobenzene	11.00	U	
87-86-5	pentachlorophenol	53.00	U	
85-01-8	phenanthrene	11.00	U	
120-12-7	anthracene	11.00	U	
84-74-2	di-n-butylphthalate	11.00	U	
206-44-0	fluoranthene	11.00	U	
129-00-0	pyrene	11.00	U	
85-68-7	butylbenzylphthalate	11.00	U	
91-94-1	3,3'-dichlorobenzidine	21.00	U	
56-55-3	benzo(a)anthracene	11.00	U	
218-01-9	chrysene	11.00	U	
117-81-7	bis(2-ethylhexyl)phthalate	11.00	U	
117-84-0	di-n-octylphthalate	11.00	U	
205-99-2	benzo(b)fluoranthene	11.00	U	
207-08-9	benzo(k)fluoranthene	11.00	U	
50-32-8	benzo(a)pyrene	11.00	U	
193-39-5	indeno(1,2,3-cd)pyrene	11.00	U	
53-70-3	dibenz(a,h)anthracene	11.00	U	
191-24-2	benzo(g,h,i)perylene	11.00	U	

(1) - Cannot be separated from Diphenylamine

By: *S. J. Lenhardt*
Date: 8/8/89

Semivolatile Organic Analysis Data

EPA SAMPLE NO.

Request Number: 0AL92805

BLANK CCLT

Procedure Number: 8270

Matrix: WATER

Series:

Frequency:

Charge Number: 33900213

Customer Name: GRIEST/8/22/89

Lab Sample ID: 890622-031

Sample Wt/vol: 900 ML

Lab File ID: >C1502

Date Sampled:

Date Received 22-Jun-1989 09:10

% Moisture: not dec.

dec:

Date Analyzed: 13-Jul-1989

Material Description

Date of Report: 20-JUL-89

Number TICs found: 2

CONCENTRATION UNITS:
(UG/L or UG/KG) UG/L

CAS NO.	COMPOUND NAME	RT	CONC	P
1.	UNKNOWN	7.10	9	J
2.	UNKNOWN	7.82	3	J

REVIEWED

Lf Wichter
8/8/89

2E

WATER PESTICIDE SURROGATE RECOVERY

LAB NAME: OAK RIDGE NATIONAL LABORATORY CONTRACT: NA
LAB CODE: ORNL CASE NO.: NA SAS NO.: NA SDG NO.: 2805R

EPA		S1		
SAMPLE NO.		(DCBP)	#	OTHER #
1	MESO CCLT1		83	
2	MESO CCLT 2		77	
3	THERMO CCLT 1		87	
4	THERMO CCLT 2		71	
5	BLANK CCLT		71	
6				
7				
8				
9				
10				
11				
12				
13				
14				
15				
16				
17				
18				
19				
20				
21				
22				
23				
24				
25				
26				
27				
28				
29				
30				

ADVISORY QC LIMITS (20-150)

S1 (DCBP)= Decachlorobiphenyl
Column used to flag recovery values
* Value outside QC limits
+ No surrogate detected in sample
D Surrogates diluted out

MESO CCLT 1

EPA SAMPLE NO.

Lab Name: Oak Ridge National Lab Contract: ORNL

CCLT Leachate
of NR
Masspulv ContLab Code: Case no: 92805
Matrix: (soil/water) WATER

SAS No: 2805R

SDG No:

Lab Sample ID: 890622-027

Aliquot 1

LONG
5-24-89

Sample wt/vol: 900 ML

Lab File ID: NR

Level: (low/med) LOW

Date Received 22-Jun-1989

% Moisture: not dec. dec.

Date Analyzed: 14-Jul-1989

Extraction: (Sepf/Cont/Sonc) CONT

Date Extracted: 26-Jun-1989

GPC Cleanup: (Y/N) N

pH: 7.8

Dilution Factor: 1.0

CAS NO.	COMPOUND	CONCENTRATION UNITS: (ug/L or ug/Kg) UG/L	Q
319-84-6	ALPHA-BHC	0.06	U
319-85-7	BETA-BHC	0.24	J
319-86-8	DELTA-BHC	0.06	U
58-89-9	GAMMA-BHC (LINDANE)	0.19	
76-44-8	HEPTACHLOR	0.06	U
309-00-2	ALDRIN	0.06	U
1024-57-3	HEPTACHLOR EPOXIDE	0.06	U
959-98-8	ENDOSULFAM I	0.06	U
60-57-1	DIELDRIN	0.01	J
72-55-9	4,4'-DDE	0.11	U
72-20-8	ENDRIN	0.11	U
33213-65-9	ENDOSULFAM II	0.11	U
72-54-8	4,4'-DDD	0.11	U
1031-07-8	ENDOSULFAM SULFATE	0.11	U
50-29-3	4,4'-DDT	0.11	U
72-43-5	METHOXYCHLOR	0.60	U
53494-70-5	ENDRIN KETONE	0.11	U
5103-71-9	ALPHA CHLORDANE	0.60	U
5103-74-2	GAMMA CHLORDANE	0.12	
8001-35-2	TOXAPHENE	1.10	U
12674-11-2	AROCLOR-1016	0.60	U
11104-28-2	AROCLOR-1221	0.60	U
11141-16-5	AROCLOR-1232	0.60	U
53469-21-9	AROCLOR-1242	0.60	U
12672-29-6	AROCLOR-1248	0.60	U
11097-69-1	AROCLOR-1254	1.10	U
11096-82-5	AROCLOR-1260	1.10	U

REVIEWED

by J. W. Wachtel
Date 8/8/89

MESO CCLT 2

EPA SAMPLE NO.

Lab Name: Oak Ridge National Lab Contract: ORNL

Lab Code: Case no: 92805
Matrix: (soil/water) WATER

SAS No: 2805R

SDG No:

Lab Sample ID: 890622-028

Sample wt/vol: 900 ML

Lab File ID: NR

Level: (low/med) LOW

Date Received 22-Jun-1989

% Moisture: not dec. dec.

Date Analyzed: 14-Jul-1989

Extraction: (Sepf/Cont/Sonc) CONT

Date Extracted: 26-Jun-1989

GPC Cleanup: (Y/N) N

pH: 7.8

Dilution Factor: 1.0

CCLT Leachate
 of NR
 Mesophilic Consort
 (Aligent 2) 6-27-90

CAS NO.	COMPOUND	CONCENTRATION UNITS: (ug/L or ug/Kg) UG/L	Q
319-84-6	ALPHA-BHC	0.06	U
319-85-7	BETA-BHC	0.40	
319-86-8	DELTA-BHC	0.06	U
58-89-9	GAMMA-BHC (LINDANE)	0.30	
76-44-8	HEPTACHLOR	0.02	J
309-00-2	ALDRIN	0.06	U
1024-57-3	HEPTACHLOR EPOXIDE	0.06	U
959-98-8	ENDOSULFAM I	0.06	U
60-57-1	DIELDRIN	0.01	J
72-55-9	4,4'-DDE	0.11	U
72-20-8	ENDRIN	0.11	U
33213-65-9	ENDOSULFAM II	0.11	U
72-54-8	4,4'-DDD	0.11	U
1031-07-8	ENDOSULFAM SULFATE	0.11	U
50-29-3	4,4'-DDT	0.11	U
72-43-5	METHOXYCHLOR	0.60	U
53494-70-5	ENDRIN KETONE	0.11	U
5103-71-9	ALPHA CHLORDANE	0.60	J
5103-74-2	GAMMA CHLORDANE	0.12	
8001-35-2	TOXAPHENE	1.10	U
12674-11-2	AROCLOR-1016	0.60	U
11104-28-2	AROCLOR-1221	0.60	U
11141-16-5	AROCLOR-1232	0.60	U
53469-21-9	AROCLOR-1242	0.60	U
12672-29-6	AROCLOR-1248	0.60	U
11097-69-1	AROCLOR-1254	1.10	U
11096-82-5	AROCLOR-1260	1.10	U

REMOVED

Sp. W. Richter
 8/8/89

EPA SAMPLE NO.

THERMO CCLT 1

Lab Name: Oak Ridge National Lab Contract: ORNL

Lab Code: Case no: 92805 SAS No: 2805R SDG No: *CCLT leachate of NR Thermophilic Count (Aliquot) wt% 5-27-80*

Matrix: (soil/water) WATER Lab Sample ID: 890622-029

Sample wt/vol: 900 ML Lab File ID: NR

Level: (low/med) LOW Date Received 22-Jun-1989

% Moisture: not dec. dec. Date Analyzed: 14-Jul-1989

Extraction: (Sepf/Cont/Sonc) COMT Date Extracted: 26-Jun-1989

GPC Cleanup: (Y/N) N pH: 7.8 Dilution Factor: 1.0

CAS NO.	COMPOUND	CONCENTRATION UNITS: (ug/L or ug/Kg) UG/L	2
319-84-6	ALPHA-BHC	0.06	U
319-85-7	BETA-BHC	0.33	
319-86-8	DELTA-BHC	0.06	U
58-89-9	GAMMA-BHC (LINDANE)	0.18	
76-44-8	HEPTACHLOR	0.06	U
309-00-2	ALDRIN	0.06	U
1024-57-3	HEPTACHLOR EPOXIDE	0.06	U
959-98-8	ENDOSULFAM I	0.06	U
60-57-1	DIELDRIN	0.01	J
72-55-9	4,4'-DDE	0.11	U
72-20-8	ENDRIN	0.11	U
33213-65-9	ENDOSULFAM II	0.11	U
72-54-8	4,4'-DDD	0.11	U
1031-07-8	ENDOSULFAM SULFATE	0.11	U
50-29-3	4,4'-DDT	0.11	U
72-43-5	METHOXYCHLOR	0.60	U
53494-70-5	ENDRIN KETONE	0.11	U
5103-71-9	ALPHA CHLORDANE	0.60	U
5103-74-2	GAMMA CHLORDANE	0.06	J
8001-35-2	TOXAPHENE	1.10	U
12674-11-2	AROCLOR-1016	0.60	U
11104-28-2	AROCLOR-1221	0.60	U
11141-16-5	AROCLOR-1232	0.60	U
53469-21-9	AROCLOR-1242	0.60	U
12672-29-6	AROCLOR-1248	0.60	U
11097-69-1	AROCLOR-1254	1.10	U
11096-82-5	AROCLOR-1260	1.10	U

REVIEWED

by *J. K. Kuchter*
Date 8/8/89

THERMO CCLT 2

EPA SAMPLE NO.

Lab Name: Oak Ridge National Lab

Contract: ORNL

CCLT Leachate
of NR
Thermophilic Compost
(Aligant) WPG
1-11-90

Lab Code: Case no: 92805
Matrix: (soil/water) WATERSAS No: 2805R SDG No:
Lab Sample ID: 890622-030

Sample wt/vol: 900 ML

Lab File ID: NR

Level: (low/med) LOW

Date Received 22-Jun-1989

% Moisture: not dec. dec.

Date Analyzed: 14-Jul-1989

Extraction: (Sepf/Cont/Sonc) COMT

Date Extracted: 26-Jun-1989

GPC Cleanup: (Y/N) N pH: 7.9

Dilution Factor: 1.0

CAS NO.	COMPOUND	CONCENTRATION UNITS:		Q
		(ug/L or ug/Kg)	UG/L	
319-84-6	ALPHA-BHC	0.06	U	
319-85-7	BETA-BHC	0.30	U	
319-86-8	DELTA-BHC	0.06	U	
58-89-9	GAMMA-BHC (LINDANE)	0.17	U	
76-44-8	HEPTACHLOR	0.01	J	
309-00-2	ALDRIN	0.06	U	
1024-57-3	HEPTACHLOR EPOXIDE	0.06	U	
959-98-8	ENDOSULFAM I	0.06	U	
60-57-1	DIELDRIN	0.01	J	
72-55-9	4,4'-DDE	0.11	U	
72-20-8	ENDRIN	0.11	U	
33213-65-9	ENDOSULFAM II	0.11	U	
72-54-8	4,4'-DDD	0.11	U	
1031-07-8	ENDOSULFAM SULFATE	0.11	U	
50-29-3	4,4'-DDT	0.11	U	
72-43-5	METHOXYCHLOR	0.60	U	
53494-70-5	ENDRIN KETONE	0.11	U	
5103-71-9	ALPHA CHLORDANE	0.60	U	
5103-74-2	GAMMA CHLORDANE	0.04	J	
8001-35-2	TOXAPHENE	1.10	U	
12674-11-2	AROCLOR-1016	0.60	U	
11104-28-2	AROCLOR-1221	0.60	U	
11141-16-5	AROCLOR-1232	0.60	U	
53469-21-9	AROCLOR-1242	0.60	U	
12672-29-6	AROCLOR-1248	0.60	U	
11097-69-1	AROCLOR-1254	1.10	U	
11096-82-5	AROCLOR-1260	1.10	U	

REVIEWED

Sp. Hachte
Date 8/8/89

BLANK CCLT

EPA SAMPLE NO.

CCLT Blank
NR 1-22-90

Lab Name: Oak Ridge National Lab Contract: ORNL

 Lab Code: Case no: 92805 SAS No: 2805R SDG No:
 Matrix: (soil/water) WATER Lab Sample ID: 890622-031

Sample Wt/vol: 900 ML Lab File ID: NR

Level: (low/med) LOW Date Received 22-Jun-1989

% Moisture: not dec. dec. Date Analyzed: 14-Jul-1989

Extraction: (Sepi/Cont/Sonc) COMT Date Extracted: 26-Jun-1989

GPC Cleanup: (Y/N) N pH: 7.2 Dilution Factor: 1.0

CAS NO.	COMPOUND	CONCENTRATION UNITS:	
		(ug/L or ug/Kg) UG/L	g
319-84-6	ALPHA-BHC	0.06	U
319-85-7	BETA-BHC	0.06	U
319-86-8	DELTA-BHC	0.06	U
58-89-9	GAMMA-BHC (LINDANE)	0.06	U
76-44-8	HEPTACHLOR	0.06	U
309-00-2	ALDRIN	0.06	U
1024-57-3	HEPTACHLOR EPOXIDE	0.06	U
959-98-8	ENDOSULFAM I	0.06	U
60-57-1	DIELDRIN	0.11	U
72-55-9	4,4'-DDE	0.11	U
72-20-8	ENDRIN	0.11	U
33213-65-9	ENDOSULFAM II	0.11	U
72-54-8	4,4'-DDD	0.11	U
1031-07-8	ENDOSULFAM SULFATE	0.11	U
50-29-3	4,4'-DDT	0.11	U
72-43-5	METHOXYCHLOR	0.60	U
53494-70-5	ENDRIN KETONE	0.11	U
5103-71-9	ALPHA CHLORDANE	0.60	U
5103-74-2	GAMMA CHLORDANE	0.60	U
8001-35-2	TOXAPHENE	1.10	U
12674-11-2	AROCLOR-1016	0.60	U
11104-28-2	AROCLOR-1221	0.60	U
11141-16-5	AROCLOR-1232	0.60	U
53469-21-9	AROCLOR-1242	0.60	U
12672-29-6	AROCLOR-1248	0.60	U
11097-69-1	AROCLOR-1254	1.10	U
11096-82-5	AROCLOR-1260	1.10	U

REVIEWED

df uschte
 8/8/89

2A
WATER VOLATILE SURROGATE RECOVERY

b Name: OAK RIDGE NATIONAL LAB Contract: 1.0

b Code: NR Case No.: ORNL SAS No.: NR SDG No.: G711

	EPA SAMPLE NO.	S1 (TOL) *	S2 (BFB) *	S3 (DCE) *	OTHER	TOT OUT
01	VBLK630	94	101	94		0
02	890707-033	106	99	106		0
03	890630-071	35.84 *	38 *	84		2
04	890630-072	39.87 *	44 *	87		2
05	890630-074	7.88 *	46 *	86		2
06	890630-075	88	86	88		10
07						
08						
09						
10						
11						
12						
13						
14						
15						
16						
17						
18						
19						
20						
21						
22						
23						
24						
25						
26						
27						
28						
29						
30						

*NOTE: SAMPLES
THAT ARE OUT
OF CONTROL
WERE PRESPIKED
BY CUSTOMER.

MOG
7/17/94

QC LIMITS
S1 (TOL) = Toluene-d8 (88-110)
S2 (BFB) = Bromofluorobenzene (86-115)
S3 (DCE) = 1,2-Dichloroethane-d4 (76-114)

- * Column to be used to flag recovery values
- * Values outside of contract required QC limits
- D Surrogates diluted out

de 1 of 1

FORM II VOA-1

1/87 Rev.
MDe
7/3/94

2A
WATER VOLATILE SURROGATE RECOVERY

Lab Name: OAK RIDGE NATIONAL LAB Contract: 16667

Lab Code: NR Case No.: ORNL SAS No.: NR SDG No.: G712

	EPA SAMPLE NO.	S1 (TOL) #	S2 (BFB) #	S3 (DCE) #	OTHER	TOT OUT
01	VBLK630	99	99	98		0
02	890630-073	11 *	37 *	79		2
03	890629-051	99	95	100		0
04	890706-105	109	101	92		0
05						
06						
07						
08						
09						
10						
11						
12						
13						
14						
15						
16						
17						
18						
19						
20						
21						
22						
23						
24						
25						
26						
27						
28						
29						
30						

*SAMPLE WAS
PRESPIKED BY
CUSTOMER.

QC LIMITS
S1 (TOL) = Toluene-d8 (88-110)
S2 (BFB) = Bromofluorobenzene (86-115)
S3 (DCE) = 1,2-Dichloroethane-d4 (76-114)

* Column to be used to flag recovery values

* Values outside of contract required QC limits

D Surrogates diluted out

VOLATILE ORGANICS ANALYSIS DATA SHEET

EPA SAMPLE NO.

MESO CCLT-1

Lab Name: Oak Ridge National Lab Contract: NR

Lab Code: Case no: ORML SAS No: 92819 SDG No: G711

Matrix: (soil/water) WATER Lab Sample ID: 890630-071

Sample wt/vol: 5 ML Lab File ID: PG0004

Level: (low/med) LOW Date Received 30-Jun-1989

% Moisture: not dec. Date Analyzed: 11-Jul-1989

Column: (pack/cap) PACK Dilution Factor: 1.0

CAS NO.	COMPOUND	CONCENTRATION UNITS: (UG/L or UG/KG) UG/L	Q
74-87-3	chloromethane	10.00	U
74-83-9	bromomethane	10.00	U
75-01-4	vinyl chloride	10.00	U
75-00-3	chloroethane	10.00	U
75-09-2	methylene chloride	9.00	B
67-64-1	acetone	15.00	B
75-15-0	carbon disulfide	5.00	U
75-35-4	1,1-dichloroethene	5.00	U
75-34-3	1,1-dichloroethane	5.00	U
540-59-0	1,2-dichloroethene (total)	5.00	U
67-66-3	chloroform	2.00	J
107-06-2	1,2-dichloroethane	5.00	U
78-93-3	2-butanone	10.00	U
71-55-6	1,1,1-trichloroethane	5.00	U
56-23-5	carbon tetrachloride	5.00	U
108-05-4	vinyl acetate	10.00	U
75-27-4	bromodichloromethane	5.00	U
78-87-5	1,2-dichloropropane	5.00	U
10061-01-5	cis-1,3-dichloropropene	5.00	U
79-01-6	trichloroethene	5.00	U
124-48-1	dibromochloromethane	5.00	U
79-00-5	1,1,2-trichloroethane	5.00	U
71-43-2	benzene	5.00	U
10061-02-06	trans-1,3-dichloropropene	5.00	U
75-25-2	bromoform	5.00	U
108-10-1	4-methyl-2-pentanone	10.00	U
591-78-6	2-hexanone	10.00	U
127-18-4	tetrachloroethene	5.00	U
79-34-5	1,1,2,2-tetrachloroethane	5.00	U
108-88-3	toluene	5.00	U
108-90-7	chlorobenzene	5.00	U
100-41-4	ethylbenzene	5.00	U
100-42-5	styrene	5.00	U
1330-20-7	xylene (total)	5.00	U

REVIEWED

Date

8/8/89

VOLATILE ORGANICS ANALYSIS DATA SHEET

EPA SAMPLE NO

MESO CCLT-2

Lab Name: Oak Ridge National Lab Contract: MR
Lab Code: Case no: ORNL SAS No: 92819 SDG No: G711
Matrix: (soil/water) WATER Lab Sample ID: 890630-072
Sample wt/vol: 5 ML Lab File ID: >G0005
Level: (low/med) LOW Date Received 30-Jun-1989
% Moisture: not dec. Date Analyzed: 11-Jul-1989
Column: (pack/cap) PACK Dilution Factor: 1.0

CAS NO.	COMPOUND	CONCENTRATION UNITS: (UG/L or UG/KG) UG/L	Q
74-87-3	chloromethane	10.00	U
74-83-9	bromomethane	10.00	U
75-01-4	vinyl chloride	10.00	U
75-00-3	chloroethane	10.00	U
75-09-2	methylene chloride	35.00	B
67-64-1	acetone	19.00	B
75-15-0	carbon disulfide	5.00	U
75-35-4	1,1-dichloroethene	5.00	U
75-34-3	1,1-dichloroethane	5.00	U
540-59-0	1,2-dichloroethane (total)	5.00	U
67-66-3	chloroform	2.00	J
107-06-2	1,2-dichloroethane	5.00	U
78-93-3	2-butanone	10.00	U
71-55-6	1,1,1-trichloroethane	5.00	U
56-23-5	carbon tetrachloride	5.00	U
108-05-4	vinyl acetate	10.00	U
75-27-4	bromodichloromethane	5.00	U
78-87-5	1,2-dichloropropane	5.00	U
10061-01-5	cis-1,3-dichloropropene	5.00	U
79-01-6	trichloroethene	5.00	U
124-48-1	dibromochloromethane	5.00	U
79-00-5	1,1,2-trichloroethane	5.00	U
71-43-2	benzene	5.00	U
10061-02-06	trans-1,3-dichloropropene	5.00	U
75-25-2	bromoform	5.00	U
108-10-1	4-methyl-2-pentanone	10.00	U
591-78-6	2-hexanone	10.00	U
127-18-4	tetrachloroethene	5.00	U
79-34-5	1,1,2,2-tetrachloroethane	5.00	U
108-88-3	toluene	5.00	U
108-90-7	chlorobenzene	5.00	U
100-41-4	ethylbenzene	5.00	U
100-42-5	styrene	5.00	U
1330-20-7	xylene (total)	5.00	U

REVIEWED

BY

Date

VOLATILE ORGANICS ANALYSIS DATA SHEET

EPA SAMPLE NO.

THERMO CCLT-1

Lab Name: Oak Ridge National Lab Contract: NR

Lab Code: Case no: ORNL SAS No: 92819 SDG No: G712

Matrix: (soil/water) WATER

Lab Sample ID: 890630-073

Sample wt/vol: 5 ML

Lab File ID: >G0012

Level: (low/med) LOW

Date Received 30-Jun-1989

% Moisture: not dec.

Date Analyzed: 12-Jul-1989

Column: (pack/cap) PACK

Dilution Factor: 1.0

CAS NO.	COMPOUND	CONCENTRATION UNITS:		Q
		(UG/L or UG/KG)	UG/L	
74-87-3	chloromethane	10.00	U	
74-83-9	bromomethane	10.00	U	
75-01-4	vinyl chloride	10.00	U	
75-00-3	chloroethane	10.00	U	
75-09-2	methylene chloride	13.00	B	
67-64-1	acetone	9.00	JB	
75-15-0	carbon disulfide	5.00	U	
75-35-4	1,1-dichloroethene	5.00	U	
75-34-3	1,1-dichloroethane	5.00	U	
540-59-0	1,2-dichloroethene (total)	5.00	U	
67-66-3	chloroform	1.00	J	
107-06-2	1,2-dichloroethane	5.00	U	
78-93-3	2-butanone	10.00	U	
71-55-6	1,1,1-trichloroethane	5.00	U	
56-23-5	carbon tetrachloride	5.00	U	
108-05-4	vinyl acetate	10.00	U	
75-27-4	bromodichloromethane	5.00	U	
78-87-5	1,2-dichloropropane	5.00	U	
10061-01-5	cis-1,3-dichloropropene	5.00	U	
79-01-6	trichloroethene	5.00	U	
124-48-1	dibromochloromethane	5.00	U	
79-00-5	1,1,2-trichloroethane	5.00	U	
71-43-2	benzene	5.00	U	
10061-02-06	trans-1,3-dichloropropene	5.00	U	
75-25-2	bromoform	5.00	U	
108-10-1	4-methyl-2-pentanone	10.00	U	
591-78-6	2-hexanone	10.00	U	
127-18-4	tetrachloroethene	5.00	U	
79-34-5	1,1,2,2-tetrachloroethane	5.00	U	
108-88-3	toluene	5.00	U	
108-90-7	chlorobenzene	5.00	U	
100-41-4	ethylbenzene	5.00	U	
100-42-5	styrene	5.00	U	
1330-20-7	xylene (total)	5.00	U	

Date

8/8/89

VOLATILE ORGANICS ANALYSIS DATA SHEET

EPA SAMPLE NO.

Lab Name: Oak Ridge National Lab Contract: NR
Lab Code: Case no: ORNL SAS No: 92819 SPG No: G711
Matrix: (soil/water) WATER Lab Sample ID: 890630-074
Sample wt/vol: 5 ML Lab File ID: G0007
Level: (low/med) LOW Date Received 30-Jun-1989
% Moisture: not dec. Date Analyzed: 11-Jul-1989
Column: (pack/cap) PACK Dilution Factor: 1.0

THERMO CCLT-2

CAS NO.	COMPOUND	CONCENTRATION UNITS: (UG/L or UG/KG) UG/L	Q
74-87-3	chloromethane	10.00	U
74-83-9	bromomethane	10.00	U
75-01-4	vinyl chloride	10.00	U
75-00-3	chloroethane	10.00	U
75-09-2	methylene chloride	13.00	B
67-64-1	acetone	19.00	B
75-15-0	carbon disulfide	5.00	U
75-35-4	1,1-dichloroethene	5.00	U
75-34-3	1,1-dichloroethane	5.00	U
540-59-0	1,2-dichloroethene (total)	5.00	U
67-66-3	chloroform	1.00	J
107-06-2	1,2-dichloroethane	5.00	U
78-93-3	2-butanone	10.00	U
71-55-6	1,1,1-trichloroethane	5.00	U
56-23-5	carbon tetrachloride	5.00	U
108-05-4	vinyl acetate	10.00	U
75-27-4	bromodichloromethane	5.00	U
78-87-5	1,2-dichloropropane	5.00	U
10061-01-5	cis-1,3-dichloropropene	5.00	U
79-01-6	trichloroethene	5.00	U
124-48-1	dibromochloromethane	5.00	U
79-00-5	1,1,2-trichloroethane	5.00	U
71-43-2	benzene	5.00	U
10061-02-06	trans-1,3-dichloropropene	5.00	U
75-25-2	bromoform	5.00	U
108-10-1	4-methyl-2-pentanone	10.00	U
591-78-6	2-hexanone	10.00	U
127-18-4	tetrachloroethene	5.00	U
79-34-5	1,1,2,2-tetrachloroethane	5.00	U
108-88-3	toluene	5.00	U
108-90-7	chlorobenzene	5.00	U
100-41-4	ethylbenzene	5.00	U
100-42-5	styrene	5.00	U
1330-20-7	xylene (total)	5.00	U

REVIEWED

By

Date

8/8/89

VOLATILE ORGANICS ANALYSIS DATA SHEET

EPA SAMPLE NO.

BLANK CCLT

Lab Name: Oak Ridge National Lab Contract: NR

Lab Code: Case no: ORNL SAS No: 92819 SDG No: G711

Matrix: (soil/water) WATER

Lab Sample ID: 890630-075

Sample wt/vol: 5 ML

Lab File ID: >G0008

Level: (low/med) LOW

Date Received 30-Jun-1989

% Moisture: not dec.

Date Analyzed: 11-Jul-1989

Column: (pack/cap) PACK

Dilution Factor: 1.0

CAS NO.	COMPOUND	CONCENTRATION UNITS:		Q
		(UG/L or UG/KG)	UG/L	
74-87-3	-----chloromethane	10.00	U	
74-83-9	-----bromomethane	10.00	U	
75-01-4	-----vinyl chloride	10.00	U	
75-00-3	-----chloroethane	10.00	U	
75-09-2	-----methylene chloride	8.00	B	
67-64-1	-----acetone	23.00	B	
75-15-0	-----carbon disulfide	5.00	U	
75-35-4	-----1,1-dichloroethene	5.00	U	
75-34-3	-----1,1-dichloroethane	5.00	U	
540-59-0	-----1,2-dichloroethane (total)	5.00	U	
67-66-3	-----chloroform	2.00	J	
107-06-2	-----1,2-dichloroethane	5.00	U	
78-93-3	-----2-butanone	10.00	U	
71-55-6	-----1,1,1-trichloroethane	5.00	U	
56-23-5	-----carbon tetrachloride	5.00	U	
108-05-4	-----vinyl acetate	10.00	U	
75-27-4	-----bromodichloromethane	5.00	U	
78-87-5	-----1,2-dichloropropane	5.00	U	
10061-01-5	-----cis-1,3-dichloropropene	5.00	U	
79-01-6	-----trichloroethene	5.00	U	
124-48-1	-----dibromochloromethane	5.00	U	
79-00-5	-----1,1,2-trichloroethane	5.00	U	
71-43-2	-----benzene	5.00	U	
10061-02-06	-----trans-1,3-dichloropropene	5.00	U	
75-25-2	-----bromoform	5.00	U	
108-10-1	-----4-methyl-2-pentanone	10.00	U	
591-78-6	-----2-hexanone	10.00	U	
127-18-4	-----tetrachloroethene	5.00	U	
79-34-5	-----1,1,2,2-tetrachloroethane	5.00	U	
108-88-3	-----toluene	5.00	U	
108-90-7	-----chlorobenzene	5.00	U	
100-41-4	-----ethylbenzene	5.00	U	
100-42-5	-----styrene	5.00	U	
1330-20-7	-----xylene (total)	5.00	U	

REVIEWED

By AP/arkto
Date 8/8/89

Oak Ridge National Laboratory
Analytical Chemistry Division
Results of Analyses
Chemical and Physical Analysis

Customer Name M.M. GRIEST Date Received 22-Jun-1989 09:45
Request Number CPA9711 Charge Number 33900213 Approved By J.C. Prie
Project Number Dept Number 3390
Series Date of Report 9-AUG-89 Date 8-9-89

Analy No.	Customers Id	Date/Time Sampled	Analysis	Result	Units	Procedure Mo	Completed
1890622-036	ME50 CCLT 1						
MATRIX							
			AG	< 1.0E-02	MG/L	EPA 200.7	14-Jul-1989
			AL	4.5E+01	MG/L	EPA 200.7	14-Jul-1989
			AS	< 1.0E-01	MG/L	EPA 200.7	14-Jul-1989
			BA	6.8E-01	MG/L	EPA 200.7	14-Jul-1989
			BE	< 8.0E-04	MG/L	EPA 200.7	14-Jul-1989
			CA	7.2E+00	MG/L	EPA 200.7	14-Jul-1989
			CD	< 4.0E-03	MG/L	EPA 200.7	14-Jul-1989
			CO	6.1E-03	MG/L	EPA 200.7	14-Jul-1989
			CR	4.8E-02	MG/L	EPA 200.7	14-Jul-1989
			CU	4.7E-02	MG/L	EPA 200.7	14-Jul-1989
			FE	7.3E+00	MG/L	EPA 200.7	14-Jul-1989
			LI	< 3.0E+01	MG/L	EPA 200.7	14-Jul-1989
			MG	6.3E+00	MG/L	EPA 200.7	14-Jul-1989
			MN	2.2E-01	MG/L	EPA 200.7	14-Jul-1989
			MO	< 8.0E-02	MG/L	EPA 200.7	14-Jul-1989
			NA	2.9E+01	MG/L	EPA 200.7	14-Jul-1989
			NI	2.5E-02	MG/L	EPA 200.7	14-Jul-1989
			P	1.1E+01	MG/L	EPA 200.7	14-Jul-1989
			PB	< 6.0E-02	MG/L	EPA 200.7	14-Jul-1989
			SB	1.4E-01	MG/L	EPA 200.7	14-Jul-1989
			SE	< 1.6E-01	MG/L	EPA 200.7	14-Jul-1989
			SI	1.2E+02	MG/L	EPA 200.7	14-Jul-1989
			SN	< 1.0E-01	MG/L	EPA 200.7	14-Jul-1989
			SR	5.2E-02	MG/L	EPA 200.7	14-Jul-1989
			TI	1.1E+00	MG/L	EPA 200.7	14-Jul-1989
			V	7.3E-02	MG/L	EPA 200.7	14-Jul-1989
			ZM	4.5E-01	MG/L	EPA 200.7	14-Jul-1989
			ZR	6.0E-02	MG/L	EPA 200.7	14-Jul-1989

Analy No. Customers Id	Date/Time Sampled	Analysis	Result	Units	Procedure No	Completed
Matrix	Material Desc.					
Frequency						
1890622-037 MESO CCLT 2						
WATER						
		AG	< 1.0E-02	MG/L	EPA 200.7	14-Jul-1989
		AL	3.5E+01	MG/L	EPA 200.7	14-Jul-1989
		AS	< 1.0E-01	MG/L	EPA 200.7	14-Jul-1989
		BA	5.8E-01	MG/L	EPA 200.7	14-Jul-1989
		BE	< 8.0E-04	MG/L	EPA 200.7	14-Jul-1989
		CA	6.4E+00	MG/L	EPA 200.7	14-Jul-1989
		CD	< 4.0E-03	MG/L	EPA 200.7	14-Jul-1989
		CO	< 6.0E-03	MG/L	EPA 200.7	14-Jul-1989
		CR	3.3E-02	MG/L	EPA 200.7	14-Jul-1989
		CU	4.3E-02	MG/L	EPA 200.7	14-Jul-1989
		FE	2.3E+01	MG/L	EPA 200.7	14-Jul-1989
		LI	< 3.0E+01	MG/L	EPA 200.7	14-Jul-1989
		MG	5.3E+00	MG/L	EPA 200.7	14-Jul-1989
		MN	1.8E-01	MG/L	EPA 200.7	14-Jul-1989
		MO	< 8.0E-02	MG/L	EPA 200.7	14-Jul-1989
		NA	2.9E+01	MG/L	EPA 200.7	14-Jul-1989
		NI	1.8E-02	MG/L	EPA 200.7	14-Jul-1989
		P	1.0E+01	MG/L	EPA 200.7	14-Jul-1989
		PB	< 6.0E-02	MG/L	EPA 200.7	14-Jul-1989
		SB	1.3E-01	MG/L	EPA 200.7	14-Jul-1989
		SE	< 1.6E-01	MG/L	EPA 200.7	14-Jul-1989
		SI	1.0E+02	MG/L	EPA 200.7	14-Jul-1989
		SK	< 1.0E-01	MG/L	EPA 200.7	14-Jul-1989
		SR	4.2E-02	MG/L	EPA 200.7	14-Jul-1989
		TI	7.7E-01	MG/L	EPA 200.7	14-Jul-1989
		V	6.1E-02	MG/L	EPA 200.7	14-Jul-1989
		ZN	4.1E-01	MG/L	EPA 200.7	14-Jul-1989
		ZR	5.1E-02	MG/L	EPA 200.7	14-Jul-1989

131

Analy No. Customers Id		Date/Time Sampled					
Matrix		Material Desc.					
Frequency			Analysis	Result	Units	Procedure No	Completed

1890622-039 THERMO CCLT 2							
WATER							
	AG	< 1.0E-02	MG/L	EPA 200.7	14-Jul-1989		
	AL	6.4E+01	MG/L	EPA 200.7	14-Jul-1989		
	AS	< 1.0E-01	MG/L	EPA 200.7	14-Jul-1989		
	BA	6.9E-01	MG/L	EPA 200.7	14-Jul-1989		
	BE	< 8.0E-04	MG/L	EPA 200.7	14-Jul-1989		
	CA	1.2E+01	MG/L	EPA 200.7	14-Jul-1989		
	CD	< 4.0E-03	MG/L	EPA 200.7	14-Jul-1989		
	CO	9.3E-03	MG/L	EPA 200.7	14-Jul-1989		
	CR	7.1E-02	MG/L	EPA 200.7	14-Jul-1989		
	CU	7.5E-02	MG/L	EPA 200.7	14-Jul-1989		
	FE	2.0E+01	MG/L	EPA 200.7	14-Jul-1989		
	LI	< 3.0E+01	MG/L	EPA 200.7	14-Jul-1989		
	MG	8.1E+00	MG/L	EPA 200.7	14-Jul-1989		
	MN	6.1E-01	MG/L	EPA 200.7	14-Jul-1989		
	MO	< 8.0E-02	MG/L	EPA 200.7	14-Jul-1989		
	NA	2.4E+01	MG/L	EPA 200.7	14-Jul-1989		
	NI	2.8E-02	MG/L	EPA 200.7	14-Jul-1989		
	P	8.8E+00	MG/L	EPA 200.7	14-Jul-1989		
	PB	6.5E-02	MG/L	EPA 200.7	14-Jul-1989		
	SB	2.0E-01	MG/L	EPA 200.7	14-Jul-1989		
	SE	< 1.6E-01	MG/L	EPA 200.7	14-Jul-1989		
	SI	1.6E+02	MG/L	EPA 200.7	14-Jul-1989		
	SN	< 1.0E-01	MG/L	EPA 200.7	14-Jul-1989		
	SR	9.2E-02	MG/L	EPA 200.7	14-Jul-1989		
	TI	1.4E+00	MG/L	EPA 200.7	14-Jul-1989		
	V	8.5E-02	MG/L	EPA 200.7	14-Jul-1989		
	ZN	5.7E-01	MG/L	EPA 200.7	14-Jul-1989		
	ZR	7.6E-02	MG/L	EPA 200.7	14-Jul-1989		

Analy No.	Customers Id	Date/Time Sampled	Analysis	Result	Units	Procedure No	Completed
Matrix	Material Desc.						
Frequency							
1890622-040	BLANK CCLT						
WATER							
	AG			< 1.0E-02	MG/L	EPA 200.7	14-Jul-1989
	AL			< 1.0E-01	MG/L	EPA 200.7	14-Jul-1989
	AS			< 1.0E-01	MG/L	EPA 200.7	14-Jul-1989
	BA			< 4.0E-03	MG/L	EPA 200.7	14-Jul-1989
	BE			< 8.0E-04	MG/L	EPA 200.7	14-Jul-1989
	CA			< 1.2E-01	MG/L	EPA 200.7	14-Jul-1989
	CD			< 4.0E-03	MG/L	EPA 200.7	14-Jul-1989
	CO			< 6.0E-03	MG/L	EPA 200.7	14-Jul-1989
	CR			< 6.0E-03	MG/L	EPA 200.7	14-Jul-1989
	CU			< 2.0E-02	MG/L	EPA 200.7	14-Jul-1989
	FE			< 2.0E-02	MG/L	EPA 200.7	14-Jul-1989
	LI			< 3.0E+01	MG/L	EPA 200.7	14-Jul-1989
	MG			< 2.0E-02	MG/L	EPA 200.7	14-Jul-1989
	MM			< 6.0E-03	MG/L	EPA 200.7	14-Jul-1989
	MO			< 8.0E-02	MG/L	EPA 200.7	14-Jul-1989
	NA			< 4.0E+00	MG/L	EPA 200.7	14-Jul-1989
	NI			< 1.2E-02	MG/L	EPA 200.7	14-Jul-1989
	P			< 6.0E-01	MG/L	EPA 200.7	14-Jul-1989
	PB			< 6.0E-02	MG/L	EPA 200.7	14-Jul-1989
	SB			< 8.0E-02	MG/L	EPA 200.7	14-Jul-1989
	SE			< 1.6E-01	MG/L	EPA 200.7	14-Jul-1989
	SI			< 4.0E-01	MG/L	EPA 200.7	14-Jul-1989
	SM			< 1.0E-01	MG/L	EPA 200.7	14-Jul-1989
	SR			< 1.0E-02	MG/L	EPA 200.7	14-Jul-1989
	TI			< 4.0E-02	MG/L	EPA 200.7	14-Jul-1989
	V			< 8.0E-03	MG/L	EPA 200.7	14-Jul-1989
	ZM			< 1.6E-02	MG/L	EPA 200.7	14-Jul-1989
	ZR			< 4.0E-02	MG/L	EPA 200.7	14-Jul-1989

End of data for Request Number CPA9711 Total pages = 5 Cust. Copy ☒ File Copy ☐

APPENDIX A-5

TOXICOLOGY LABORTORY REPORTS FOR FATHEAD MINNOW LARVAE AND
CERIODAPHNIA DUBIA TESTS

TOXICITY TEST REPORT

TOXICOLOGY LABORATORY

ENVIRONMENTAL SCIENCES DIVISION
OAK RIDGE NATIONAL LABORATORY
P.O. BOX 2008, MS 6351
OAK RIDGE, TN 37831-6351

EXPERIMENT NO. C-431

Clean Closure Leachate Test
of
Thermophilic and Mesophilic Explosives-Contaminated Compost

June 22-29, 1989

STANDARD REPORT FORM

CERIODAPHNIA 7-DAY SURVIVAL AND REPRODUCTION TEST

Experiment no. C-431

Starting date: June 22, 1989.

Ending date: June 29, 1989.

1. SAMPLE

- 1.1 Sample description: Clean Closure Leachate Test leachates of thermophilic and mesophilic explosives-contaminated compost, and of artificial rainwater (AR) used to prepare the leachates.
- 1.2 Sampling point: Not applicable: Samples were collected from the laboratory where they were prepared.
- 1.3 Sampling date: Samples were picked up from the analytical chemistry laboratory on the morning of June 22, 1989, by A. J. Stewart.
- 1.4 Sampling method: Does not apply.
- 1.5 Sample was received at Environmental Sciences Division's toxicology laboratory on June 22, 1989 at 9:25 a.m., by A. J. Stewart; Registered Water Log Book A-103384, pp. 32.
- 1.6 Sample was used immediately.
- 1.7 Pretreatment: None.

2. TEST ORGANISMS

- 2.1 Species: Ceriodaphnia dubia.
- 2.2 Source: Environmental Sciences Division cultures.
- 2.3 Incubation water for cultures: Well/spring water.
- 2.4 Temperature of cultures: $25 \pm 2^{\circ}\text{C}$ (mean \pm SD)

3. TEST METHODS

- 3.1 Toxicity test method: Ceriodaphnia survival and reproduction test. Reference: EPA Test Method 1002.0, in W. B. Horning and C. I. Weber (eds.), Short-Term Methods for Estimating the Chronic Toxicity of Effluent and Receiving Water to Freshwater Organisms, EPA/600/4-85/014 (December 1985).
- 3.2 Deviations from reference: None.
- 3.3 Date/time test started: June 22, 1989; 12:40 p.m.
- 3.4 Date/time test terminated: June 29, 1989; 11:45 a.m.
- 3.5 Type of test chambers: 12 mm diameter x 75 mm borosilicate test tubes. Previous tests (C-416) showed that the 12 x 75 mm test chambers yielded results that were virtually identical to those for Ceriodaphnia in 17-mL polystyrene test chambers.
- Volume per chamber: 2.5 mL.
- 3.6 Number of Ceriodaphnia per test chamber: 1.
- 3.7 Number of replicates per treatment: 12.
- 3.8 Dilution/Control water: 1:9 (v:v) ratio of degassed mineral water to deionized distilled water.
- 3.9 Renewal period: 24 h.
- 3.10 Test temperature: Mean = 24.7°C; range = 24.5-24.9°C.
- 3.11 Treatment groups/concentrations: Control; Thermophillic 100%, 80%, 60%, 40%; Mesophillic 100%, 80%, 60%, 40%; and artificial rainwater 100%, 60%, and 40% of full-strength effluent.
- 3.12 Feeding regime during test: 35 uL of trout chow-yeast-cerophyl (TCO) mixture (EPA-600/4-85-014; section 7.10.6.2) per 15 mL of test solution every 24 h. The TCO mixture was augmented with the green alga, Ankistrodesmus falcatus (TCA lot TCA1-24-89; TCO lot 1-24-89).

4. QUALITY ASSURANCE

- 4.1 Standard toxicant used: Sodium lauryl sulfate (SLS).
- Source: Environmental Protection Agency, Environmental Monitoring and Support Laboratory, Cincinnati, OH.
- 4.2 Date/time of most recent test: June 14-15, 1989; 9:23 a.m.
- 4.3 Dilution water used: 1:9 (v:v) ratio of degassed mineral water to deionized distilled water augmented with trace metals.

4.4 24-h LC_{50} : 15.50 mg/L SLS; 95% C.L. = 12.35 - 18.17 mg/L SLS.

The LC_{50} was calculated by the moving average method. Reference: SAS User's Guide: Statistics, Version 5. Cary, NC: SAS Institute Inc., 1985, 956p. Raw data for the reference test is given in section 7.0.

4.5 Data from control chart prepared from standard toxicity tests:

Number of standard SLS reference tests completed by laboratory: 9. Central tendency: 10.00 ± 8.02 mg/L SLS (mean \pm 2 SD; moving average method).

4.6 Physical and chemical methods

The pH was measured by EPA method 150.1 with an Orion 700 pH meter. The meter was calibrated with pH 4.0 and pH 10.0 buffers.

Conductance ($\mu\text{mho/cm}$) was measured by EPA method 120.1 with a YSI model 31 meter. All values were corrected to 25°C.

Dissolved oxygen (mg/L) was measured by EPA method 360.1 with a YSI model 54AR dissolved oxygen meter. The meter was air calibrated.

Alkalinity was measured by titrating 50-mL samples with 0.01 N HCl to an endpoint pH of 4.5 (EPA method 130.1).

Hardness was determined by titrating 50-mL samples with EDTA to a colorimetric endpoint using Eriochrome Black T (EPA method 130.2).

Instruments were calibrated and standardized according to manufacturer's instructions.

All measurements were made on fresh samples before daily water replacement.

5. CERIODAPHNIA SURVIVAL AND FECUNDITY TEST RESULTS

5.1 Daily results from the Ceriodaphnia toxicity test:

Dilution	Day	Replicate ^a												Number of live Adults	Total live Young
		1	2	3	4	5	6	7	8	9	10	11	12		
Control	1	-	-	-	-	-	-	-	-	-	-	-	-	12	0
	2	-	-	-	-	-	-	-	-	-	-	-	-	12	0
	3	3	4	4	4	-	4	4	3	4	5	-	-	12	35
	4	0	x	6	0	8	8	8	6	0	7	6	2	11	51
	5	6	x	0	6	8	0	0	6	1	0	0	6	11	33
	6	6	x	9	7	2	4	8	6	10	8	9	8	11	77
	7	0	x	0	0	7	8	3	0	0	8	7	1	11	34
Total		15	4	19	17	25	24	23	21	15	28	22	17		230

5.1 Daily results from the Ceriodaphnia toxicity test (continued):

Dilution	Day	Replicate ^a												Number of live Adults	Total live Young
		1	2	3	4	5	6	7	8	9	10	11	12		
T-100%	1	-	-	-	-	-	-	-	-	-	-	-	-	12	0
	2	-	-	-	-	-	-	-	-	-	-	-	-	12	0
	3	-	-	-	-	-	-	-	-	-	-	-	-	12	0
	4	3	0	0	1	1	0	0	1	0	2	0	0	12	8
	5	0	0	0	0	3	0	x	0	0	0	0	0	11	3
	6	0	0	0	0	2	2	x	2	2	2	10	5	11	25
	7	4	5	0	5	0	3	x	4	0	2	3	8	11	34
	Total	7	5	0	6	6	5	0	7	2	6	13	13		70
T-80%	1	-	-	-	-	-	-	-	-	-	-	-	-	12	0
	2	-	-	-	-	-	-	-	-	-	-	-	-	12	0
	3	1	1	-	-	-	-	-	3	x	-	-	x	10	5
	4	3	3	3	2	0	4	2	1	x	2x	0	x	9	20
	5	0	4	0	0	3	2	x	0	x	x	0	x	8	9
	6	0	0	8	2	6	0	x	5	x	x	0	x	8	21
	7	3	1	3	5	6	3	x	7	x	x	3	x	8	31
	Total	7	9	14	9	15	9	2	16	0	2	3	0		86
T-60%	1	-	-	-	-	-	-	-	-	-	-	-	-	12	0
	2	-	-	-	-	-	-	-	-	-	-	-	-	12	0
	3	4	-	3	3	1	-	3	-	-	4	2	-	12	20
	4	4	1	4	1	4	6	5	5	7	4	1	0	12	42
	5	4	6	5	7	1	0	2	4	1	1	4	1	12	36
	6	0	0	0	9	7	4	0	8	6	6	6	0	12	46
	7	8	15	0	0	9	8	11	19	16	0	16	10	12	112
	Total	20	22	12	20	22	18	21	36	30	15	29	11		256
T-40%	1	-	-	-	-	-	-	-	-	-	-	-	-	12	0
	2	-	-	-	-	-	-	-	-	-	-	-	-	12	0
	3	4	4	2	4	3	3	2	4	2	M	4	-	12	32
	4	6	6	4	5	8	5	7	4	0	M	7	3	12	55
	5	0	11	5	5	0	0	0	0	8	M	0	6	12	35
	6	12	0	0	0	8	6	4	8	8	M	68	0	12	54
	7	5	11	11	0	11	16	15	16	14	M	14	0	12	113
	Total		27	32	22	14	30	30	28	32	32	0	33	9	289

Dilution	Day	Replicate ^a												Number of live Adults	Total live Young
		1	2	3	4	5	6	7	8	9	10	11	12		
M-100%	1	x	-	-	x	-	-	x	-	-	-	-	x	8	0
	2	x	-	-	x	x	-	x	-	-	-	-	x	7	0
	3	x	0	x	x	x	x	x	x	x	0	0	x	3	0
	4	x	x	x	x	x	x	x	x	x	x	x	x	0	0
	5	x	x	x	x	x	x	x	x	x	x	x	x	0	0
	6	x	x	x	x	x	x	x	x	x	x	x	x	0	0
	7	x	x	x	x	x	x	x	x	x	x	x	x	0	0
	Total		0	0	0	0	0	0	0	0	0	0	0	0	0
M-80%	1	-	-	-	-	-	-	-	-	-	-	-	-	12	0
	2	-	-	-	-	-	-	-	-	-	-	-	-	12	0
	3	2	-	-	-	-	-	-	-	-	-	-	-	12	2
	4	0	0	0	0	x	0	0	0	0	0	0	0	11	0
	5	0	0	0	0	x	0	0	0	0	0	0	0	11	0
	6	5	0	0	0	x	0	0	0	0	0	0	0	11	5
	7	7	0	0	0	x	0	0	0	0	0	0	0	11	7
	Total	14	0	0	0	0	0	0	0	0	0	0	0		14
M-60%	1	-	-	-	-	-	-	-	-	-	-	-	-	12	0
	2	-	-	-	-	-	-	-	-	-	-	-	-	12	0
	3	1	2	-	1	-	1	1	-	-	-	-	-	12	6
	4	0	2x	x	0	0	0	0	0	0	0	0	x	9	2
	5	0	x	x	0	0	0	0	0	0	0	0	x	9	0
	6	0	x	x	0	0	0	0	x	0	0	0	x	8	0
	7	3	x	x	0	1	6	x	x	0	0	0	x	7	10
	Total	4	4	0	1	1	7	1	0	0	0	0	0		18
M-40%	1	-	-	-	-	-	-	-	-	-	-	-	-	12	0
	2	-	-	-	-	-	-	-	-	-	-	-	-	12	0
	3	-	2	-	4	2	1	1	2	2	0	0	1	12	15
	4	0	0	0	0	0	3	0	4	x	0	3	0	11	10
	5	4	5	5	6	3	3	5	0	x	3	0	6	11	40
	6	2	0	0	8	6	6	x	4	x	5	5	8	10	44
	7	0	0	7	9	0	0	x	0	x	6	0	3	10	25
	Total	6	7	12	27	11	13	6	10	2	14	8	18		134
AR-60%	1	-	-	-	-	-	-	-	-	-	-	-	-	12	0
	2	-	-	-	-	-	-	-	-	-	-	-	-	12	0
	3	4	2	3	4	4	4	3	4	4	4	4	3	12	43
	4	8	6	8	0	0	0	0	0	0	0	6	x	11	28
	5	0	0	0	6	7	8	4	4	8	8	4	x	11	49
	6	4	6	7	7	6	4	7	6	6	6	4	x	11	63
	7	6	0	5	0	0	0	0	0	0	0	0	x	11	11
	Total	22	14	23	17	17	16	14	14	18	18	18	3		194

Dilution	Day	Replicate ^a												Number of live Adults	Total live Young
		1	2	3	4	5	6	7	8	9	10	11	12		
AR-40%	1	-	-	-	-	-	-	-	-	-	-	-	-	12	0
	2	-	-	-	-	-	-	-	4	1	-	-	-	12	5
	3	2	4	3	0	3	5	2	1	0	4	2	4	12	30
	4	5	9	0	0	0	0	5	8	1	0	6	x	11	34
	5	0	0	8	M	6	6	0	5	6	7	5	x	11	43
	6	6	8	6	M	9	5	3	6	4	4	0	x	11	51
	7	0	8	1	M	0	6	9	2	0	8	4	x	11	38
Total		13	29	18	0	18	22	19	26	12	23	17	4		201

^a - = live female too young to produce offspring;

x = dead adult, no young produced before death;

Nx = Dead adult, with no young produced before death;

M = Male.

5.2 Summary of results from the Ceriodaphnia toxicity tests:

Effluent concentration	Number of replicates	Number of females surviving for 7 d	Mean number of offspring per female (\pm SD)
Control	12	11	20.6 \pm 4.3
Thermophillic			
100%	12	11	6.4 \pm 3.9 ^a
80%	12	8	10.3 \pm 4.4 ^a
60%	12	12	21.3 \pm 7.4
40%	12	11	26.3 \pm 8.0
Mesophillic			
100%	12	0 ^c	-- \pm --
80%	12	11	1.3 \pm 4.2 ^a
60%	12	7	1.9 \pm 2.7 ^{ac}
40%	12	10	12.6 \pm 6.2 ^{ab}
Artificial Rainwater			
60%	12	11	17.4 \pm 3.0
40%	12	11	19.7 \pm 5.4

^aSurvival significantly lower than control, based on Fisher's Exact test.

^aFecundity is significantly lower than control.

^bFecundity is significantly lower than control and the 40% concentration of the thermophillic leachate and the 40% concentration of the artificial rainwater.

^cFecundity is significantly lower than control, the 60% concentration of the thermophillic leachate, and the 60% concentration of the artificial rainwater.

5.3 Statistical analyses: Control vs thermophillic leachate.

The data were first analyzed by using Fisher's Exact Test (for survival), followed by SAS-GLM (General Linear Models) procedure and Dunnett's Test (for reproduction), as recommended by Horning and Weber (1985). The GLM procedure is recommended over ANOVA when the data set contains missing values (note table in section 5.1).

Asterisks in the summary table (section 5.2) show concentrations significantly reducing survivorship of Ceriodaphnia based on Fisher's Exact Test, and/or concentrations significantly lowering fecundity based on Dunnett's test ($p = 0.05$, one tailed, $t_d = 2.22$).

For the data above, the mean number of offspring per female was computed for all surviving females. The least significant difference in fecundity at the 5% level ($LSD_{0.05}$) is 5.83 offspring per female, using the error MS value obtained with the GLM analysis (below). The LSD is a 28.3% reduction in fecundity compared to the control.

GLM Analysis of Ceriodaphnia Fecundity

Variance Source	D.F.	Sum of Squares	Mean Square	F	F _(0.05)
Among	4	3796.287	949.072	22.92	2.59
Within	54	2235.848		41.405	
Total	58	6032.135		(Pr > F = 0.0001)	

- 5.3 Statistical analyses: Control vs mesophillic leachate (all concentrations except for 100%, where survival was zero).

The data were first analyzed by using Fisher's Exact Test (for survival), followed by SAS-GLM (General Linear Models) procedure and Dunnett's Test (for reproduction), as recommended by Horning and Weber (1985). The GLM procedure is recommended over ANOVA when the data set contains missing values (note table in section 5.1).

Asterisks in the summary table (section 5.2, above), show concentrations significantly reducing survivorship of Ceriodaphnia based on Fisher's Exact Test, and/or concentrations significantly lowering fecundity based on Dunnett's test ($p = 0.05$, one tailed, $t_d = 2.12$).

For the data above, the mean number of offspring per female was computed for all surviving females. The least significant difference in fecundity at the 5% level (LSD_{0.05}) is 4.42 offspring per female, using the error MS value obtained with the GLM analysis (below). The LSD is a 21.5% reduction in fecundity compared to the control.

GLM Analysis of Ceriodaphnia Fecundity

Variance Source	D.F.	Sum of Squares	Mean Square	F	F _(0.05)
Among	3	2681.000	893.667	34.25	2.82
Within	44	1148.000	26.091		
Total	47	3829.000		(Pr > F = 0.0001)	

- 5.3 Statistical analyses: Control vs artificial rainwater (all concentrations).

The data were first analyzed by using Fisher's Exact Test (for survival), followed by SAS-GLM (General Linear Models) procedure and Dunnett's Test (for reproduction), as recommended by Horning and Weber (1985). The GLM procedure is recommended over ANOVA when the data set contains missing values (note table in section 5.1).

Asterisks in the summary table (section 5.2, above), show concentrations significantly reducing survivorship of Ceriodaphnia based on Fisher's Exact Test, and/or concentrations significantly lowering fecundity based on Dunnett's test ($p = 0.05$, one tailed, $L_d = 1.99$).

For the data set above, the mean number of offspring per female was computed for all surviving females. The least significant difference in fecundity at the 5% level ($LSD_{0.05}$) is 4.97 offspring per female, using the error MS value obtained with the GLM procedure (below). The LSD is a 24.1% reduction in fecundity relative to the control.

GLM Analysis of Ceriodaphnia Fecundity

Variance Source	D.F.	Sum of Squares	Mean Square	F	$F_{(0.05)}$
Among	2	56.771	28.385	0.76	3.32
Within	32	1199.515	37.484		
Total	34	1256.286		(Pr > F = 0.4772)	

- 5.3 Statistical analyses: Control vs 40% concentrations of the thermophilic leachate, the mesophilic leachate, and the artificial rainwater used to produce the two leachates.

The data were first analyzed by using Fisher's Exact Test (for survival), followed by SAS-GLM (General Linear Models) procedure and Dunnett's Test (for reproduction), as recommended by Horning and Weber (1985). The GLM procedure is recommended over ANOVA when the data set contains missing values (note table in section 5.1).

Asterisks in the summary table (section 5.2, above), show concentrations significantly reducing survivorship of Ceriodaphnia based on Fisher's Exact Test, and/or concentrations significantly lowering fecundity based on Dunnett's test ($p = 0.05$, one tailed, $L_d = 2.13$).

For the data above, the mean number of offspring per female was computed for all surviving females. The least significant difference in fecundity at the 5% level ($LSD_{0.05}$) is 6.06 offspring per female, using the error MS value from the GLM procedure (below). The LSD is a 29.4% reduction in fecundity relative to the control.

GLM Analysis of Ceriodaphnia Fecundity

Variance Source	D.F.	Sum of Squares	Mean Square	F	F _(0.05)
Among	3	1315.607	438.536	9.04	2.84
Within	42	2037.697	48.516		
Total	45	3353.304		(Pr > F = 0.0001)	

- 5.3 Statistical analyses: Control vs 60% concentrations of the thermophilic leachate, the mesophilic leachate, and the artificial rainwater used to produce the two leachates.

The data were first analyzed by using Fisher's Exact Test (for survival), followed by SAS-GLM (General Linear Models) procedure and Dunnett's Test (for reproduction), as recommended by Horning and Weber (1985). The GLM procedure is recommended over ANOVA when the data set contains missing values (note table in section 5.1).

Asterisks in the summary table (section 5.2, above), show concentrations significantly reducing survivorship of Ceriodaphnia based on Fisher's Exact Test, and/or concentrations significantly lowering fecundity based on Dunnett's test ($p = 0.05$, one tailed, $t_d = 2.12$).

For the data above, the mean number of offspring per female is computed for all surviving females. The least significant difference in fecundity at the 5% level ($LSD_{0.05}$) is 4.83 offspring per female, using the error MS value from the GLM procedure (below). The LSD is a 23.5% reduction in fecundity relative to the control.

GLM Analysis of Ceriodaphnia Fecundity

Variance Source	D.F.	Sum of Squares	Mean Square	F	F _(0.05)
Among	3	2882.917	960.972	30.84	2.82
Within	44	1371.000	31.159		
Total	47	4253.917		(Pr > F = 0.0001)	

5.4 Summary of Ceriodaphnia toxicity test results:

A direct comparison between leachates from the mesophilic and thermophilic composts is possible because the same concentration series and type of water was used in the tests. This comparison shows that the mesophilic leachate is about twice as toxic as the thermophilic leachate: Fecundity of Ceriodaphnia in the 40% concentration of the mesophilic leachate, for example, was similar to fecundity of the Ceriodaphnia in the 80% concentration of the thermophilic leachate (Summary Table, section 5.2).

The No-observed-effect concentration (NOEC) for the thermophilic leachate was 60% (relative to the control); the lowest-observed-effect concentration (LOEC) for this leachate was 80%.

The NOEC for the mesophilic leachate was <40% (relative to the control). Because 40% was the lowest concentration tested, the NOEC and LOEC for this leachate could not be more accurately determined. Because Ceriodaphnia fecundity was moderately high in the 40% concentration of the mesophilic leachate, it is reasonable to suppose that the "true" NOEC for this material would be about 15-25% of full-strength.

Based on visual appearance, both leachates contained very high concentrations of dissolved and/or colloidal organic matter. This material could have interfered mechanically with the Ceriodaphnia's capacity to filter food items from the water or adversely affected the "taste" of the items that they captured (cf. [1-3]). Either of these situations would have tended to increase the apparent toxicity of the leachates, for rate of food supply and reproduction are closely linked for Ceriodaphnia dubia.

6. CHEMICAL ANALYSES

6.1 Results of the daily chemical analyses:

Day	Concentration	pH	Cond. ^a	Alk. ^b	Hardness ^c	New D.O. ^d
1	Control	7.53	88	30.0	44	8.4
	T-100%	7.90	312	78.0	40	8.3
	T-40%	7.79	185	48.0	---	8.3
	M-100%	7.87	335	78.0	40	8.3
	M-40%	7.80	176	52.0	---	8.3
	AR-100%	5.84	4	1.5	≤2	8.3
	AR-40%	6.98	54	18.0	22	8.3
2	Control	7.78	87	30.0	38	8.2
	AR-40%	7.14	52	19.0	6	8.0
	T-100%	8.01	310	---	---	---
	T-80%	7.81	266	---	---	---
	T-60%	7.78	220	---	---	---
	T-40%	7.80	152	---	---	---
	M-100%	7.89	325	---	---	---
	M-80%	7.76	282	---	---	---
	M-60%	7.78	235	---	---	---
	M-40%	7.78	182	---	---	---
3	Control	7.26	86	34.0	38	8.4
	AR-40%	7.11	54	20.0	26	8.4
	T-100%	7.92	311	---	---	---
	T-80%	7.90	266	---	---	---
	T-60%	7.88	218	---	---	---
	T-40%	7.84	175	---	---	---
	M-100%	7.92	336	---	---	---
	M-80%	7.86	286	---	---	---
	M-60%	7.84	236	---	---	---
	M-40%	7.82	186	---	---	---
4	Control	7.66	87	29.0	40	8.2
	AR-40%	7.12	55	18.5	24	8.2
	T-100%	7.87	314	---	---	---
	T-80%	7.86	272	---	---	---
	T-60%	7.80	225	---	---	---
	T-40%	7.77	179	---	---	---
	M-100%	7.84	334	---	---	---
	M-80%	7.78	291	---	---	---
	M-60%	7.75	239	---	---	---
	M-40%	7.74	187	---	---	---

(continued)

6.1 Results of the daily chemical analyses (continued):

Day	Concentration	pH	Cond. ^a	Alk. ^b	Hardness ^c	New D.O. ^d
5	Control	7.63	83	29.0	42	8.2
	AR-40%	7.48	56	19.0	24	8.2
	T-100%	7.67	310	---	---	---
	T-80%	7.73	266	---	---	---
	T-60%	7.70	218	---	---	---
	T-40%	7.73	175	---	---	---
	M-100%	7.83	301	---	---	---
	M-80%	7.77	280	---	---	---
	M-60%	7.72	225	---	---	---
	M-40%	7.69	182	---	---	---
6	Control	7.92	87	30.0	44	8.4
	AR-40%	7.41	54	18.0	26	8.2
	T-100%	7.90	299	---	---	---
	T-80%	7.82	267	---	---	---
	T-60%	7.80	222	---	---	---
	T-40%	7.75	178	---	---	---
	M-100%	7.82	338	---	---	---
	M-80%	7.80	287	---	---	---
	M-60%	7.74	235	---	---	---
	M-40%	7.74	187	---	---	---
7	Control	8.12	87	29.0	42	8.3
	AR-40%	7.00	55	17.0	26	8.2
	T-100%	7.87	313	---	---	---
	T-80%	7.80	268	---	---	---
	T-60%	7.77	226	---	---	---
	T-40%	7.72	178	---	---	---
	M-100%	7.73	339	---	---	---
	M-80%	7.69	292	---	---	---
	M-60%	7.69	239	---	---	---
	M-40%	7.69	189	---	---	---

^aCond. = conductivity expressed as umho/cm, corrected to 25 °C.

^bAlk. = alkalinity expressed as mg/L CaCO₃.

^cHardness expressed as mg/L CaCO₃.

^dD.O. = mg/L dissolved oxygen.

6.2 Comments regarding chemical analysis data

The high concentrations of dissolved and/or colloidal organic matter seriously interfered with the colorimetric endpoint of the hardness assay. The accuracy of alkalinity measurements can be compromised by the presence of high concentrations of recalcitrant dissolved organic matter, as well [4]. Thus, the hardness and alkalinity values for in the table above may not be correct for the mesophilic and thermophilic leachates.

7. STANDARD REFERENCE TOXICANT TEST RESULTS

7.1 Date test conducted: June 14-15, 1989.

Time test initiated: 9:23 a.m.

7.2 Record of survival for 24-h test:

Concentration SLS (mg/L)	Replicate	Number at beginning	Number of dead
Control	1	6	0
	2	6	0
	3	6	0
	4	6	0
10 mg/L	1	6	1
	2	6	2
	3	6	1
	4	6	2
15 mg/L	1	6	3
	2	6	3
	3	6	3
	4	6	3
20 mg/L	1	6	4
	2	6	4
	3	6	4
	4	6	3
25 mg/L	1	6	4
	2	6	6
	3	6	6
	4	6	4
30 mg/L	1	6	6
	2	6	6
	3	6	6
	4	6	6

8. REFERENCES

- [1] DeMott, W. R. 1988. Discrimination between algae and artificial by freshwater and marine copepods. *Limnol. Oceanogr.* 33:397-408.
- [2] DeMott, W. R. 1986. The role of taste in food selection by freshwater zooplankton. *Oecologia* 69:334-340.
- [3] Gerritsen, J. and K. G. Porter. 1982. The role of surface chemistry in filter feeding by zooplankton. *Science* 216: 225-227.
- [4] Wilson, D. E. 1979. The influence of humic compounds on titrimetric determinations of total inorganic carbon in freshwater. *Arch. Hydrobiol.* 87:379-384.

Report prepared by: L. F. Wicker and A. J. Stewart

TOXICITY TEST REPORT

TOXICOLOGY LABORATORY
ENVIRONMENTAL SCIENCES DIVISION
OAK RIDGE NATIONAL LABORATORY
P.O. BOX 2008, MS-6351
OAK RIDGE, TN 37831-6351

EXPERIMENT NO. 340

Clean Closure Leachate Test
of
Thermophilic and Mesophilic Explosives-Contaminated Compost

June 22-29, 1989

STANDARD REPORT FORM

FATHEAD MINNOW LARVAL SURVIVAL AND GROWTH TEST

Experiment No. 340

Starting date: June 22, 1989.

Ending date: June 29, 1989.

1. SAMPLE

- 1.1 Sample description: Clean Closure Leachate Test leachates of thermophilic and mesophilic explosives-contaminated compost, and of artificial rainwater (AR) used to prepare the leachates.
- 1.2 Sampling point: Not applicable: Samples were collected from the laboratory where they were prepared.
- 1.3 Sampling date: Samples were picked up from the analytical chemistry laboratory on the morning of June 22, 1989, by A. J. Stewart.
- 1.4 Sampling method: Does not apply.
- 1.5 Sample was received at Environmental Sciences Division's toxicology laboratory on June 22, 1989 at 9:25 a.m. by A. J. Stewart; Registered Water Log Book A-103384, pp. 32.
- 1.6 Sample was used immediately.
- 1.7 Pretreatment: None.

2. TEST ORGANISMS

- 2.1 Species: Fathead minnow (Pimephales promelas).
- 2.2 Hatching date: June 21, 1989.
- 2.3 Incubation water: Dechlorinated tap water.
Incubation temperature: 25.0°C (mean \pm S).
- 2.4 Source: Environmental Sciences Division cultures.
- 2.5 Mean dry weight at start of test: 0.0937 \pm 0.0065 mg (mean \pm S)
- 2.6 Diseases and Treatment: none

3. TEST METHODS

- 3.1 Toxicity test method used: Fathead minnow larval survival and growth test.
Reference: EPA Test Method 1000.0, in W. B. Horning and C. I. Weber (eds.), Short-term Methods for Estimating the Chronic Toxicity of Effluents and Receiving Waters to Freshwater Organisms, EPA/600/4-85/014 (December 1985).
- 3.2 Deviations from reference: none.
- 3.3 Date test started: June 22, 1989.
Time test started: 1:10 p.m.
- 3.4 Date test terminated: June 29, 1989.
Time test terminated: 12:50 p.m.
- 3.5 Type of test chambers: Borosilicate 600-mL beakers.
Volume per chamber: 250 mL.
- 3.6 Number of organisms per test chamber: 10.
- 3.7 Number of replicates per treatment: 4.
- 3.8 Dilution/Control water: 1:9 (v:v) ratio of degassed mineral water to deionized distilled water.
- 3.9 Renewal period: 24 h.
- 3.10 Test temperature: Mean = 24.7°C; range = 24.6 - 24.8°C.
- 3.11 Treatment groups/concentrations: Control; Thermophilic 100%, 80%, 60%, 40%; Mesophilic 100%, 80%, 60%, 40%; and artificial rainwater 60%, and 40% of full-strength effluent.
- 3.12 Feeding Regime During Test: Brine shrimp (Artemia) nauplii less than 24 hours old; fed 600 ± 100 per beaker twice daily.

4. QUALITY ASSURANCE

- 4.1 Standard toxicant used: Sodium lauryl sulfate.

Source: Sigma Chemical Company.

4.2 Date of most recent test: June 14-15, 1989.

Time of most recent test: 9:49 a.m.

4.3 Dilution water used: 1:9 (v:v) ratio of degassed mineral water to deionized distilled water.

4.4 24-h LC_{50} : 23.053 mg/L SLS; 95% C.L. = 21.75 - 24.52 mg/L SLS.

6 The LC_{50} was calculated by the Moving Average method.

Reference: SAS User's Guide: Statistics, Version 5. Cary, NC: SAS Institute Inc., 1985, 956p.

Raw data for this reference test is provided in section 6.2.

4.5 Data from control chart prepared from standard toxicity tests:

Number of SLS standard tests completed by laboratory: 2.

Central Tendency: 25.7 ± 7.5 mg/L SLS (mean ± 2 S) mg/L.

4.6 Physical and chemical methods

The pH was measured by EPA method 150.1 with an Orion 700 pH meter. The meter was calibrated with pH 4 and pH 10 buffers.

Conductance ($\mu\text{mho/cm}$) was measured by EPA method 120.1 with a YSI model 31 meter. All values were corrected to 25°C.

Dissolved oxygen (mg/L) was measured by EPA method 360.1 with a YSI model 54AR dissolved oxygen meter. The meter was air calibrated.

Alkalinity was measured by titrating 50-mL samples with 0.1 N HCl to an endpoint pH of 4.5 (EPA method 130.1).

Hardness was determined by titrating 50-mL samples with EDTA to a colorimetric endpoint using Eriochrome Black T (EPA method 130.2).

Instruments were calibrated and standardized according to manufacturer's instructions.

All measurements were made on fresh samples before daily water replacement, except for oxygen (old), which was measured in the test solutions at the end of the replacement period.

5. FATHEAD MINNOW SURVIVAL TEST RESULTS

5.1 Daily results of the fathead minnow toxicity test:

Concentration	Replicate	Number of Larvae Surviving Each Day						
		1	2	3	4	5	6	7
Control	1	10	10	10	10	10	10	10
	2	9	9	9	9	9	9	9
	3	10	10	10	10	10	10	10
	4	10	10	10	10	10	10	10
T-100%	1	9	9	8	8	7	7	6
	2	9	9	8	7	7	6	6
	3	9	9	9	9	9	9	8
T-80%	1	10	10	9	9	9	9	9
	2	9	9	9	9	9	9	9
	3	10	10	10	10	9	9	9
T-60%	1	10	10	10	10	10	10	9
	2	10	10	10	9	9	9	9
	3	10	10	10	10	10	10	10
T-40%	1	10	10	10	10	10	10	10
	2	10	10	10	10	10	10	10
	3	10	10	10	10	10	10	10
M-100%	1	2	0	0	0	0	0	0
	2	3	0	0	0	0	0	0
	3	3	1	0	0	0	0	0
M-80%	1	8	5	2	0	0	0	0
	2	7	3	0	0	0	0	0
	3	6	5	1	0	0	0	0
M-60%	1	9	7	7	5	3	3	2
	2	10	10	7	4	3	3	3
	3	8	8	7	7	4	4	3
M-40%	1	10	9	9	9	9	9	7
	2	10	10	10	10	9	9	9
	3	10	9	8	8	8	7	6

(continued)

5.1 Daily results of the fathead minnow toxicity test (continued):

Concentration	Replicate	<u>Number of Larvae Surviving Each Day</u>						
		1	2	3	4	5	6	7
AR-60%	1	10	10	10	10	10	10	10
	2	10	10	10	10	10	10	10
	3	10	10	9	9	9	9	9
AR-40%	1	10	10	10	10	10	10	10
	2	10	10	10	10	10	10	10
	3	9	9	9	9	9	8	8

5.2 Summary of results from the fathead minnow toxicity test:

Survival (%)					
Sample Concentration	<u>Survival (%) per Replicate</u>				Mean
	1	2	3	4	
Control	100	90	100	100	97.5
T-100%	60	60	80	---	66.7
T- 80%	90	90	90	---	90.0
T- 60%	90	90	100	---	93.3
T- 40%	100	100	100	---	100.0
M-100%	0	0	0	---	0.0
M- 80%	0	0	0	---	0.0
M- 60%	20	30	30	---	26.7
M- 40%	70	90	60	---	73.3
AR-60%	100	100	90	---	96.7
AR-40%	100	100	80	---	93.3

Growth (Dry Weight)

Sample Concentration	Weight (mg) per Replicate				Mean \pm \bar{S}
	1	2	3	4	
Control	0.34	0.39	0.36	0.36	0.36 \pm 0.021
T-100%	0.53	0.57	0.46	---	0.52 \pm 0.051
T- 80%	0.41	0.36	0.46	---	0.41 \pm 0.049
T- 60%	0.39	0.47	0.39	---	0.41 \pm 0.042
T- 40%	0.48	0.43	0.49	---	0.47 \pm 0.027
M-100%	---	---	---	---	---
M- 80%	---	---	---	---	---
M- 60%	0.39	0.26	0.23	---	0.30 \pm 0.081
M- 40%	0.36	0.40	0.44	---	0.40 \pm 0.039
AR-60%	0.43	0.50	0.44	---	0.45 \pm 0.036
AR-40%	0.35	0.51	0.38	---	0.41 \pm 0.082

5.3 Statistical analyses of survival data: Control vs thermophilic leachate.

The data were analyzed using the SAS-GLM (General Linear Models) procedure and Dunnett's Test, as recommended by Horning and Weber (1985). The GLM procedure is recommended over ANOVA when the data set contains missing values (note table in section 5.1). Arcsin transformation of the data was used before analysis of survival. For this set of data, the least significant difference is 12.6 (arcsin transformed). This represents a 14.8% reduction in survival. $t_d = 2.44$, $p = 0.05$ (one-tailed test). Asterisks in the summary table (section 5.2, above), show concentrations significantly reducing survival of fathead minnows based on Dunnett's Test.

GLM Analysis of Fathead Minnow Survival

Source	D.F.	Sum of Squares	Mean Square	\bar{F}	$F_{(0.05)}$
Among	4	2328.390	582.097	10.88	3.36
Within	11	588.414	53.492		
Total	15	2916.804		(Pr > \bar{F} = 0.0008)	

5.4 Statistical Analyses of Dry Weight Data: Control vs thermophilic leachate.

The data were analyzed by using the SAS-GLM (General Linear Models) procedure and Dunnett's Test, as recommended by Horning and Weber (1985). The GLM procedure is recommended over ANOVA when the data set contains missing values (note table in section 5.1).

No transformation was used before data analysis. For this set of data, the least significant difference is 0.071 mg. This represents a 19.8% reduction in dry weight. $t_d = 2.44$, $p = 0.05$ (one-tailed test). The mean dry weight of larvae at the start of the test was 0.094 mg/fish.

GLM Analysis of Fathead Minnow Dry Weight

Source	D.F.	Sum of Squares	Mean Square	F	$F_{(0.05)}$
Among	4	0.0483	0.0121	7.06	3.36
Within	11	0.0188	0.0017		
Total	15	0.0671		(Pr > F = 0.0045)	

5.3 Statistical analyses of survival data: Control vs mesophilic leachate.

The data were analyzed using the SAS-GLM (General Linear Models) procedure and Dunnett's Test, as recommended by Horning and Weber (1985). The GLM procedure is recommended over ANOVA when the data set contains missing values (note table in section 5.1).

Arcsin transformation of the data was used before analysis of survival. For this set of data, the least significant difference is 11.6 (arcsin transformed). This represents a 13.6% reduction in survival. $t_d = 2.37$, $p = 0.05$ (one-tailed test). Asterisks in the summary table (section 5.2, above), show concentrations significantly reducing survival of fathead minnows based on Dunnett's Test.

GLM Analysis of Fathead Minnow Survival

Source	D.F.	Sum of Squares	Mean Square	F	F _(0.05)
Among	3	14405.868	4801.956	100.53	4.35
Within	9	429.886	47.765		
Total	12	14835.754		(Pr > F = 0.0001)	

5.4 Statistical Analyses of Dry Weight Data: Control vs mesophilic leachate.

The data were analyzed by using the SAS-GLM (General Linear Models) procedure and Dunnett's Test, as recommended by Horning and Weber (1985). The GLM procedure is recommended over ANOVA when the data set contains missing values (note table in section 5.1).

No transformation was used before data analysis. For this set of data, the least significant difference is 0.083 mg. This represents a 23.2% reduction in dry weight. $t_d = 2.27$, $p = 0.05$ (one-tailed test). The mean dry weight of larvae at the start of the test was 0.094 mg/fish.

GLM Analysis of Fathead Minnow Dry Weight

Source	D.F.	Sum of Squares	Mean Square	F	F _(0.05)
Among	2	0.0177	0.0088	3.26	4.74
Within	7	0.0189	0.0027		
Total	9	0.0366		(Pr > F = 0.0996)	

5.3 Statistical analyses of survival data: Control vs artificial rainwater.

The data were analyzed using the SAS-GLM (General Linear Models) procedure and Dunnett's Test (Horning and Weber, 1985). The GLM procedure is recommended over ANOVA when the data set contains missing values (note table in section 5.1).

Arcsin transformation of the data was used before analysis of survival. For this set of data, the least significant difference is 18.7 (arcsin transformed). This represents a 21.9% reduction in survival. $t_d = 2.27$, $p = 0.05$ (one-tailed test). Asterisks in the summary table (section 5.2, above), show concentrations significantly reducing survival of fathead minnows based on Dunnett's Test.

GLM Analysis of Fathead Minnow Survival

Source	D.F.	Sum of Squares	Mean Square	F	$F_{(0.05)}$
Among	2	31.079	15.539	0.11	4.74
Within	7	951.926	135.989		
Total	9	983.005		(Pr > F = 0.8936)	

5.4 Statistical Analyses of Dry Weight Data: Control vs artificial rainwater.

The data were analyzed by using the SAS-GLM (General Linear Models) procedure and Dunnett's Test, as recommended by Horning and Weber (1985). The GLM procedure is recommended over ANOVA when the data set contains missing values (note table in section 5.1).

No transformation was used before data analysis. For this set of data, the least significant difference is 0.083 mg. This represents a 23.2% reduction in dry weight. $t_d = 2.27$, $p = 0.05$ (one-tailed test). The mean dry weight of larvae at the start of the test was 0.094 mg/fish.

GLM Analysis of Fathead Minnow Dry Weight

Source	D.F.	Sum of Squares	Mean Square	F	$F_{(0.05)}$
Among	2	0.0154	0.0077	2.90	4.74
Within	7	0.0186	0.0026		
Total	9	0.0340		(Pr > F = 0.1208)	

- 5.3 Statistical analyses of survival data: Control vs 40% concentrations of the thermophilic leachate, the mesophilic leachate, and the artificial rainwater used to produce the two leachates.

The data were analyzed using the SAS-GLM (General Linear Models) procedure and Dunnett's Test, as recommended by Horning and Weber (1985). The GLM procedure is recommended over ANOVA when the data set contains missing values (note table in section 5.1).

Arcsin transformation of the data was used before analysis of survival. For this set of data, the least significant difference is 17.3 (arcsin transformed). This represents a 20.2% reduction survival. $t_d = 2.37$, $p = 0.05$ (one-tailed test). Asterisks in the summary table (section 5.2, above), show concentrations significantly reducing survival of fathead minnows based on Dunnett's Test.

GLM Analysis of Fathead Minnow Survival

Source	D.F.	Sum of Squares	Mean Square	\bar{F}	$\bar{F}_{(0.05)}$
Among	3	1652.903	550.968	5.20	3.86
Within	9	954.386	106.043		
Total	12	2607.289		$(Pr > \bar{F} = 0.0235)$	

- 5.4 Statistical Analyses of Dry Weight Data: Control vs 40% concentrations of the thermophilic leachate, the mesophilic leachate, and the artificial rainwater used to prepare the two leachates.

The data were analyzed by using the SAS-GLM (General Linear Models) procedure and Dunnett's Test, as recommended by Horning and Weber (1985). The GLM procedure is recommended over ANOVA when the data set contains missing values (note table in section 5.1).

No transformation was used before data analysis. For this set of data, the least significant difference is 0.080 mg. This represents a 22.3% reduction in dry weight. $t_d = 2.37$, $p = 0.05$ (one-tailed test). The mean dry weight of larvae at the start of the test was 0.094 mg/fish.

GLM Analysis of Fathead Minnow Dry Weight

Source	D.F.	Sum of Squares	Mean Square	F	$F_{(0.05)}$
Among	3	0.0189	0.0063	2.69	3.86
Within	9	0.0210	0.0023		
Total	12	0.0399		(Pr > F = 0.1089)	

- 5.3 Statistical analyses of survival data: Control vs 60% concentrations of the thermophilic leachate, the mesophilic leachate, and the artificial rainwater used to produce the two leachates.

The data were analyzed using the SAS-GLM (General Linear Models) procedure and Dunnett's Test, as recommended by Horning and Weber (1985). The GLM procedure is recommended over ANOVA when the data set contains missing values (note table in section 5.1).

Arcsin transformation of the data was used before analysis of survival. For this set of data, the least significant difference is 15.2 (arcsin transformed). This represents a 17.8% reduction in survival. $t_d = 2.37$, $p = 0.05$ (one-tailed test). Asterisks in the summary table (section 5.2, above), show concentrations significantly reducing survival of fathead minnows based on Dunnett's Test.

GLM Analysis of Fathead Minnow Survival

Source	D.F.	Sum of Squares	Mean Square	F	$F_{(0.05)}$
Among	3	6259.277	2086.426	25.46	3.86
Within	9	737.466	81.941		
Total	12	6996.743		(Pr > F = 0.0001)	

- 5.4 Statistical Analyses of Dry Weight Data: Control vs 60% concentrations of the thermophilic leachate, the mesophilic leachate, and the artificial rainwater used to produce the two leachates.

The data were analyzed by using the SAS-GLM (General Linear Models) procedure and Dunnett's Test (Hornig and Weber, 1985). The GLM procedure is recommended over ANOVA when the data set contains missing values (note table in section 5.1).

No transformation was used before data analysis. For this set of data, the least significant difference is 0.084 mg. This represents a 23.3% reduction in dry weight. $t_d = 2.37$, $p = 0.05$ (one-tailed test). The mean dry weight of larvae at the start of the test was 0.094 mg/fish.

GLM Analysis of Fathead Minnow Dry Weight

Source	D.F.	Sum of Squares	Mean Square	F	$F_{(0.05)}$
Among	3	0.0454	0.0151	5.96	3.86
Within	9	0.0229	0.0025		
Total	12	0.0683		(Pr > F = 0.0160)	

5.5 Summary of toxicity test results:

No-observed-effect concentration (NOEC):

80% concentration for thermophilic leachate, <40% for mesophilic leachate, based on survival of the fish relative to controls.

Lowest-observed-effect-concentration (LOEC):

100% concentration for thermophilic leachate, 40% for mesophilic leachate, based on survival of the fish relative to controls.

48-h LC_{50} concentration (by graphic interpolation):

>100% concentration for thermophilic leachate, 77% for mesophilic leachate, >100% for artificial rainwater.

7-d median lethal concentration (LC_{50}) (by graphic interpolation):

>100% for thermophilic leachate, 50% for mesophilic leachate, >100% for artificial rainwater.

7-d effective concentration (EC_{50}): Not determined.

6. CHEMICAL ANALYSES

6.1 Results of the daily chemical analyses:

Day	Concentration	pH	Cond. ^a	Alk. ^b	Hardness ^c	Oxygen ^d	
						New	Old
1	Control	7.53	88	30.0	44	8.4	7.8
	T-100%	7.90	312	78.0	40	8.3	7.6
	T-40%	7.79	185	48.0	---	8.3	7.5
	M-100%	7.87	335	78.0	40	8.3	7.6
	M-40%	7.80	176	52.0	---	8.3	7.6
	AR-100%	5.84	4	1.5	≤2	8.3	---
	AR-40%	6.98	54	18.0	22	8.3	7.6
2	Control	7.78	87	30.0	38	8.2	8.0
	AR-40%	7.14	52	19.0	6	8.0	7.2
	T-100%	8.01	310	---	---	---	7.6
	T-80%	7.81	266	---	---	---	---
	T-60%	7.78	220	---	---	---	---
	T-40%	7.80	152	---	---	---	7.4
	M-100%	7.89	325	---	---	---	7.3
	M-80%	7.76	282	---	---	---	---
	M-60%	7.78	235	---	---	---	---
	M-40%	7.78	182	---	---	---	7.2
3	Control	7.26	86	34.0	38	8.4	7.6
	AR-40%	7.11	54	20.0	26	8.4	6.7
	T-100%	7.92	311	---	---	---	6.8
	T-80%	7.90	266	---	---	---	---
	T-60%	7.88	218	---	---	---	---
	T-40%	7.84	175	---	---	---	6.8
	M-100%	7.92	336	---	---	---	6.8
	M-80%	7.86	286	---	---	---	---
	M-60%	7.84	236	---	---	---	---
	M-40%	7.82	186	---	---	---	6.8
4	Control	7.66	87	29.0	40	8.2	7.5
	AR-40%	7.12	55	18.5	24	8.2	7.4
	T-100%	7.87	314	---	---	---	7.1
	T-80%	7.86	272	---	---	---	---
	T-60%	7.80	225	---	---	---	---
	T-40%	7.77	179	---	---	---	7.1
	M-100%	7.84	334	---	---	---	---
	M-80%	7.78	291	---	---	---	---
	M-60%	7.75	239	---	---	---	---
	M-40%	7.74	187	---	---	---	7.5

(continued)

6.1 Results of the daily chemical analyses (continued):

Day	Concentration	pH	Cond. ^a	Alk. ^b	Hardness ^c	Oxygen ^d	
						New	Old
5	Control	7.63	83	29.0	42	8.2	7.6
	AR-40%	7.48	56	19.0	24	8.2	7.5
	T-100%	7.67	310	---	---	---	7.7
	T-80%	7.73	266	---	---	---	---
	T-60%	7.70	218	---	---	---	---
	T-40%	7.73	175	---	---	---	7.7
	M-100%	7.83	301	---	---	---	---
	M-80%	7.77	280	---	---	---	---
	M-60%	7.72	225	---	---	---	---
	M-40%	7.69	182	---	---	---	7.7
6	Control	7.92	87	30.0	44	8.4	7.7
	AR-40%	7.41	54	18.0	26	8.2	7.3
	T-100%	7.90	299	---	---	---	7.1
	T-80%	7.82	267	---	---	---	---
	T-60%	7.80	222	---	---	---	---
	T-40%	7.75	178	---	---	---	---
	M-100%	7.82	338	---	---	---	---
	M-80%	7.80	287	---	---	---	---
	M-60%	7.74	235	---	---	---	---
	M-40%	7.7	187	---	---	---	7.3
7	Control	8.12	87	29.0	42	8.3	7.7
	AR-40%	7.00	55	17.0	26	8.2	7.4
	T-100%	7.87	313	---	---	---	7.4
	T-80%	7.80	268	---	---	---	---
	T-60%	7.77	226	---	---	---	---
	T-40%	7.72	178	---	---	---	7.4
	M-100%	7.73	339	---	---	---	---
	M-80%	7.69	292	---	---	---	---
	M-60%	7.69	239	---	---	---	---
	M-40%	7.69	189	---	---	---	7.4

^aCond. = conductivity expressed as umho/cm, corrected to 25 °C. ^bAlk. = alkalinity expressed as mg/L CaCO₃. ^cHardness expressed as mg/L CaCO₃. ^dD.O. = mg/L dissolved oxygen.

7. STANDARD REFERENCE TOXICANT TEST RESULTS

7.1 Date test conducted: June 14-15, 1989.

Time test initiated: 9:49 a.m.

7.2 Record of survival for 24-h test (sodium lauryl sulfate):

Concentration SLS (mg/L)	Rep.	Number at Beginning	Number of Dead
Control	1	6	0
	2	6	0
	3	6	0
	4	6	0
15	1	6	0
	2	6	0
	3	6	0
	4	6	0
20	1	6	1
	2	6	1
	3	6	1
	4	6	0
25	1	6	5
	2	6	4
	3	6	4
	4	6	4
30	1	6	6
	2	6	6
	3	6	6
	4	6	5
40	1	6	6
	2	6	6
	3	6	6
	4	6	6

Report prepared by: L. F. Wicker and A. J. Stewart

APPENDIX A-6
AMES TEST DATA

TABLE 6A-1. AMES TEST OF TNT AND TETRYL

	$\mu\text{g/plate}$	<u>TA 98</u> Revertants/plate	<u>TA 100</u> Revertants/plate
Control ^a	-	15	68
Nitrofluorene	5	536	-
Sodium azide	1	-	603
	20	98	238
	50	222	544
TNT	100	290	921
	150	343	1266
	200 ^b	381	1318
	2	30	142
	5	47	178
Tetryl	10	89	305
	15 ^b	146	419
	20 ^b	175	722

^aDMSO solvent.^bToxicity toward Salmonella tester strain observed.

TABLE 6A-2. AMES TEST OF TNT, 2-4-DA,6-NT AND 2-6-DA,4-NT

Sample	$\mu\text{g/plate}$	Revertants/Plate	
		TA98	TA100
Control*	60	6	117
Nitrofluorene	5	551	NT
Sodium Azide	0.5	NT	506
	20	88	232
TNT	40	172	366
	80	290	577
	120	399	932
	50	15	134
2,4-DA-6-NT	100	16	131
	150	18	120
	200	15	131
	250	22	148
	50	26	173
2,6-DA-4-NT	100	49	178
	150	56	228
	200	72	241
	250	83	247

*DMSO

NT = Not tested

TABLE 6A-3. AMES TEST OF TNT, 2-A-4,6-DNT AND 6-A-2,6-DNT

Sample	$\mu\text{g/plate}$	Revertants/Plate	
		TA98	TA100
Control ^a	100	22	98
Nitrofluorene	5	590	NT
Sodium azide	0.5	NT	506
TNT	20	155	209
	40	303	371
	80	501	777
	120	653	890
	160	679	1112
	200	644	774
2-A-4,6-DNT	20	27	141
	40	30	169
	80	63	279
	120	71	334
	160	83	495
	200	129	545
4-A-2,6-DNT	20	40	106
	40	60	131
	80	103	138
	120	138	138
	160	161	176
	200	202	198

^aAcetonitrile solvent

NT = Not tested

TABLE 6A-4. RESULTS FROM AMES TESTING OF CONTROLS AND PRELIMINARY CCLT LEACHATES

Sample	TA98 ^c revertants/plate		TA100 ^c revertants/plate	
	-S9	+S9	-S9	+S9
1. Control	24	28	117	111
2. Nitrofluorene ^a	874	NT	NT	NT
3. Acetylaminofluorene ^a	NT	533	NT	227
4. Sodium Azide ^a	NT ^b	NT	586	NT
5. 50 μ L Mesophilic Leachate	26	24	109	109
6. 200 μ L Mesophilic Leachate	29	35	113	107
7. 350 μ L Mesophilic Leachate	45	NT	108	NT
8. 700 μ L Mesophilic Leachate	56	NT	97	NT
9. 50 μ L Thermophilic Leachate	19	29	101	103
10. 200 μ L Thermophilic Leachate	30	33	95	106
11. 350 μ L Thermophilic Leachate	32	NT	111	NT
12. 700 μ L Thermophilic Leachate	35	NT	NT	NT

^aPositive controls. Nitrofluorene (10 μ g/plate). Acetylaminofluorene (10 μ g/plate). Sodium azide (2 μ g/plate).

^bNT - not tested

^cSalmonella overnight culture - 2×10^9 cells/ml. Ampicillin resistant, UV sensitive, crystal violet sensitive.

TABLE 6A-5. AMES TEST OF CCLT LEACHATES OF THE MESOPHILIC
AND THERMOPHILIC COMPOSTS

	$\mu\text{g/plate}$	<u>TA 98</u> Revertants/plate	<u>TA 100</u> Revertants/plate
Control*	1800	30	147
Nitrofluorene	5 ^b	348	NT
Sodium azide	0.5 ^b	NT	262
Mesophilic	200	44	141
Leachate	600	35	150
	1000	86	174
	1400	105	202
	1800	95	152
Thermophilic	200	26	128
Leachate	600	39	116
	1000	33	177
	1400	38	187
	1800	41	146

*CCLT solvent.

^bNumber of $\mu\text{g/plate}$.

NT = Not tested.

TABLE 6A-6. AMES TEST OF THE DMSO CONCENTRATE OF THE MESOPHILIC AND THERMOPHILIC COMPOST CCLT EXTRACTS

Sample	$\mu\text{g}/\text{plate}$	Revertants/Plate	
		TA98	TA100
Control ^a	50	34	74
Nitrofluorene	5 ^b	722	NT
Sodium azide	0.5 ^b	NT	480
Mesophilic	10	623	219
	20	1106	344
	30	2160	504
	40	2880	703
	50	3552	607
Thermophilic	10	185	126
	20	340	172
	30	540	201
	40	678	252
	50	862	278

^aDMSO solvent

^b $\mu\text{g}/\text{plate}$

NT = Not tested

TABLE 6A-7. AMES TEST OF THE DMSO CONCENTRATE OF THE MESOPHILIC AND THERMOPHILIC COMPOST CCLT EXTRACT

Sample	μ L	μ g	<u>Revertants</u>	
			TA-98	TA-100
Spontaneous	-	-	27	119
DMSO Control	320	-	23	75
Sodium Azide	-	0.5	-	520
Nitrofluorene	-	5	709	-
	20	-	47	129
Mesophilic	40	-	51	127
Compost	80	-	60	157
Leachate	160	-	101	184
	320	-	169	196
	20	-	55	124
DMSO	40	-	79	143
Extract of	80	-	173	169
Particles	160	-	305	195
	320	-	496	299
Spontaneous	-	-	24	97
DMSO Control	320	-	19	89
Sodium Azide	-	0.5	-	550
Nitro fluorene	-	5	920	-
Centrifuged	20	-	16	117
Mesophilic	40	-	15	115
Compost	80	-	33	123
Leachate	160	-	34	130
	320	-	49	136

TABLE 6A-8. AMES TEST OF PRELIMINARY ACETONITRILE EXTRACTS OF
MESOPHILIC AND THERMOPHILIC COMPOSTS

Sample	$\mu\text{g/plate}$	Revertants/Plate	
		TA98	TA100
Control ^a	100	12	88
Nitrofluorene	5	308	NT
Sodium azide	0.5	NT	342
Mesophilic	10	40	107
	100	351	230
	200	441	340
Thermophilic	10	41	95
	100	266	195
	200	327	275

^aAcetonitrile solvent

NT = Not tested

TABLE 6A-9. AMES TEST OF FINAL ACETONITRILE EXTRACTS OF MESOPHILIC AND THERMOPHILIC COMPOSTS

Sample	$\mu\text{g/plate}$	Revertants/Plate	
		TA98	TA100
Control*	80	11	106
Nitrofluorene	5	911	NT
Sodium Azide	0.5	NT	575
Mesophilic	40	418	232
	80	723	311
	120	1084	364
	160	1227	450
Thermophilic	40	1458	197
	80	2190	296
	120	2976	363
	160	2780	450

*DMSO

NT = Not tested

TABLE 6A-10. AMES TEST OF FINAL ETHYL ACETATE EXTRACTS OF MESOPHILIC AND THERMOPHILIC COMPOSTS

Sample	$\mu\text{g/plate}$	Revertants/Plate	
		TA98	TA100
Control ^a	80	11	106
Nitrofluorene	5	911	NT
Sodium Azide	0.5	NT	575
Mesophilic	40	464	209
	80	781	286
	120	984	345
	160	1266	364
Thermophilic	40	3408	357
	80	TNTC	594
	120	TNTC	878
	160	TNTC	982

DMSO

NT = Not tested

TNTC = Too numerous to count, i.e. greater than 4,000.

TABLE 6A-11. AMES TEST OF ORGANIC SOLVENT EXTRACTS: SECOND SET 12/89

Sample/Solvent	μL	μg	Revertants	
			TA-98	TA-100
Spontaneous	-	-	27	98
Sodium Azide	5	0.5	-	551
Nitrofluorene	5	5	962	-
	10	-	616	592
	20	-	1258	1198
Mesophilic	40	-	1972	1830
Acetonitrile	80	-	2112	1672
	160	-	T*	851*
	10	-	626	476
Mesophilic	20	-	1108	668
Ethylacetate	40	-	1831	1438
	80	-	2201	1600
	160	-	3949	2368
	10	-	1874	369
Thermophilic	20	-	2894	640
Acetonitrile	40	-	3528	1067
	80	-	4060	1544
	160	-	7056	2352
	10	-	4364	721
Thermophilic	20	-	4177	1304
Ethylacetate	40	-	5820	1756
	80	-	TNTC	2480
	160	-	TNTC	2832

*Toxic

TNTC > 6000

TABLE 6A-12. AMES TEST OF ACID HYDROLYSATES OF BOUND FRACTION
RESULTING FROM MESOPHILIC AND THERMOPHILIC COMPOSTS

Sample	$\mu\text{g/plate}$	Revertants/Plate	
		TA98	TA100
Control ^a	NA	33	109
Nitrofluorene	5	894	1180
Sodium Azide	0.5	27	538
Mesophilic	360	42	113
	720	49	128
	1440	61	121
	2880	53	138
	5760	36	147
Thermophilic	292	38	123
	584	49	127
	1168	42	151
	2336	45	147
	4672	46	142
Blank ^b	32	165	133

^aSpontaneous revertants, not treated
NA = Not applicable

^bAcid hydrolysate reagents in the absence of compost
NT = Not tested

DISTRIBUTION LIST

	<u>No. of Copies</u>
Commander U.S. Army Biomedical Research and Development Laboratory ATTN: SGRD-UBZ-RA Fort Detrick Frederick, MD 21701-5010	20
Commander U.S. Army Medical Research and Development Command ATTN: SGRD-RMI-S Fort Detrick Frederick, MD 21701-5012	2
Central Resarch Library Bldg. 4500N, MS-6191 Oak Ridge National Laboratory P.O. Box 2008 Oak Ridge, TN 37831-6286	1
Document Reference Section Bldg. 9711-1, MS-6107 Oak Ridge National Laboratory P.O. Box 2008 Oak Ridge, TN 37831-6286	1
Mr. R. L. Egli, Acting Assistant Manager Energy Research and Development U.S. Department of Energy, Oak Ridge Operations P.O. Box E Oak Ridge, TN 37831-8600	1
W. H. Griest Bldg. 4500S, MS-6120 Oak Ridge National Laboratory P.O. Box 2008 Oak Ridge, TN 37831-6120	10
M. R. Guerin Bldg. 4500S, MS-6120 Oak Ridge National Laboratory P.O. Box 2008 Oak Ridge, TN 37831-6120	10
C.-h. Ho Bldg. 4500S, MS-6120 Oak Ridge National Laboratory P. O. Box 2008 Oak Ridge, TN 37831-6120	1

DISTRIBUTION LIST (CONTD)

	<u>No. of Copies</u>
Laboratory Records Bldg. 4500N, MS-6285 Oak Ridge National Laboratory P.O. Box 2008 Oak Ridge, TN 37831-6285	3
C. P. McGinnis Bldg. K1006, MS-7274 ORGDP Oak Ridge, TN 37831-7274	1
ORNL Patent Office Bldg. 4500N, MS-6258 Oak Ridge National Laboratory P.O. Box 2008 Oak Ridge, TN 37831-6258	1
A. J. Stewart Bldg. 1504, MS-6351 Oak Ridge National Laboratory P. O. Box 2008 Oak Ridge, TN 37831-6351	1
E. Tan Oak Ridge Associated Universities Medical P. O. Box 117 Oak Ridge, TN 37831-0117	1
R. L. Tyndall Bldg. 9207, Y-12 Oak Ridge National Laboratory P. O. Box 2008 Oak Ridge, TN 37831-8077	1
U.S. Department of Energy Office of Scientific and Technical Information P.O. Box 62 Oak Ridge, TN 37831	10